

THE STABLE DERIVED CATEGORY OF A NOETHERIAN SCHEME

HENNING KRAUSE

ABSTRACT. For a noetherian scheme, we introduce its unbounded stable derived category. This leads to a recollement which reflects the passage from the bounded derived category of coherent sheaves to the quotient modulo the subcategory of perfect complexes. Some applications are included, for instance an analogue of maximal Cohen-Macaulay approximations, a construction of Tate cohomology, and an extension of the classical Grothendieck duality.

1. INTRODUCTION

Let \mathbb{X} be a separated noetherian scheme and denote by $\mathbf{Qcoh}\ \mathbb{X}$ the category of quasi-coherent sheaves on \mathbb{X} . We consider the derived category $\mathbf{D}(\mathbf{Qcoh}\ \mathbb{X})$ and two full subcategories

$$\mathbf{D}^{\mathrm{perf}}(\mathrm{coh}\ \mathbb{X}) \subseteq \mathbf{D}^b(\mathrm{coh}\ \mathbb{X}) \subseteq \mathbf{D}(\mathbf{Qcoh}\ \mathbb{X})$$

which are of particular interest. Here, $\mathbf{D}^b(\mathrm{coh}\ \mathbb{X})$ denotes the bounded derived category of coherent sheaves, and $\mathbf{D}^{\mathrm{perf}}(\mathrm{coh}\ \mathbb{X})$ denotes the subcategory of perfect complexes.

Now let $\mathbf{Inj}\ \mathbb{X}$ be the full subcategory of injective objects in $\mathbf{Qcoh}\ \mathbb{X}$, and denote by $\mathbf{K}(\mathbf{Inj}\ \mathbb{X})$ its homotopy category. The composite

$$Q: \mathbf{K}(\mathbf{Inj}\ \mathbb{X}) \xrightarrow{\mathrm{inc}} \mathbf{K}(\mathbf{Qcoh}\ \mathbb{X}) \xrightarrow{\mathrm{can}} \mathbf{D}(\mathbf{Qcoh}\ \mathbb{X})$$

gives rise to a localization sequence

$$\mathbf{S}(\mathbf{Qcoh}\ \mathbb{X}) \xrightarrow{I} \mathbf{K}(\mathbf{Inj}\ \mathbb{X}) \xrightarrow{Q} \mathbf{D}(\mathbf{Qcoh}\ \mathbb{X})$$

where $\mathbf{S}(\mathbf{Qcoh}\ \mathbb{X})$ denotes the full subcategory of all acyclic complexes in $\mathbf{K}(\mathbf{Inj}\ \mathbb{X})$. Thus Q induces an equivalence

$$\mathbf{K}(\mathbf{Inj}\ \mathbb{X})/\mathbf{S}(\mathbf{Qcoh}\ \mathbb{X}) \xrightarrow{\sim} \mathbf{D}(\mathbf{Qcoh}\ \mathbb{X}).$$

Next we recall that an object X in some category with coproducts is *compact* if every map $X \rightarrow \coprod_i Y_i$ into an arbitrary coproduct factors through a finite coproduct. For instance, an object in $\mathbf{D}(\mathbf{Qcoh}\ \mathbb{X})$ is compact if and only if it is isomorphic to a perfect complex. It is well known that the derived category $\mathbf{D}(\mathbf{Qcoh}\ \mathbb{X})$ is *compactly generated*, that is, there is a set of compact objects which generate $\mathbf{D}(\mathbf{Qcoh}\ \mathbb{X})$ [26]. To formulate our main result, let us denote by $\mathbf{K}^c(\mathbf{Inj}\ \mathbb{X})$ and $\mathbf{S}^c(\mathbf{Qcoh}\ \mathbb{X})$ the full subcategories of compact objects in $\mathbf{K}(\mathbf{Inj}\ \mathbb{X})$ and $\mathbf{S}(\mathbf{Qcoh}\ \mathbb{X})$ respectively.

Theorem 1.1. *Let \mathbb{X} be a separated noetherian scheme.*

2000 *Mathematics Subject Classification.* Primary: 14F05, 18E30; Secondary: 16E45, 16G50, 55U35.
Version from March 2004.

- (1) The functors I, Q have left adjoints I_λ, Q_λ and right adjoints I_ρ, Q_ρ respectively. We have therefore a recollement

$$\mathbf{S}(\mathrm{Qcoh}\ \mathbb{X}) \rightleftarrows \mathbf{K}(\mathrm{Inj}\ \mathbb{X}) \rightleftarrows \mathbf{D}(\mathrm{Qcoh}\ \mathbb{X}).$$

- (2) The triangulated category $\mathbf{K}(\mathrm{Inj}\ \mathbb{X})$ is compactly generated, and Q induces an equivalence $\mathbf{K}^c(\mathrm{Inj}\ \mathbb{X}) \rightarrow \mathbf{D}^b(\mathrm{coh}\ \mathbb{X})$.
 (3) The sequence

$$\mathbf{D}(\mathrm{Qcoh}\ \mathbb{X}) \xrightarrow{Q_\lambda} \mathbf{K}(\mathrm{Inj}\ \mathbb{X}) \xrightarrow{I_\lambda} \mathbf{S}(\mathrm{Qcoh}\ \mathbb{X})$$

is a localization sequence. Therefore $\mathbf{S}(\mathrm{Qcoh}\ \mathbb{X})$ is compactly generated, and $I_\lambda \circ Q_\rho$ induces (up to direct factors) an equivalence

$$\mathbf{D}^b(\mathrm{coh}\ \mathbb{X}) / \mathbf{D}^{\mathrm{perf}}(\mathrm{coh}\ \mathbb{X}) \xrightarrow{\sim} \mathbf{S}^c(\mathrm{Qcoh}\ \mathbb{X}).$$

Note that this theorem is a special case of a general result about Grothendieck categories. All we need is a locally noetherian Grothendieck category \mathcal{A} , for instance $\mathcal{A} = \mathrm{Qcoh}\ \mathbb{X}$, such that $\mathbf{D}(\mathcal{A})$ is compactly generated. There is a surprising consequence which seems worth mentioning.

Corollary 1.2. *Let \mathbb{X} be a separated noetherian scheme. Then a product of acyclic complexes of injective objects in $\mathrm{Qcoh}\ \mathbb{X}$ is acyclic.*

We call the category $\mathbf{S}(\mathrm{Qcoh}\ \mathbb{X})$ the *stable derived category* of $\mathrm{Qcoh}\ \mathbb{X}$. A first systematic study of the bounded stable derived category

$$\mathbf{D}^b(\mathrm{coh}\ \mathbb{X}) / \mathbf{D}^{\mathrm{perf}}(\mathrm{coh}\ \mathbb{X})$$

can be found in work of Buchweitz [12]. Unfortunately this beautiful paper has never been published; see however [11]. He identifies for a Gorenstein ring Λ the bounded derived category of finitely generated Λ -modules modulo perfect complexes

$$\mathbf{D}^b(\mathrm{mod}\ \Lambda) / \mathbf{D}^{\mathrm{perf}}(\mathrm{mod}\ \Lambda)$$

with the stable category of maximal Cohen-Macaulay Λ -modules and with the category of acyclic complexes of finitely generated projective Λ -modules. The same identification appears in [30] for selfinjective algebras and plays an important role in modular representation theory of finite groups; see also [23]. The approach in the present paper differs from that of Buchweitz substantially because we work in the unbounded setting and we use injective objects instead of projectives. This has some advantages. For instance, we have in any Grothendieck category enough injectives but often not enough projectives. On the other hand, we obtain a recollement in the unbounded setting which does not exist in the bounded setting. In fact, the celebrated theory of maximal Cohen-Macaulay approximations [2] is described as ‘decomposition’ [2] or ‘glueing’ [12], but finds a natural interpretation as ‘recollement’ in the sense of [6] if one passes to the unbounded setting. To be precise, the recollement

$$\mathbf{S}(\mathrm{Mod}\ \Lambda) \begin{array}{c} \xleftarrow{I_\rho} \\ \rightleftarrows \\ \xrightarrow{I_\lambda} \end{array} \mathbf{K}(\mathrm{Inj}\ \Lambda) \begin{array}{c} \xleftarrow{Q_\rho} \\ \rightleftarrows \\ \xrightarrow{Q_\lambda} \end{array} \mathbf{D}(\mathrm{Mod}\ \Lambda)$$

induces for any Gorenstein ring Λ the Gorenstein injective approximation functor

$$T: \mathrm{Mod}\ \Lambda \xrightarrow{\mathrm{can}} \mathbf{D}(\mathrm{Mod}\ \Lambda) \xrightarrow{I_\lambda \circ Q_\rho} \mathbf{S}(\mathrm{Mod}\ \Lambda) \xrightarrow{Z^0} \underline{\mathrm{Mod}}\ \Lambda$$

where $\underline{\text{Mod}} \Lambda$ denotes the stable category modulo injective objects. For any Λ -module A , the Gorenstein injective approximation $A \rightarrow TA$ is the ‘dual’ of the maximal Cohen-Macaulay approximation which is based on projective resolutions. Let us stress again that this approach generalizes to any locally noetherian Grothendieck category \mathcal{A} provided that $\mathbf{D}(\mathcal{A})$ is compactly generated.

Next we explain the connection between Gorenstein injective approximations and Tate cohomology. We fix a locally noetherian Grothendieck category \mathcal{A} and pass from the stable derived category $\mathbf{S}(\mathcal{A})$ to the full subcategory $\mathbf{T}(\mathcal{A})$ of *totally acyclic* complexes. An object in \mathcal{A} is by definition *Gorenstein injective* if it is of the form $\text{Ker}(X^0 \rightarrow X^1)$ for some X in $\mathbf{T}(\mathcal{A})$. The inclusion $J: \mathbf{T}(\mathcal{A}) \rightarrow \mathbf{K}(\text{Inj } \mathcal{A})$ has a left adjoint J_λ . Given an object A in \mathcal{A} with injective resolution iA , we may think of $J_\lambda iA$ as a complete injective resolution of A . This leads to the following definition of *Tate cohomology groups*

$$\widehat{\text{Ext}}_{\mathcal{A}}^n(A, B) = H^n \text{Hom}_{\mathbf{K}(\mathcal{A})}(A, \Sigma^n(J_\lambda iB))$$

for any A, B in \mathcal{A} and $n \in \mathbb{Z}$. This cohomology theory is symmetric in the sense that for any A in \mathcal{A} , we have $\widehat{\text{Ext}}_{\mathcal{A}}^*(A, -) = 0$ iff $\widehat{\text{Ext}}_{\mathcal{A}}^*(-, A) = 0$ iff $\widehat{\text{Ext}}_{\mathcal{A}}^0(A, A) = 0$. Let \mathcal{X} denote the class of all objects A such that $\widehat{\text{Ext}}_{\mathcal{A}}^*(A, -)$ vanishes, and let \mathcal{Y} be the class of Gorenstein injective objects in \mathcal{A} .

Theorem 1.3. *Let \mathcal{A} be a locally noetherian Grothendieck category and suppose that $\mathbf{D}(\mathcal{A})$ is compactly generated.*

- (1) $\mathcal{X} = \{A \in \mathcal{A} \mid \text{Ext}_{\mathcal{A}}^1(A, B) = 0 \text{ for all } B \in \mathcal{Y}\}$.
- (2) $\mathcal{Y} = \{B \in \mathcal{A} \mid \text{Ext}_{\mathcal{A}}^1(A, B) = 0 \text{ for all } A \in \mathcal{X}\}$.
- (3) *Every object A in \mathcal{A} fits into exact sequences*

$$0 \rightarrow Y_A \rightarrow X_A \rightarrow A \rightarrow 0 \quad \text{and} \quad 0 \rightarrow A \rightarrow Y^A \rightarrow X^A \rightarrow 0$$

in \mathcal{A} with X_A, X^A in \mathcal{X} and Y_A, Y^A in \mathcal{Y} .

- (4) $\mathcal{X} \cap \mathcal{Y} = \text{Inj } \mathcal{A}$.

After explaining some historical background, let us mention more recent work on stable derived categories. For instance, Beligiannis develops a general theory of ‘stabilization’ in the framework of relative homological algebra [4], and Jørgensen studies the category of ‘spectra’ for a module category [19]. Also, Orlov discusses the category

$$\mathbf{D}^b(\text{coh } \mathbb{X}) / \mathbf{D}^{\text{perf}}(\text{coh } \mathbb{X})$$

under the name ‘triangulated category of singularities’ and points out some connection with the Homological Mirror Symmetry Conjecture [28]. In any case, our notation $\mathbf{S}(\text{Qcoh } \mathbb{X})$ reflects this terminology.

Our main results suggests that the homotopy category $\mathbf{K}(\text{Inj } \mathbb{X})$ deserves some more attention. We may think of this category as the ‘compactly generated completion’ of the category $\mathbf{D}^b(\text{coh } \mathbb{X})$. In fact, the category $\text{coh } \mathbb{X}$ of coherent sheaves carries a natural DG structure and its derived category $\mathbf{D}_{\text{dg}}(\text{coh } \mathbb{X})$ is equivalent to $\mathbf{K}(\text{Inj } \mathbb{X})$. This follows from Keller’s work [22] and complements a recent result of Bondal and van den Bergh [9] which says that $\mathbf{D}(\text{Qcoh } \mathbb{X})$ is equivalent to $\mathbf{D}_{\text{dg}}(A)$ for some DG algebra A .

As another application of our main result, let us mention that the adjoint pair of functors $\mathbf{R}f_*$ and $f^!$ which establish the Grothendieck duality for a morphism $f: \mathbb{X} \rightarrow \mathbb{Y}$ between schemes [16, 26], can be extended to a pair of adjoint functors between $\mathbf{K}(\text{Inj } \mathbb{X})$ and $\mathbf{K}(\text{Inj } \mathbb{Y})$.

Theorem 1.4. *Let $f: \mathbb{X} \rightarrow \mathbb{Y}$ be a morphism between separated noetherian schemes. Denote by $\mathbf{R}f_*: \mathbf{D}(\mathrm{Qcoh} \mathbb{X}) \rightarrow \mathbf{D}(\mathrm{Qcoh} \mathbb{Y})$ the right derived direct image functor and by $f^!$ its right adjoint. Then there is an adjoint pair of functors $\hat{\mathbf{R}}f_*$ and $\hat{f}^!$ making the following diagram commutative.*

$$\begin{array}{ccc}
\mathbf{D}(\mathrm{Qcoh} \mathbb{X}) & \xrightarrow{\mathbf{R}f_*} & \mathbf{D}(\mathrm{Qcoh} \mathbb{Y}) & & \mathbf{D}(\mathrm{Qcoh} \mathbb{Y}) & \xrightarrow{f^!} & \mathbf{D}(\mathrm{Qcoh} \mathbb{X}) \\
\downarrow Q_\lambda & & \uparrow Q & & \downarrow Q_\rho & & \uparrow Q \\
\mathbf{K}(\mathrm{Inj} \mathbb{X}) & \xrightarrow{\hat{\mathbf{R}}f_*} & \mathbf{K}(\mathrm{Inj} \mathbb{Y}) & & \mathbf{K}(\mathrm{Inj} \mathbb{Y}) & \xrightarrow{\hat{f}^!} & \mathbf{K}(\mathrm{Inj} \mathbb{X})
\end{array}$$

Again, this theorem is really a lot more general. It is irrelevant that the functor f_* comes from a morphism $f: \mathbb{X} \rightarrow \mathbb{Y}$. All we need is that f_* and its right derived functor $\mathbf{R}f_*$ preserve coproducts. On the other hand, there is a strengthened version of Theorem 1.4 which uses the special properties of f_* . I am grateful to Amnon Neeman for pointing out the following.

Theorem 1.5 (Neeman). *Let $f: \mathbb{X} \rightarrow \mathbb{Y}$ be a morphism between separated noetherian schemes. Then $\hat{\mathbf{R}}f_*$ sends acyclic complexes to acyclic complexes. Thus we have an adjoint pair of functors between $\mathbf{S}(\mathrm{Qcoh} \mathbb{X})$ and $\mathbf{S}(\mathrm{Qcoh} \mathbb{Y})$, making the following diagrams commutative.*

$$\begin{array}{ccccc}
\mathbf{S}(\mathrm{Qcoh} \mathbb{X}) & \xrightarrow{I} & \mathbf{K}(\mathrm{Inj} \mathbb{X}) & \xrightarrow{Q} & \mathbf{D}(\mathrm{Qcoh} \mathbb{X}) \\
\downarrow \mathbf{S}f_* & & \downarrow \hat{\mathbf{R}}f_* & & \downarrow \mathbf{R}f_* \\
\mathbf{S}(\mathrm{Qcoh} \mathbb{Y}) & \xrightarrow{I} & \mathbf{K}(\mathrm{Inj} \mathbb{Y}) & \xrightarrow{Q} & \mathbf{D}(\mathrm{Qcoh} \mathbb{Y}) \\
\\
\mathbf{S}(\mathrm{Qcoh} \mathbb{Y}) & \xleftarrow{I_\rho} & \mathbf{K}(\mathrm{Inj} \mathbb{Y}) & \xleftarrow{Q_\rho} & \mathbf{D}(\mathrm{Qcoh} \mathbb{Y}) \\
\downarrow \mathbf{S}f^! & & \downarrow \hat{f}^! & & \downarrow f^! \\
\mathbf{S}(\mathrm{Qcoh} \mathbb{X}) & \xleftarrow{I_\rho} & \mathbf{K}(\mathrm{Inj} \mathbb{X}) & \xleftarrow{Q_\rho} & \mathbf{D}(\mathrm{Qcoh} \mathbb{X})
\end{array}$$

It seems an interesting project to study the functor $\mathbf{S}f_*$, for instance to find out when it is an equivalence. The following result demonstrates the geometric content of this question; it generalizes a result of Orlov for the bounded stable derived category [28].

Theorem 1.6. *Let \mathbb{Y} be a separated noetherian scheme of finite Krull dimension. If $f: \mathbb{X} \rightarrow \mathbb{Y}$ denotes the inclusion of an open subscheme which contains all singular points of \mathbb{Y} , then $\mathbf{S}f_*: \mathbf{S}(\mathrm{Qcoh} \mathbb{X}) \rightarrow \mathbf{S}(\mathrm{Qcoh} \mathbb{Y})$ is an equivalence.*

Having stated some of the main results, let us sketch the outline of this paper. The paper deals with locally noetherian Grothendieck categories and covers therefore various applications, for instance in algebraic geometry or representation theory. Thus we fix a locally noetherian Grothendieck category \mathcal{A} and study the recollement

$$(1.1) \quad \mathbf{S}(\mathcal{A}) \rightleftarrows \mathbf{K}(\mathrm{Inj} \mathcal{A}) \rightleftarrows \mathbf{D}(\mathcal{A}).$$

More specifically, we begin in Section 2 with the basic properties of the homotopy category $\mathbf{K}(\mathrm{Inj} \mathcal{A})$. The recollement (1.1) is established in Sections 3 and 4. In Section 5, we discuss the essential properties of the stable derived category $\mathbf{S}(\mathcal{A})$. Then we extend

derived functors in Section 6, and Section 7 is devoted to studying Gorenstein injective approximations and Tate cohomology. An appendix provides additional material about DG categories.

2. THE HOMOTOPY CATEGORY OF INJECTIVES

We fix a *locally noetherian* Grothendieck category \mathcal{A} . Thus \mathcal{A} is an abelian Grothendieck category and has a set \mathcal{A}_0 of noetherian objects which *generate* \mathcal{A} , that is, every object in \mathcal{A} is a quotient of a coproduct of objects in \mathcal{A}_0 . We denote by $\text{noeth } \mathcal{A}$ the full subcategory formed by the noetherian objects in \mathcal{A} , and $\text{Inj } \mathcal{A}$ denotes the full subcategory of injective objects. Note that $\text{Inj } \mathcal{A}$ is closed under taking coproducts.

We write $\mathbf{K}(\mathcal{A})$ for the homotopy category and $\mathbf{D}(\mathcal{A})$ for the derived category of unbounded complexes in \mathcal{A} ; for their definitions and basic properties, we refer to [32]. We do not distinguish between an object in \mathcal{A} and the corresponding complex concentrated in degree zero in the homotopy category $\mathbf{K}(\mathcal{A})$. The inclusion $\text{noeth } \mathcal{A} \rightarrow \mathcal{A}$ induces a fully faithful functor

$$\mathbf{D}^b(\text{noeth } \mathcal{A}) \longrightarrow \mathbf{D}(\mathcal{A})$$

which identifies $\mathbf{D}^b(\text{noeth } \mathcal{A})$ with the full subcategory of objects X in $\mathbf{D}(\mathcal{A})$ such that $H^n X$ is noetherian for all n and $H^n X = 0$ for almost all $n \in \mathbb{Z}$.

In this section we study the basic properties of the homotopy category $\mathbf{K}(\text{Inj } \mathcal{A})$. We shall see that this category solves a completion problem for the triangulated category $\mathbf{D}^b(\text{noeth } \mathcal{A})$. Let us begin with some elementary observations.

Lemma 2.1. *Let A be an object in \mathcal{A} and denote by iA an injective resolution. Then the natural map*

$$(2.1) \quad \text{Hom}_{\mathbf{K}(\mathcal{A})}(iA, X) \longrightarrow \text{Hom}_{\mathbf{K}(\mathcal{A})}(A, X)$$

is an isomorphism for all X in $\mathbf{K}(\text{Inj } \mathcal{A})$. Therefore iA is a compact object in $\mathbf{K}(\text{Inj } \mathcal{A})$ if A is noetherian.

Proof. Denote for any $n \in \mathbb{Z}$ by $\sigma^{\geq n} X$ the truncation satisfying

$$(\sigma^{\geq n} X)^p = \begin{cases} X^p & \text{if } p \geq n, \\ 0 & \text{if } p < n. \end{cases}$$

We complete the map $A \rightarrow iA$ to an exact triangle

$$aA \longrightarrow A \longrightarrow iA \longrightarrow \Sigma(aA)$$

and obtain

$$0 = \text{Hom}_{\mathbf{K}(\mathcal{A})}(aA, \sigma^{\geq -1} X) \cong \text{Hom}_{\mathbf{K}(\mathcal{A})}(aA, X)$$

since aA is acyclic and concentrated in non-negative degrees. Thus $\text{Hom}_{\mathbf{K}(\mathcal{A})}(iA, X) \cong \text{Hom}_{\mathbf{K}(\mathcal{A})}(A, X)$.

Now assume that A is noetherian. Clearly, A is a compact object in \mathcal{A} and therefore a compact object in $\mathbf{K}(\mathcal{A})$. The isomorphism (2.1) shows that iA is a compact object in $\mathbf{K}(\text{Inj } \mathcal{A})$. \square

Lemma 2.2. *Let X be a non-zero object in $\mathbf{K}(\text{Inj } \mathcal{A})$. Then there exists a noetherian object A in \mathcal{A} such that $\text{Hom}_{\mathbf{K}(\mathcal{A})}(A, \Sigma^n X) \neq 0$ for some $n \in \mathbb{Z}$.*

Proof. Suppose first $H^n X \neq 0$ for some n . Choose a noetherian object A and a map $A \rightarrow Z^n X$ inducing a non-zero map $A \rightarrow H^n X$. We obtain a chain map $A \rightarrow \Sigma^n X$ which induces a non-zero element in $\mathrm{Hom}_{\mathbf{K}(\mathcal{A})}(A, \Sigma^n X)$.

Now suppose $H^n X = 0$ for all n . We can choose n such that $Z^n X$ is non-injective. Using Baer's criterion, there exists a noetherian object A in \mathcal{A} such that $\mathrm{Ext}_{\mathcal{A}}^1(A, Z^n X)$ is non-zero. Now observe that

$$\mathrm{Hom}_{\mathbf{K}(\mathcal{A})}(A, \Sigma^{n+p} X) \cong \mathrm{Ext}_{\mathcal{A}}^p(A, Z^n X)$$

for all $p \geq 1$. Thus $\mathrm{Hom}_{\mathbf{K}(\mathcal{A})}(A, \Sigma^{n+1} X) \neq 0$. This completes the proof. \square

Let \mathcal{T} be a triangulated category with arbitrary coproducts. Recall that an object X in \mathcal{T} is *compact* if $\mathrm{Hom}_{\mathcal{T}}(X, -)$ preserves all coproducts. The triangulated category is *compactly generated* if there is a set \mathcal{T}_0 of compact objects such that $\mathrm{Hom}_{\mathcal{T}}(X, \Sigma^n Y) = 0$ for all $X \in \mathcal{T}_0$ and $n \in \mathbb{Z}$ implies $Y = 0$ for every object Y in \mathcal{T} .

Proposition 2.3. *Let \mathcal{A} be a locally noetherian Grothendieck category, and let $\mathbf{K}^c(\mathrm{Inj} \mathcal{A})$ denote the full subcategory of compact objects in $\mathbf{K}(\mathrm{Inj} \mathcal{A})$.*

- (1) *The triangulated category $\mathbf{K}(\mathrm{Inj} \mathcal{A})$ is compactly generated.*
- (2) *The canonical functor $\mathbf{K}(\mathcal{A}) \rightarrow \mathbf{D}(\mathcal{A})$ induces an equivalence*

$$\mathbf{K}^c(\mathrm{Inj} \mathcal{A}) \xrightarrow{\sim} \mathbf{D}^b(\mathrm{noeth} \mathcal{A}).$$

Proof. It follows from Lemma 2.1 and Lemma 2.2 that $\mathbf{K}(\mathrm{Inj} \mathcal{A})$ is compactly generated. A standard argument shows that $\mathbf{K}^c(\mathrm{Inj} \mathcal{A})$ equals the thick subcategory of $\mathbf{K}(\mathrm{Inj} \mathcal{A})$ which is generated by the injective resolutions of the noetherian objects in \mathcal{A} ; see [25, 2.2]. The equivalence $\mathbf{K}^+(\mathrm{Inj} \mathcal{A}) \rightarrow \mathbf{D}^+(\mathcal{A})$ restricts to an equivalence $\mathbf{K}^{+,b}(\mathrm{Inj} \mathcal{A}) \rightarrow \mathbf{D}^b(\mathcal{A})$ and identifies $\mathbf{K}^c(\mathrm{Inj} \mathcal{A})$ with $\mathbf{D}^b(\mathrm{noeth} \mathcal{A})$. \square

Note that we obtain a functor $\mathbf{D}^b(\mathrm{noeth} \mathcal{A}) \rightarrow \mathbf{K}(\mathrm{Inj} \mathcal{A})$ which identifies $\mathbf{D}^b(\mathrm{noeth} \mathcal{A})$ with the full subcategory of compact objects. Therefore the formation of the category $\mathbf{K}(\mathrm{Inj} \mathcal{A})$ solves a completion problem which we explain by an analogy. The category \mathcal{A} is a completion of $\mathrm{noeth} \mathcal{A}$ in the following sense.

- \mathcal{A} is an additive category with filtered colimits.
- The inclusion $\mathrm{noeth} \mathcal{A} \rightarrow \mathcal{A}$ identifies $\mathrm{noeth} \mathcal{A}$ with the full subcategory of finitely presented objects.
- \mathcal{A} coincides with the smallest subcategory which contains all finitely presented objects and is closed under forming filtered colimits.

Recall that an object X in \mathcal{A} is *finitely presented* if the functor $\mathrm{Hom}_{\mathcal{A}}(X, -)$ preserves filtered colimits. Similarly, we have the following for $\mathcal{T} = \mathbf{K}(\mathrm{Inj} \mathcal{A})$.

- \mathcal{T} is a triangulated category with coproducts.
- The functor $\mathbf{D}^b(\mathrm{noeth} \mathcal{A}) \rightarrow \mathcal{T}$ identifies $\mathbf{D}^b(\mathrm{noeth} \mathcal{A})$ with the full subcategory of compact objects.
- \mathcal{T} coincides with the smallest subcategory which contains all compact objects and is closed under forming triangles and coproducts.

The category \mathcal{A} is, up to an equivalence, uniquely determined by $\mathrm{noeth} \mathcal{A}$. It would be interesting to know to what extent $\mathbf{K}(\mathrm{Inj} \mathcal{A})$ is uniquely determined by $\mathbf{D}^b(\mathrm{noeth} \mathcal{A})$.

Example 2.4. Suppose there is a noetherian object A in \mathcal{A} such that $\mathbf{D}^b(\mathrm{noeth} \mathcal{A})$ is generated by A , that is, there is no proper thick subcategory containing A . Take an

injective resolution iA and denote by $\mathcal{E}nd_{\mathcal{A}}(A)$ the endomorphism DG algebra of iA . Then $\mathcal{H}om_{\mathcal{A}}(iA, -)$ induces an equivalence between $\mathbf{K}(\text{Inj } \mathcal{A})$ and the derived category $\mathbf{D}_{\text{dg}}(\mathcal{E}nd_{\mathcal{A}}(A))$ of DG $\mathcal{E}nd_{\mathcal{A}}(A)$ -modules; see [22]. If one replaces a single generator by a set of generating objects, then one obtains an analogue which involves a DG category instead of a DG algebra. In particular, $\text{noeth } \mathcal{A}$ carries the structure of a DG category such that $\mathbf{K}(\text{Inj } \mathcal{A})$ and $\mathbf{D}_{\text{dg}}(\text{noeth } \mathcal{A})$ are equivalent. We refer to Appendix A for details.

Example 2.5. Let G be a finite p -group and k be a field of characteristic $p > 0$. We consider the category $\mathcal{A} = \text{Mod } kG$ of modules over the group algebra kG . Take an injective resolution ik of the trivial representation k , and denote by $\mathcal{E}nd_{kG}(k)$ the endomorphism DG algebra of ik . Then its derived category $\mathbf{D}_{\text{dg}}(\mathcal{E}nd_{kG}(k))$ is equivalent to $\mathbf{K}(\text{Inj } \mathcal{A})$. The tensor product \otimes_k on \mathcal{A} restricts to a product on $\text{Inj } \mathcal{A}$ and induces therefore a (total) tensor product on $\mathbf{K}(\text{Inj } \mathcal{A})$. On the other hand, the E_{∞} -structure of $\mathcal{E}nd_{kG}(k)$ induces a product on $\mathbf{D}_{\text{dg}}(\mathcal{E}nd_{kG}(k))$. We conjecture that these products are naturally isomorphic.

Example 2.6. Let Λ be a finite dimensional algebra over a field k . Then $E = \text{Hom}_k(\Lambda^{\text{op}}, k)$ is an injective cogenerator for $\mathcal{A} = \text{Mod } \Lambda$, and $\text{Hom}_{\Lambda}(E, -)$ induces an equivalence $\text{Inj } \mathcal{A} \rightarrow \text{Proj } \mathcal{A}$ since $\text{Hom}_{\Lambda}(E, E) \cong \Lambda$. Thus the homotopy category $\mathbf{K}(\text{Proj } \mathcal{A})$ is compactly generated. For more on $\mathbf{K}(\text{Proj } \mathcal{A})$, see [19, 20].

3. A LOCALIZATION SEQUENCE

Let \mathcal{A} be a locally noetherian Grothendieck category and let

$$\mathbf{K}_{\text{ac}}(\text{Inj } \mathcal{A}) = \mathbf{K}(\text{Inj } \mathcal{A}) \cap \mathbf{K}_{\text{ac}}(\mathcal{A}).$$

In this section we prove that the canonical functors

$$I: \mathbf{K}_{\text{ac}}(\text{Inj } \mathcal{A}) \xrightarrow{\text{inc}} \mathbf{K}(\text{Inj } \mathcal{A}) \quad \text{and} \quad Q: \mathbf{K}(\text{Inj } \mathcal{A}) \xrightarrow{\text{inc}} \mathbf{K}(\mathcal{A}) \xrightarrow{\text{can}} \mathbf{D}(\mathcal{A})$$

form a localization sequence

$$(3.1) \quad \mathbf{K}_{\text{ac}}(\text{Inj } \mathcal{A}) \xrightarrow{I} \mathbf{K}(\text{Inj } \mathcal{A}) \xrightarrow{Q} \mathbf{D}(\mathcal{A}).$$

Let us start with some preparations. In particular, we need to give the definition of a localization sequence.

Definition 3.1. We say that a sequence

$$\mathcal{T}' \xrightarrow{F} \mathcal{T} \xrightarrow{G} \mathcal{T}''$$

of exact functors between triangulated categories is a *localization sequence* if the following holds.

- (L1) The functor F has a right adjoint $F_{\rho}: \mathcal{T} \rightarrow \mathcal{T}'$ satisfying $F_{\rho} \circ F \cong \text{Id}_{\mathcal{T}'}$.
- (L2) The functor G has a right adjoint $G_{\rho}: \mathcal{T}'' \rightarrow \mathcal{T}$ satisfying $G \circ G_{\rho} \cong \text{Id}_{\mathcal{T}''}$.
- (L3) Let X be an object in \mathcal{T} . Then $GX = 0$ if and only if $X \cong FX'$ for some $X' \in \mathcal{T}'$.

The sequence (F, G) of functors is called *colocalization sequence* if the sequence $(F^{\text{op}}, G^{\text{op}})$ of opposite functors is a localization sequence.

The basic properties of a localization sequence are the following [32, II.2].

- (1) The functors F and G_{ρ} are fully faithful.

(2) Identify $\mathcal{T}' = \text{Im } F$ and $\mathcal{T}'' = \text{Im } G_\rho$. Given objects $X, Y \in \mathcal{T}$, then

$$\begin{aligned} X \in \mathcal{T}' &\iff \text{Hom}_{\mathcal{T}}(X, \mathcal{T}'') = 0, \\ Y \in \mathcal{T}'' &\iff \text{Hom}_{\mathcal{T}}(\mathcal{T}', Y) = 0. \end{aligned}$$

(3) Identify $\mathcal{T}' = \text{Im } F$. Then the functor G induces an equivalence $\mathcal{T}/\mathcal{T}' \rightarrow \mathcal{T}''$.

(4) Let X be an object in \mathcal{T} . Then there is an exact triangle

$$(F \circ F_\rho)X \longrightarrow X \longrightarrow (G_\rho \circ G)X \longrightarrow \Sigma((F \circ F_\rho)X)$$

which is functorial in X .

(5) The sequence

$$\mathcal{T}'' \xrightarrow{G_\rho} \mathcal{T} \xrightarrow{F_\rho} \mathcal{T}'$$

is a colocalization sequence.

The next lemma is well known; it provides useful criteria for a sequence to be a localization sequence. Recall that a full subcategory of a triangulated category is *thick* if it is a triangulated subcategory which is closed under taking direct factors.

Lemma 3.2. *Let \mathcal{T} be a triangulated category and \mathcal{S} be a thick subcategory. Then the following are equivalent.*

- (1) *The sequence $\mathcal{S} \xrightarrow{\text{inc}} \mathcal{T} \xrightarrow{\text{can}} \mathcal{T}/\mathcal{S}$ is a localization sequence.*
- (2) *The inclusion functor $\mathcal{S} \rightarrow \mathcal{T}$ has a right adjoint.*
- (3) *The quotient functor $\mathcal{T} \rightarrow \mathcal{T}/\mathcal{S}$ has a right adjoint.*

Proof. Condition (1) implies (2) and (3). Also, (2) and (3) together imply (1). Thus we need to show that (2) and (3) are equivalent. Let us write $F: \mathcal{S} \rightarrow \mathcal{T}$ and $G: \mathcal{T} \rightarrow \mathcal{T}/\mathcal{S}$ for the functors which are involved.

(2) \Rightarrow (3): Let \mathcal{T}'' denote the full subcategory of objects Y in \mathcal{T} such that $\text{Hom}_{\mathcal{T}}(\mathcal{S}, Y) = 0$. We obtain a left adjoint $L: \mathcal{T} \rightarrow \mathcal{T}''$ for the inclusion by completing for each X in \mathcal{T} the natural map $F_\rho X \rightarrow X$ to an exact triangle

$$F_\rho X \longrightarrow X \longrightarrow LX \longrightarrow \Sigma(F_\rho X).$$

The functor L annihilates \mathcal{S} and factors therefore through G via an exact functor $\bar{L}: \mathcal{T}/\mathcal{S} \rightarrow \mathcal{T}''$. The composite of \bar{L} with the inclusion $\mathcal{T}'' \rightarrow \mathcal{T}$ is a right adjoint for G .

(3) \Rightarrow (2): We obtain a right adjoint $F_\rho: \mathcal{T} \rightarrow \mathcal{S}$ for the inclusion F by completing for each X in \mathcal{T} the natural map $X \rightarrow (G_\rho \circ G)X$ to an exact triangle

$$F_\rho X \longrightarrow X \longrightarrow (G_\rho \circ G)X \longrightarrow \Sigma(F_\rho X).$$

Note that $F_\rho X$ belongs to \mathcal{S} since $G(F_\rho X) = 0$. □

We need to construct left and right adjoints for functors starting in a compactly generated triangulated category. Our basic tool for this is the following result which is due to Neeman.

Proposition 3.3. *Let $F: \mathcal{S} \rightarrow \mathcal{T}$ be an exact functor between triangulated categories, and suppose \mathcal{S} is compactly generated.*

- (1) *There is a right adjoint $\mathcal{T} \rightarrow \mathcal{S}$ if and only if F preserves all coproducts.*
- (2) *There is a left adjoint $\mathcal{T} \rightarrow \mathcal{S}$ if and only if F preserves all products.*

Proof. For (1), see [26, Theorem 4.1]. The proof of (2) is analogous and uses covariant Brown representability [27, Theorem 8.6.1]; see also [24]. \square

We record a similar result for later use.

Proposition 3.4. *Let \mathcal{T} be a compactly generated triangulated category and \mathcal{S}_0 be a set of objects in \mathcal{T} . Denote by \mathcal{U} the full subcategory of objects Y in \mathcal{T} such that $\mathrm{Hom}_{\mathcal{T}}(\Sigma^n X, Y) = 0$ for all $X \in \mathcal{S}_0$ and $n \in \mathbb{Z}$. Then the inclusion $\mathcal{U} \rightarrow \mathcal{T}$ has a left adjoint.*

Proof. The localizing subcategory \mathcal{S} generated by \mathcal{S}_0 is well generated and the inclusion $\mathcal{S} \rightarrow \mathcal{T}$ has therefore a right adjoint; see [27]. We obtain a localization sequence $\mathcal{S} \xrightarrow{\mathrm{inc}} \mathcal{T} \xrightarrow{\mathrm{can}} \mathcal{T}/\mathcal{S}$ by Lemma 3.2, and the right adjoint of the canonical functor $\mathcal{T} \rightarrow \mathcal{T}/\mathcal{S}$ identifies \mathcal{T}/\mathcal{S} with \mathcal{U} . \square

There is a useful criterion when a left adjoint preserves compactness.

Lemma 3.5. *Let $F: \mathcal{S} \rightarrow \mathcal{T}$ be an exact functor between compactly generated triangulated categories which has a right adjoint G . Then F preserves compactness if and only if G preserves coproducts.*

Proof. See [26, Theorem 5.1]. \square

The following result establishes the localization sequence for the homotopy category of injective objects.

Proposition 3.6. *Let \mathcal{A} be a locally noetherian Grothendieck category. Then the canonical functors $\mathbf{K}_{\mathrm{ac}}(\mathrm{Inj} \mathcal{A}) \rightarrow \mathbf{K}(\mathrm{Inj} \mathcal{A})$ and $\mathbf{K}(\mathrm{Inj} \mathcal{A}) \rightarrow \mathbf{D}(\mathcal{A})$ form a localization sequence*

$$\mathbf{K}_{\mathrm{ac}}(\mathrm{Inj} \mathcal{A}) \xrightarrow{I} \mathbf{K}(\mathrm{Inj} \mathcal{A}) \xrightarrow{Q} \mathbf{D}(\mathcal{A}).$$

Proof. We know from Proposition 2.3 that $\mathbf{K}(\mathrm{Inj} \mathcal{A})$ is compactly generated. In addition, we use Lemma 3.2 and Proposition 3.3. The inclusion $J: \mathbf{K}(\mathrm{Inj} \mathcal{A}) \rightarrow \mathbf{K}(\mathcal{A})$ preserves products and has therefore a left adjoint J_λ satisfying $J_\lambda \circ J \cong \mathrm{Id}_{\mathbf{K}(\mathrm{Inj} \mathcal{A})}$. We obtain a localization sequence

$$\mathcal{K} \xrightarrow{\mathrm{inc}} \mathbf{K}(\mathcal{A}) \xrightarrow{J_\lambda} \mathbf{K}(\mathrm{Inj} \mathcal{A})$$

where \mathcal{K} denotes the kernel of J_λ . Thus

$$\mathrm{Hom}_{\mathbf{K}(\mathcal{A})}(X, Y) = 0 \quad \text{for all } X \in \mathcal{K} \quad \text{and} \quad Y \in \mathbf{K}(\mathrm{Inj} \mathcal{A}).$$

This implies $\mathcal{K} \subseteq \mathbf{K}_{\mathrm{ac}}(\mathcal{A})$ and gives the following commutative diagram of exact functors.

$$\begin{array}{ccccc} \mathcal{K} & \xrightarrow{\mathrm{inc}} & \mathbf{K}(\mathcal{A}) & \xrightarrow{J_\lambda} & \mathbf{K}(\mathrm{Inj} \mathcal{A}) \\ \downarrow \mathrm{inc} & & \parallel & & \downarrow F \\ \mathbf{K}_{\mathrm{ac}}(\mathcal{A}) & \xrightarrow{\mathrm{inc}} & \mathbf{K}(\mathcal{A}) & \xrightarrow{\mathrm{can}} & \mathbf{D}(\mathcal{A}) \end{array}$$

The functor F is induced by the canonical functor $\mathbf{K}(\mathcal{A}) \rightarrow \mathbf{D}(\mathcal{A})$, and we have $F \cong Q$ since $J_\lambda \circ J \cong \mathrm{Id}_{\mathbf{K}(\mathrm{Inj} \mathcal{A})}$. Moreover, F preserves coproducts and has therefore a right adjoint F_ρ . The composite $J \circ F_\rho$ is a right adjoint for the canonical functor $\mathbf{K}(\mathcal{A}) \rightarrow \mathbf{D}(\mathcal{A})$. This implies $F \circ F_\rho \cong \mathrm{Id}_{\mathbf{D}(\mathcal{A})}$. On the other hand, $\mathbf{K}_{\mathrm{ac}}(\mathrm{Inj} \mathcal{A})$ is the kernel of F . Thus we conclude that the sequence (3.1) is a localization sequence. \square

We add some useful remarks which are immediate consequences.

Remark 3.7. Let $J_\lambda: \mathbf{K}(\mathcal{A}) \rightarrow \mathbf{K}(\text{Inj } \mathcal{A})$ be the left adjoint of the inclusion $\mathbf{K}(\text{Inj } \mathcal{A}) \rightarrow \mathbf{K}(\mathcal{A})$. Then the composite $Q \circ J_\lambda$ is naturally isomorphic to the canonical functor $\mathbf{K}(\mathcal{A}) \rightarrow \mathbf{D}(\mathcal{A})$.

Remark 3.8. The right adjoint Q_ρ of Q induces an equivalence

$$\mathbf{D}^b(\text{noeth } \mathcal{A}) \xrightarrow{\sim} \mathbf{K}^c(\text{Inj } \mathcal{A})$$

which is a quasi-inverse for the equivalence $\mathbf{K}^c(\text{Inj } \mathcal{A}) \rightarrow \mathbf{D}^b(\text{noeth } \mathcal{A})$ induced by Q .

Let us denote by $\mathbf{K}_{\text{inj}}(\mathcal{A})$ the full subcategory of complexes Y in $\mathbf{K}(\text{Inj } \mathcal{A})$ such that $\text{Hom}_{\mathbf{K}(\mathcal{A})}(X, Y) = 0$ for all acyclic complexes X in $\mathbf{K}(\mathcal{A})$. Following Spaltenstein's terminology [31], the objects in $\mathbf{K}_{\text{inj}}(\mathcal{A})$ are precisely the *K-injective* complexes having injective components. There are various results about K-injective resolutions in the literature; see for instance [31, 10]. The following is certainly not the most general one; however it is sufficient in our context.

Corollary 3.9. *The inclusion $\mathbf{K}_{\text{inj}}(\mathcal{A}) \rightarrow \mathbf{K}(\mathcal{A})$ has a left adjoint $i: \mathbf{K}(\mathcal{A}) \rightarrow \mathbf{K}_{\text{inj}}(\mathcal{A})$ which has the following properties.*

- (1) *Every object X in $\mathbf{K}(\mathcal{A})$ fits into an exact triangle*

$$aX \longrightarrow X \longrightarrow iX \longrightarrow \Sigma(aX)$$

such that aX is an acyclic complex.

- (2) *The functor $i: \mathbf{K}(\mathcal{A}) \rightarrow \mathbf{K}_{\text{inj}}(\mathcal{A})$ induces an equivalence*

$$\mathbf{D}(\mathcal{A}) = \mathbf{K}(\mathcal{A})/\mathbf{K}_{\text{ac}}(\mathcal{A}) \xrightarrow{\sim} \mathbf{K}_{\text{inj}}(\mathcal{A}).$$

- (3) *We have for all X, Y in $\mathbf{K}(\mathcal{A})$*

$$\text{Hom}_{\mathbf{D}(\mathcal{A})}(X, Y) \cong \text{Hom}_{\mathbf{K}(\mathcal{A})}(X, iY).$$

Proof. Put $iX = Q_\rho X$ for each X in $\mathbf{K}(\mathcal{A})$, where Q_ρ denotes the right adjoint of $Q: \mathbf{K}(\text{Inj } \mathcal{A}) \rightarrow \mathbf{D}(\mathcal{A})$. \square

The functor

$$R: \mathbf{D}(\mathcal{A}) = \mathbf{K}(\mathcal{A})/\mathbf{K}_{\text{ac}}(\mathcal{A}) \xrightarrow{\sim} \mathbf{K}_{\text{inj}}(\mathcal{A}) \xrightarrow{\text{inc}} \mathbf{K}(\mathcal{A})$$

provides a right adjoint for the canonical functor $\mathbf{K}(\mathcal{A}) \rightarrow \mathbf{D}(\mathcal{A})$. Let us mention as an application that the right derived functor of any additive functor $F: \mathcal{A} \rightarrow \mathcal{B}$ is obtained as composite

$$\mathbf{R}F: \mathbf{D}(\mathcal{A}) \xrightarrow{R} \mathbf{K}(\mathcal{A}) \xrightarrow{\mathbf{K}(F)} \mathbf{K}(\mathcal{B}) \xrightarrow{\text{can}} \mathbf{D}(\mathcal{B}).$$

Example 3.10. Suppose every object in \mathcal{A} has finite injective dimension. Then the functor $\mathbf{K}(\text{Inj } \mathcal{A}) \rightarrow \mathbf{D}(\mathcal{A})$ is an equivalence since $\mathbf{K}_{\text{ac}}(\text{Inj } \mathcal{A}) = 0$. In particular, the compact objects in $\mathbf{D}(\mathcal{A})$ are precisely those from $\mathbf{D}^b(\text{noeth } \mathcal{A})$.

Example 3.11. Suppose products in \mathcal{A} are exact. For instance, let \mathcal{A} be a module category. Then one can show that $\mathbf{K}_{\text{inj}}(\mathcal{A})$ is the smallest triangulated subcategory of $\mathbf{K}(\mathcal{A})$ which is closed under taking products and contains the injective objects of \mathcal{A} (viewed as complexes concentrated in degree zero).

4. A RECOLLEMENT

In this section we provide a criterion for \mathcal{A} such that the sequence

$$\mathbf{K}_{\text{ac}}(\text{Inj } \mathcal{A}) \xrightarrow{I} \mathbf{K}(\text{Inj } \mathcal{A}) \xrightarrow{Q} \mathbf{D}(\mathcal{A})$$

induces a recollement

$$\mathbf{K}_{\text{ac}}(\text{Inj } \mathcal{A}) \rightleftarrows \mathbf{K}(\text{Inj } \mathcal{A}) \rightleftarrows \mathbf{D}(\mathcal{A})$$

in the sense of [6]. It is important to note that one cannot expect a recollement

$$(4.1) \quad \mathbf{K}_{\text{ac}}(\mathcal{A}) \rightleftarrows \mathbf{K}(\mathcal{A}) \rightleftarrows \mathbf{D}(\mathcal{A})$$

without severe restrictions on \mathcal{A} ; see Example 4.9. In fact, a recollement (4.1) implies that a product of exact sequences in \mathcal{A} remains exact.

We begin with a lemma.

Lemma 4.1. *Let \mathcal{A} be a locally noetherian Grothendieck category. Then a compact object in $\mathbf{D}(\mathcal{A})$ belongs to $\mathbf{D}^b(\text{noeth } \mathcal{A})$.*

Proof. Suppose X is compact in $\mathbf{D}(\mathcal{A})$. We need to show that $H^n X$ is noetherian for all n , and that $H^n X$ vanishes for almost all n in \mathbb{Z} . We have for any injective object E in \mathcal{A} an isomorphism

$$\text{Hom}_{\mathbf{D}(\mathcal{A})}(X, E) \cong \text{Hom}_{\mathcal{A}}(H^0 X, E).$$

Therefore $\text{Hom}_{\mathcal{A}}(H^0 X, -)$ preserves coproducts in $\text{Inj } \mathcal{A}$. This implies that each $H^n X$ is noetherian; see [29]. Now fix for each n an injective envelope $H^n X \rightarrow E(H^n X)$ and consider the induced map

$$\alpha: X \longrightarrow \prod_{n \in \mathbb{Z}} \Sigma^{-n} E(H^n X)$$

in $\mathbf{D}(\mathcal{A})$. The canonical map

$$\prod_{n \in \mathbb{Z}} \Sigma^{-n} E(H^n X) \longrightarrow \prod_{n \in \mathbb{Z}} \Sigma^{-n} E(H^n X)$$

is an isomorphism in $\mathbf{D}(\mathcal{A})$, and therefore α factors through a finite number of factors in

$$\prod_{n \in \mathbb{Z}} \Sigma^{-n} E(H^n X).$$

Thus $H^n X$ vanishes for almost all n in \mathbb{Z} , and the proof is complete. \square

We denote by $\mathbf{D}^c(\mathcal{A})$ the full subcategory of $\mathbf{D}(\mathcal{A})$ which is formed by all compact objects.

Theorem 4.2. *Let \mathcal{A} be a locally noetherian Grothendieck category and suppose $\mathbf{D}(\mathcal{A})$ is compactly generated. Then the canonical functor $Q: \mathbf{K}(\text{Inj } \mathcal{A}) \rightarrow \mathbf{D}(\mathcal{A})$ has a left adjoint and therefore the sequence*

$$\mathbf{K}_{\text{ac}}(\text{Inj } \mathcal{A}) \xrightarrow{I} \mathbf{K}(\text{Inj } \mathcal{A}) \xrightarrow{Q} \mathbf{D}(\mathcal{A})$$

is a colocalization sequence.

Proof. Let \mathcal{K} be the localizing subcategory of $\mathbf{K}(\text{Inj } \mathcal{A})$ which is generated by all compact objects X in $\mathbf{K}(\text{Inj } \mathcal{A})$ such that QX is compact in $\mathbf{D}(\mathcal{A})$. We claim that $Q|_{\mathcal{K}}: \mathcal{K} \rightarrow \mathbf{D}(\mathcal{A})$ is an equivalence. First note that \mathcal{K} and $\mathbf{D}(\mathcal{A})$ are both compactly generated. We have seen in Lemma 4.1 that

$$\mathbf{D}^c(\mathcal{A}) \subseteq \mathbf{D}^b(\text{noeth } \mathcal{A}),$$

and Q induces an equivalence

$$\mathbf{K}^c(\text{Inj } \mathcal{A}) \xrightarrow{\sim} \mathbf{D}^b(\text{noeth } \mathcal{A}),$$

by Proposition 2.3. Thus Q induces an equivalence between the subcategories of compact objects in \mathcal{K} and $\mathbf{D}(\mathcal{A})$. Then a standard argument shows that $Q|_{\mathcal{K}}$ is an equivalence since Q preserves all coproducts. Now fix a left adjoint $L: \mathbf{D}(\mathcal{A}) \rightarrow \mathcal{K}$. We claim that the composite

$$\mathbf{D}(\mathcal{A}) \xrightarrow{L} \mathcal{K} \xrightarrow{\text{inc}} \mathbf{K}(\text{Inj } \mathcal{A})$$

is a left adjoint for Q . To see this, consider for objects X in $\mathbf{D}(\mathcal{A})$ and Y in $\mathbf{K}(\text{Inj } \mathcal{A})$ the natural map

$$\alpha_{X,Y}: \text{Hom}_{\mathbf{K}(\text{Inj } \mathcal{A})}(LX, Y) \longrightarrow \text{Hom}_{\mathbf{D}(\mathcal{A})}(QLX, QY) \xrightarrow{\sim} \text{Hom}_{\mathbf{D}(\mathcal{A})}(X, QY)$$

which is induced by Q . If X and Y are compact, then $\alpha_{X,Y}$ is bijective by Proposition 2.3. We use a standard argument to show that $\alpha_{X,Y}$ is bijective for arbitrary X and Y . Fix a compact object X . Then the objects Y such that $\alpha_{X,Y}$ is bijective form a triangulated subcategory which is closed under taking coproducts and contains all compact objects. Thus $\alpha_{X,Y}$ is bijective for all Y because $\mathbf{K}(\text{Inj } \mathcal{A})$ is compactly generated. Now fix any object Y . The same argument shows that $\alpha_{X,Y}$ is bijective for all X because $\mathbf{D}(\mathcal{A})$ is compactly generated. We conclude that Q has a left adjoint. Moreover, Lemma 3.2 implies that I and Q form a colocalization sequence. \square

Following Beilinson, Bernstein, and Deligne [6], we say that a sequence

$$(4.2) \quad \mathcal{T}' \longrightarrow \mathcal{T} \longrightarrow \mathcal{T}''$$

of exact functors between triangulated categories induces a *recollement*

$$\mathcal{T}' \rightleftarrows \mathcal{T} \rightleftarrows \mathcal{T}''$$

if the sequence (4.2) is a localization sequence and a colocalization sequence in the sense of Definition 3.1.

Corollary 4.3. *Let \mathcal{A} be a locally noetherian Grothendieck category and suppose $\mathbf{D}(\mathcal{A})$ is compactly generated. Then the sequence*

$$\mathbf{K}_{\text{ac}}(\text{Inj } \mathcal{A}) \xrightarrow{I} \mathbf{K}(\text{Inj } \mathcal{A}) \xrightarrow{Q} \mathbf{D}(\mathcal{A})$$

induces a recollement

$$\mathbf{K}_{\text{ac}}(\text{Inj } \mathcal{A}) \rightleftarrows \mathbf{K}(\text{Inj } \mathcal{A}) \rightleftarrows \mathbf{D}(\mathcal{A}).$$

Corollary 4.4. *Let \mathcal{A} be a locally noetherian Grothendieck category and suppose $\mathbf{D}(\mathcal{A})$ is compactly generated. Then a product of acyclic complexes of injective objects in \mathcal{A} is acyclic.*

Let us give a criterion for \mathcal{A} such that the derived category $\mathbf{D}(\mathcal{A})$ is compactly generated.

Lemma 4.5. *Let \mathcal{A} be a locally noetherian Grothendieck category. Suppose there is a set \mathcal{A}_0 of objects in \mathcal{A} which are compact when viewed as objects in $\mathbf{D}(\mathcal{A})$. If \mathcal{A}_0 generates \mathcal{A} , then $\mathbf{D}(\mathcal{A})$ is compactly generated by \mathcal{A}_0 .*

The lemma is an immediate consequence of the following statement.

Lemma 4.6. *Let \mathcal{A} be a locally noetherian Grothendieck category and fix a set \mathcal{A}_0 of generating objects. Let X be a complex in \mathcal{A} such that $H^0 X \neq 0$. Then there exists some object A in \mathcal{A}_0 such that*

$$\mathrm{Hom}_{\mathbf{K}(\mathcal{A})}(A, X) \neq 0 \quad \text{and} \quad \mathrm{Hom}_{\mathbf{D}(\mathcal{A})}(A, X) \neq 0.$$

Proof. Choose A in \mathcal{A}_0 and a map $A \rightarrow Z^0 X$ such that the composite with $Z^0 X \rightarrow H^0 X$ is non-zero. This induces a non-zero element in

$$H^0(\mathrm{Hom}_{\mathcal{A}}(A, X)) \cong \mathrm{Hom}_{\mathbf{K}(\mathcal{A})}(A, X).$$

The second assertion follows from the first since for any object A in \mathcal{A} we have

$$\mathrm{Hom}_{\mathbf{D}(\mathcal{A})}(A, X) \cong \mathrm{Hom}_{\mathbf{K}(\mathcal{A})}(A, iX)$$

and $H^0(iX) \cong H^0 X$. □

We give examples of Grothendieck categories such that objects in \mathcal{A} become compact objects in $\mathbf{D}(\mathcal{A})$.

Example 4.7. Let Λ be an associative ring. Denote by $\mathcal{A} = \mathrm{Mod} \Lambda$ the category of (right) Λ -modules and by $\mathrm{proj} \Lambda$ the full subcategory of finitely generated projective Λ -modules. Then every object in $\mathrm{proj} \Lambda$ is compact when viewed as object in $\mathbf{D}(\mathcal{A})$. Thus the inclusion $\mathbf{D}^b(\mathrm{proj} \Lambda) \rightarrow \mathbf{D}(\mathcal{A})$ identifies $\mathbf{D}^b(\mathrm{proj} \Lambda)$ with the full subcategory of compact objects in $\mathbf{D}(\mathcal{A})$. Suppose now that Λ is right noetherian. Then the fully faithful functor $Q_\lambda: \mathbf{D}(\mathcal{A}) \rightarrow \mathbf{K}(\mathrm{Inj} \Lambda)$ identifies $\mathbf{D}(\mathcal{A})$ with the localizing subcategory of $\mathbf{K}(\mathrm{Inj} \Lambda)$ which is generated by the injective resolution $i\Lambda$ of Λ .

Let us return to the completion problem for triangulated categories which has been adressed in Section 2. Keeping the analogy between the completion with respect to filtered colimits and the completion with respect to triangles und coproducts, we obtain the following diagram for a right noetherian ring Λ . The vertical arrows denote completions and the horizontal ones the appropriate inclusions.

$$\begin{array}{ccc} \mathrm{proj} \Lambda & \longrightarrow & \mathrm{mod} \Lambda \\ \downarrow & & \downarrow \\ \mathrm{Flat} \Lambda & \longrightarrow & \mathrm{Mod} \Lambda \end{array} \quad \begin{array}{ccc} \mathbf{D}^b(\mathrm{proj} \Lambda) & \longrightarrow & \mathbf{D}^b(\mathrm{mod} \Lambda) \\ \downarrow & & \downarrow \\ \mathbf{D}(\mathrm{Mod} \Lambda) & \xrightarrow{Q_\lambda} & \mathbf{K}(\mathrm{Inj} \Lambda) \end{array}$$

Here, $\mathrm{Flat} \Lambda$ denotes the full subcategory of flat Λ -modules, which is the closure of $\mathrm{proj} \Lambda$ under forming filtered colimits.

Example 4.8. Let \mathbb{X} be a quasi-compact and separated scheme, and let L be a locally free sheaf of finite rank. Then

$$\mathrm{Hom}_{\mathbf{D}(\mathrm{Qcoh} \mathbb{X})}(L, -) \cong \mathrm{Hom}_{\mathbf{D}(\mathrm{Qcoh} \mathbb{X})}(\mathcal{O}_{\mathbb{X}}, L^\vee \otimes_{\mathcal{O}_{\mathbb{X}}} -) \cong H^0(L^\vee \otimes_{\mathcal{O}_{\mathbb{X}}} -),$$

where $L^\vee = \mathcal{H}om_{\mathcal{O}_{\mathbb{X}}}(L, \mathcal{O}_{\mathbb{X}})$. Thus L is a compact object in $\mathbf{D}(\mathrm{Qcoh} \mathbb{X})$; see [26]. If \mathbb{X} has an ample family of line bundles, then the locally free sheaves of finite rank generate $\mathrm{Qcoh} \mathbb{X}$.

Our final example shows that products in $\mathbf{Qcoh} \mathbb{X}$ need not to be exact. We include this as an illustration for Corollary 4.4

Example 4.9. Let k be a field and $\mathbb{X} = \mathbb{P}_k^1$ the projective line with homogeneous coordinate ring $S = k[x_0, x_1]$. For each $n \in \mathbb{Z}$, we have an exact sequence

$$S(n-1) \amalg S(n-1) \longrightarrow S(n) \longrightarrow k(n) \longrightarrow 0$$

of S -modules which gives rise to an epimorphism $\mathcal{O}(n-1) \amalg \mathcal{O}(n-1) \rightarrow \mathcal{O}(n)$ in $\mathbf{Qcoh} \mathbb{X}$. We claim that the product

$$\pi: \prod_{n \in \mathbb{Z}} (\mathcal{O}(n-1) \amalg \mathcal{O}(n-1)) \longrightarrow \prod_{n \in \mathbb{Z}} \mathcal{O}(n)$$

is not an epimorphism. In fact, the cokernel of π is isomorphic to the sheaf corresponding to the module $\prod_{n \in \mathbb{Z}} k(n)$. But this module is not torsion. For instance, it contains $\prod_{n \geq 0} k(n)$ as a finitely generated submodule which is not of finite length. We conclude that π has a non-trivial cokernel.

5. THE STABLE DERIVED CATEGORY

Let \mathcal{A} be a locally noetherian Grothendieck category. We suppose that $\mathbf{D}(\mathcal{A})$ is compactly generated.

Definition 5.1. The *stable derived category* $\mathbf{S}(\mathcal{A})$ of \mathcal{A} is by definition the full subcategory of $\mathbf{K}(\mathcal{A})$ which is formed by all acyclic complexes of injective objects in \mathcal{A} . The full subcategory of compact objects is denoted by $\mathbf{S}^c(\mathcal{A})$.

In this section, we show that the stable derived category is compactly generated, and the description of the category of compact objects justifies our terminology. Our basic tool is the (co)localization sequence

$$\mathbf{S}(\mathcal{A}) \xrightarrow{I} \mathbf{K}(\mathrm{Inj} \mathcal{A}) \xrightarrow{Q} \mathbf{D}(\mathcal{A}) .$$

Thus we use the fact that I and Q have left adjoints I_λ, Q_λ and right adjoints I_ρ, Q_ρ . The *stabilization functor* is by definition the composite

$$S: \mathbf{D}(\mathcal{A}) \xrightarrow{I_\lambda \circ Q_\rho} \mathbf{S}(\mathcal{A}).$$

We begin with the following lemma.

Lemma 5.2. *Let \mathcal{A} be a locally noetherian Grothendieck category and suppose $\mathbf{D}(\mathcal{A})$ is compactly generated. The functors $Q_\lambda, Q_\rho: \mathbf{D}(\mathcal{A}) \rightarrow \mathbf{K}(\mathrm{Inj} \mathcal{A})$ admit a natural transformation $\eta: Q_\lambda \rightarrow Q_\rho$, and η is an isomorphism when restricted to the subcategory of compact objects in $\mathbf{D}(\mathcal{A})$.*

Proof. Each object in $\mathbf{D}(\mathcal{A})$ is isomorphic to one in the image of Q , and each map $QX \rightarrow QY$ is a fraction of maps in the image of Q . Thus the natural transformation

$$Q_\lambda \circ Q \longrightarrow \mathrm{Id}_{\mathbf{K}(\mathrm{Inj} \mathcal{A})} \longrightarrow Q_\rho \circ Q$$

induces a natural transformation $\eta: Q_\lambda \rightarrow Q_\rho$. Now observe that $Q(\eta)$ induces an isomorphism

$$Q \circ Q_\lambda \xrightarrow{\sim} Q \circ Q_\rho.$$

We know from Proposition 2.3 that Q induces an equivalence

$$\mathbf{K}^c(\mathrm{Inj} \mathcal{A}) \xrightarrow{\sim} \mathbf{D}^b(\mathrm{noeth} \mathcal{A}).$$

On the other hand,

$$Q_\lambda(\mathbf{D}^c(\mathcal{A})) \subseteq \mathbf{K}^c(\mathrm{Inj} \mathcal{A})$$

since a left adjoint preserves compactness if the right adjoint preserves coproducts; see Lemma 3.5. Also,

$$Q_\rho(\mathbf{D}^c(\mathcal{A})) \subseteq \mathbf{K}^c(\mathrm{Inj} \mathcal{A}),$$

since $\mathbf{D}^c(\mathcal{A}) \subseteq \mathbf{D}^b(\mathrm{noeth} \mathcal{A})$ by Lemma 4.1, and

$$Q_\rho(\mathbf{D}^b(\mathrm{noeth} \mathcal{A})) = \mathbf{K}^c(\mathrm{Inj} \mathcal{A})$$

by Remark 3.8. We conclude that $\eta|_{\mathbf{D}^c(\mathcal{A})}$ is an isomorphism. \square

Proposition 5.3. *Let \mathcal{A} be a locally noetherian Grothendieck category, and suppose $\mathbf{D}(\mathcal{A})$ is compactly generated. Then we have a localization sequence*

$$(5.1) \quad \mathbf{D}(\mathcal{A}) \xrightarrow{Q_\lambda} \mathbf{K}(\mathrm{Inj} \mathcal{A}) \xrightarrow{I_\lambda} \mathbf{S}(\mathcal{A})$$

which induces the following commutative diagram.

$$\begin{array}{ccccc} \mathbf{D}^c(\mathcal{A}) & \xrightarrow{\mathrm{inc}} & \mathbf{D}^b(\mathrm{noeth} \mathcal{A}) & \xrightarrow{\mathrm{can}} & \mathbf{D}^b(\mathrm{noeth} \mathcal{A})/\mathbf{D}^c(\mathcal{A}) \\ \parallel & & \downarrow \wr Q_\rho|_{\mathbf{D}^b(\mathrm{noeth} \mathcal{A})} & & \downarrow F \\ \mathbf{D}^c(\mathcal{A}) & \longrightarrow & \mathbf{K}^c(\mathrm{Inj} \mathcal{A}) & \longrightarrow & \mathbf{S}^c(\mathcal{A}) \\ \downarrow \mathrm{inc} & & \downarrow \mathrm{inc} & & \downarrow \mathrm{inc} \\ \mathbf{D}(\mathcal{A}) & \xrightarrow{Q_\lambda} & \mathbf{K}(\mathrm{Inj} \mathcal{A}) & \xrightarrow{I_\lambda} & \mathbf{S}(\mathcal{A}) \end{array}$$

Proof. It follows from Theorem 4.2 that the sequence (5.1) is a localization sequence. Let us explain the commutativity of the diagram. First observe that a left adjoint preserves compactness if the right adjoint preserves coproducts; see Lemma 3.5. Therefore I_λ and Q_λ preserve compactness, and this explains the commutativity of the lower squares. Now observe that

$$\mathbf{D}^c(\mathcal{A}) \subseteq \mathbf{D}^b(\mathrm{noeth} \mathcal{A}),$$

by Lemma 4.1, and that $Q_\rho|_{\mathbf{D}^b(\mathrm{noeth} \mathcal{A})}$ is a quasi-inverse for $Q|_{\mathbf{K}^c(\mathrm{Inj} \mathcal{A})}$. It follows from Lemma 5.2 that the upper left hand square commutes. The functor F is by definition the unique functor making the upper right hand square commutative. It exists because $I_\lambda \circ Q_\lambda = 0$. \square

We have seen that the stable derived category $\mathbf{S}(\mathcal{A})$ is a localization of the homotopy category $\mathbf{K}(\mathrm{Inj} \mathcal{A})$. This has some interesting consequences.

Corollary 5.4. *The stable derived category $\mathbf{S}(\mathcal{A})$ is compactly generated, and the functor $I_\lambda \circ Q_\rho: \mathbf{D}(\mathcal{A}) \rightarrow \mathbf{S}(\mathcal{A})$ induces (up to direct factors) an equivalence*

$$F: \mathbf{D}^b(\mathrm{noeth} \mathcal{A})/\mathbf{D}^c(\mathcal{A}) \xrightarrow{\sim} \mathbf{S}^c(\mathcal{A}).$$

Proof. We know from Proposition 2.3 that $\mathbf{K}(\text{Inj } \mathcal{A})$ is compactly generated. This property carries over to $\mathbf{S}(\mathcal{A})$ since I_λ sends a set of compact generators of $\mathbf{K}(\text{Inj } \mathcal{A})$ to a set of compact generators of $\mathbf{S}(\mathcal{A})$. The functor Q_λ identifies $\mathbf{D}(\mathcal{A})$ with the localizing subcategory of $\mathbf{K}(\text{Inj } \mathcal{A})$ which is generated by all compact objects in the image of Q_λ . Now apply the localization theorem of Neeman-Ravenel-Thomason-Trobaugh-Yao [25]. This result describes the category of compact objects of the quotient $\mathbf{S}(\mathcal{A})$, up to direct factors, as the quotient of the compact objects in $\mathbf{K}(\text{Inj } \mathcal{A})$ modulo those from the localizing subcategory. To be precise, F is fully faithful and every object in $\mathbf{S}^c(\mathcal{A})$ is a direct factor of some object in the image of F . \square

Corollary 5.5. *The composite*

$$\mathcal{A} \xrightarrow{\text{can}} \mathbf{D}(\mathcal{A}) \xrightarrow{I_\lambda \circ Q_\rho} \mathbf{S}(\mathcal{A})$$

preserves all coproducts and annihilates the objects in $\mathcal{A} \cap \mathbf{D}^c(\mathcal{A})$.

Proof. The diagram in Proposition 5.3 shows that $I_\lambda \circ Q_\rho$ annihilates $\mathcal{A} \cap \mathbf{D}^c(\mathcal{A})$. To show that $I_\lambda \circ Q_\rho$ preserves all coproducts, observe that Q_ρ sends an object in \mathcal{A} to an injective resolution. A coproduct of injective resolutions is again an injective resolution, and the left adjoint I_λ preserves all coproducts. This finishes the proof. \square

Using the stabilization functor $S: \mathbf{D}(\mathcal{A}) \rightarrow \mathbf{S}(\mathcal{A})$, we define for objects X, Y in $\mathbf{D}(\mathcal{A})$ and $n \in \mathbb{Z}$ the *stable cohomology group*

$$\underline{\text{Ext}}_{\mathcal{A}}^n(X, Y) = \text{Hom}_{\mathbf{K}(\mathcal{A})}(SX, \Sigma^n(SY)).$$

Note that in both arguments each exact sequence in \mathcal{A} induces a long exact sequence in stable cohomology. We do not go into details but refer to our discussion of Tate cohomology in Section 7. In fact, both cohomology theories coincide in case \mathcal{A} satisfies some appropriate Gorenstein property, and we shall see explicit formulae for the Tate cohomology groups.

Example 5.6. Suppose \mathcal{A} is a module category. Then the stabilization functor annihilates all modules of finite projective dimension. Similarly, if \mathcal{A} is a category of quasi-coherent sheaves, then the stabilization functor annihilates all sheaves having a finite resolution with locally free sheaves.

Given a noetherian scheme \mathbb{X} , the stable derived category $\mathbf{S}(\text{Qcoh } \mathbb{X})$ vanishes if \mathbb{X} is regular. Nonetheless, a classical result of Bernstein-Gelfand-Gelfand [8] shows that stable derived categories are relevant when one studies regular schemes. This is sketched in the following example.

Example 5.7. Let Λ be a Koszul algebra and $\Lambda^!$ its Koszul dual. Then we have under appropriate assumptions an equivalence $\mathbf{K}(\text{Inj } \Lambda) \xrightarrow{\sim} \mathbf{K}(\text{Inj } \Lambda^!)$ which induces an equivalence $\mathbf{D}^b(\text{mod } \Lambda) \xrightarrow{\sim} \mathbf{D}^b(\text{mod } \Lambda^!)$ when restricted to the full subcategories of compact objects [7, 22]. Note that we consider the categories of graded modules over Λ and $\Lambda^!$ respectively. The classical example is the symmetric algebra $\Lambda = SV$ of a d -dimensional space V over a field k , where $\Lambda^!$ is the exterior algebra $\bigwedge V^*$ of the dual space V^* . The equivalence $\mathbf{K}(\text{Inj } \Lambda) \xrightarrow{\sim} \mathbf{K}(\text{Inj } \Lambda^!)$ takes an injective resolution ik of $\Lambda_0 = k$ to $\Lambda^!$ and identifies the localizing subcategory \mathcal{K} generated by ik with the localizing subcategory generated by $\Lambda^!$, which is $\mathbf{D}(\text{Mod } \Lambda^!)$. Note that the quotient $\mathbf{K}(\text{Inj } \Lambda)/\mathcal{K}$ identifies with the derived category of the quotient $\text{Mod } \Lambda/(\text{Mod } \Lambda)_0$, where $(\text{Mod } \Lambda)_0$ denotes

the subcategory of torsion modules. This quotient is equivalent to $\mathrm{Qcoh} \mathbb{P}_k^{d-1}$ by Serre's Theorem. Thus we obtain an equivalence

$$\mathbf{D}(\mathrm{Qcoh} \mathbb{P}_k^{d-1}) \xrightarrow{\sim} \mathbf{S}(\mathrm{Mod} \wedge k^d).$$

Note that $\mathbf{S}(\mathrm{Mod} \wedge k^d)$ is equivalent to the stable module category $\underline{\mathrm{Mod}} \wedge k^d$ because the exterior algebra is self-injective; see Example 7.15. Passing to the subcategory of compact objects, one obtains the equivalence

$$\mathbf{D}^b(\mathrm{coh} \mathbb{P}_k^{d-1}) \xrightarrow{\sim} \underline{\mathrm{mod}} \wedge k^d$$

of Bernstein-Gelfand-Gelfand [8], where $\underline{\mathrm{mod}} \wedge k^d$ denotes the stable category of all finite dimensional $\wedge k^d$ -modules. This example generalizes to non-commutative algebras, for instance, to Artin-Schelter regular algebras [21].

6. EXTENDING DERIVED FUNCTORS

An additive functor $F: \mathcal{A} \rightarrow \mathcal{B}$ between locally noetherian Grothendieck categories admits a right derived functor $\mathbf{R}F: \mathbf{D}(\mathcal{A}) \rightarrow \mathbf{D}(\mathcal{B})$. In this section we extend this to a functor $\hat{\mathbf{R}}F: \mathbf{K}(\mathrm{Inj} \mathcal{A}) \rightarrow \mathbf{K}(\mathrm{Inj} \mathcal{B})$ and investigate its right and left adjoints. As an application, we consider for F the direct image functor $f_*: \mathrm{Qcoh} \mathbb{X} \rightarrow \mathrm{Qcoh} \mathbb{Y}$ corresponding to a morphism $f: \mathbb{X} \rightarrow \mathbb{Y}$ between noetherian schemes. We use the following functors

$$I: \mathbf{K}(\mathrm{Inj} \mathcal{A}) \xrightarrow{\mathrm{inc}} \mathbf{K}(\mathcal{A}) \quad \text{and} \quad Q: \mathbf{K}(\mathrm{Inj} \mathcal{A}) \xrightarrow{\mathrm{inc}} \mathbf{K}(\mathcal{A}) \xrightarrow{\mathrm{can}} \mathbf{D}(\mathcal{A})$$

simultaneously for \mathcal{A} and \mathcal{B} . Moreover, we use the fact that both functors have left and right adjoints.

Theorem 6.1. *Let $F: \mathcal{A} \rightarrow \mathcal{B}$ be an additive functor between locally noetherian Grothendieck categories. Suppose $\mathbf{D}(\mathcal{A})$ and $\mathbf{D}(\mathcal{B})$ are compactly generated. Then the composite*

$$\hat{\mathbf{R}}F: \mathbf{K}(\mathrm{Inj} \mathcal{A}) \xrightarrow{I} \mathbf{K}(\mathcal{A}) \xrightarrow{\mathbf{K}(F)} \mathbf{K}(\mathcal{B}) \xrightarrow{I_\lambda} \mathbf{K}(\mathrm{Inj} \mathcal{B})$$

makes the following diagram commutative.

$$\begin{array}{ccc} \mathbf{D}(\mathcal{A}) & \xrightarrow{\mathbf{R}F} & \mathbf{D}(\mathcal{B}) \\ \downarrow Q_\rho & & \uparrow Q \\ \mathbf{K}(\mathrm{Inj} \mathcal{A}) & \xrightarrow{\hat{\mathbf{R}}F} & \mathbf{K}(\mathrm{Inj} \mathcal{B}) \end{array}$$

- (1) *Suppose F preserves coproducts. Then $\hat{\mathbf{R}}F$ preserves coproducts and has therefore a right adjoint $(\hat{\mathbf{R}}F)_\rho$.*
- (2) *Suppose F and $\mathbf{R}F$ preserve coproducts. Then $\mathbf{R}F$ has a right adjoint $(\mathbf{R}F)_\rho$ making the following diagram commutative.*

$$\begin{array}{ccc} \mathbf{D}(\mathcal{A}) & \xrightarrow{\mathbf{R}F} & \mathbf{D}(\mathcal{B}) & \xrightarrow{(\mathbf{R}F)_\rho} & \mathbf{D}(\mathcal{A}) \\ \downarrow Q_\lambda & & \uparrow Q & & \downarrow Q_\rho \\ \mathbf{K}(\mathrm{Inj} \mathcal{A}) & \xrightarrow{\hat{\mathbf{R}}F} & \mathbf{K}(\mathrm{Inj} \mathcal{B}) & \xrightarrow{(\hat{\mathbf{R}}F)_\rho} & \mathbf{K}(\mathrm{Inj} \mathcal{A}) \end{array}$$

Proof. The composite

$$\mathbf{K}(\mathcal{A}) \xrightarrow{I_\lambda} \mathbf{K}(\mathrm{Inj} \mathcal{A}) \xrightarrow{Q} \mathbf{D}(\mathcal{A})$$

is naturally isomorphic to the canonical functor $\mathbf{K}(\mathcal{A}) \rightarrow \mathbf{D}(\mathcal{A})$; see Remark 3.7. Clearly, $I \circ Q_\rho$ is its right adjoint. We denote by $\mathbf{R}F$ the right derived functor of F and have

$$\mathbf{R}F = Q \circ I_\lambda \circ \mathbf{K}(F) \circ I \circ Q_\rho.$$

Using the definition $\hat{\mathbf{R}}F = I_\lambda \circ \mathbf{K}(F) \circ I$, we obtain $\mathbf{R}F = Q \circ \hat{\mathbf{R}}F \circ Q_\rho$.

(1) Suppose F preserves coproducts. Then $\mathbf{K}(F)$ preserves coproducts. It follows that $\hat{\mathbf{R}}F$ preserves coproducts since I and I_λ preserve coproducts. Now apply Proposition 3.3 to obtain a right adjoint for $\hat{\mathbf{R}}F$.

(2) Suppose F and $\mathbf{R}F$ preserve coproducts. Then $\mathbf{R}F$ has a right adjoint by Proposition 3.3. Next we show that

$$Q \circ \hat{\mathbf{R}}F \circ Q_\lambda \cong Q \circ \hat{\mathbf{R}}F \circ Q_\rho.$$

We have a natural transformation $Q_\lambda \rightarrow Q_\rho$ which is induced from the natural transformation $Q_\lambda \circ Q \rightarrow Q_\rho \circ Q$. Now apply $Q \circ \hat{\mathbf{R}}F$ to get a natural transformation

$$\mu: Q \circ \hat{\mathbf{R}}F \circ Q_\lambda \longrightarrow Q \circ \hat{\mathbf{R}}F \circ Q_\rho.$$

It is shown in Lemma 5.2 that $Q_\lambda \rightarrow Q_\rho$ is an isomorphism when restricted to compact objects in $\mathbf{D}(\mathcal{A})$. On the other hand, $Q \circ \hat{\mathbf{R}}F \circ Q_\lambda$ and $Q \circ \hat{\mathbf{R}}F \circ Q_\rho$ both preserve coproducts by our assumption on $\mathbf{R}F$. It follows that μ is an isomorphism since $\mathbf{D}(\mathcal{A})$ is compactly generated. Clearly, $Q \circ (\hat{\mathbf{R}}F)_\rho \circ Q_\rho$ is a right adjoint for $Q \circ \hat{\mathbf{R}}F \circ Q_\lambda$. This completes the proof. \square

The extended derived functor and its right adjoint admit some alternative description. I am indebted to Bernhard Keller for providing this remark.

Remark 6.2. It is possible to express $\hat{\mathbf{R}}F$ as the tensor functor and its right adjoint $(\hat{\mathbf{R}}F)_\rho$ as the Hom functor with respect to a bimodule of DG categories; see [22, 6.4]. This depends on the appropriate choice of DG categories \mathcal{A}_0 and \mathcal{B}_0 such that $\mathbf{K}(\mathrm{Inj} \mathcal{A}) \cong \mathbf{D}_{\mathrm{dg}}(\mathcal{A}_0)$ and $\mathbf{K}(\mathrm{Inj} \mathcal{B}) \cong \mathbf{D}_{\mathrm{dg}}(\mathcal{B}_0)$ respectively.

Next we consider as an example a morphism $f: \mathbb{X} \rightarrow \mathbb{Y}$ between separated noetherian schemes. Let $f_*: \mathrm{Qcoh} \mathbb{X} \rightarrow \mathrm{Qcoh} \mathbb{Y}$ denote the direct image functor. Note that the right derived functor $\mathbf{R}f_*: \mathbf{D}(\mathrm{Qcoh} \mathbb{X}) \rightarrow \mathbf{D}(\mathrm{Qcoh} \mathbb{Y})$ preserves coproducts [26, Lemma 1.4]. Thus $\mathbf{R}f_*$ and its right adjoint Grothendieck duality functor $f^!$ extend to functors between $\mathbf{K}(\mathrm{Inj} \mathbb{X})$ and $\mathbf{K}(\mathrm{Inj} \mathbb{Y})$, by Theorem 6.1. This is the statement of Theorem 1.4 from the introduction. In fact, the situation is in this case much nicer. I am grateful to Amnon Neeman for pointing out that the functor $\hat{\mathbf{R}}f_*$ and its right adjoint $(\hat{\mathbf{R}}f_*)_\rho$ make the following diagram commutative.

$$(6.1) \quad \begin{array}{ccc} \mathbf{K}(\mathrm{Inj} \mathbb{X}) & \xrightarrow{\hat{\mathbf{R}}f_*} & \mathbf{K}(\mathrm{Inj} \mathbb{Y}) \\ \downarrow Q & & \downarrow Q \\ \mathbf{D}(\mathrm{Qcoh} \mathbb{X}) & \xrightarrow{\mathbf{R}f_*} & \mathbf{D}(\mathrm{Qcoh} \mathbb{Y}) \end{array} \quad \begin{array}{ccc} \mathbf{K}(\mathrm{Inj} \mathbb{Y}) & \xrightarrow{(\hat{\mathbf{R}}f_*)_\rho} & \mathbf{K}(\mathrm{Inj} \mathbb{X}) \\ \uparrow Q_\rho & & \uparrow Q_\rho \\ \mathbf{D}(\mathrm{Qcoh} \mathbb{Y}) & \xrightarrow{f^!} & \mathbf{D}(\mathrm{Qcoh} \mathbb{X}) \end{array}$$

This is essentially the statement of Theorem 1.5 from the introduction. The proof which is due to Neeman uses the following lemma.

Lemma 6.3. *Keep the assumptions from Theorem 6.1 and consider the following diagram.*

$$\begin{array}{ccc} \mathbf{K}(\mathrm{Inj} \mathcal{A}) & \xrightarrow{\hat{\mathbf{R}}F} & \mathbf{K}(\mathrm{Inj} \mathcal{B}) \\ \downarrow Q & & \downarrow Q \\ \mathbf{D}(\mathcal{A}) & \xrightarrow{\mathbf{R}F} & \mathbf{D}(\mathcal{B}) \end{array}$$

There is a natural transformation $Q \circ \hat{\mathbf{R}}F \rightarrow \mathbf{R}F \circ Q$ which is an isomorphism if and only if F sends every acyclic complex of injective objects to an acyclic complex.

Proof. We apply the the localization sequence

$$\mathbf{K}_{\mathrm{ac}}(\mathrm{Inj} \mathcal{A}) \xrightarrow{I} \mathbf{K}(\mathrm{Inj} \mathcal{A}) \xrightarrow{Q} \mathbf{D}(\mathcal{A})$$

from Proposition 3.6. Let X be an object in $\mathbf{K}(\mathrm{Inj} \mathcal{A})$ and consider the triangle

$$(I \circ I_\rho)X \rightarrow X \rightarrow (Q_\rho \circ Q)X \rightarrow \Sigma(I \circ I_\rho)X$$

in $\mathbf{K}(\mathrm{Inj} \mathcal{A})$. Now apply $Q \circ \hat{\mathbf{R}}F$ which gives a map

$$(Q \circ \hat{\mathbf{R}}F)X \rightarrow (\mathbf{R}F \circ Q)X$$

since $Q \circ \hat{\mathbf{R}}F \circ Q_\rho \cong \mathbf{R}F$, by Theorem 6.1. Clearly, this map is an isomorphism if and only if $Q \circ \hat{\mathbf{R}}F$ annihilates $(I \circ I_\rho)X$. \square

We include a simple example which illustrates the preceding lemma.

Example 6.4. Let k be a field and $\Lambda = k[t]/(t^2)$. We take the functor

$$F: \mathrm{Mod} \Lambda \rightarrow \mathrm{Mod} k, \quad X \mapsto \mathrm{Hom}_\Lambda(k, X),$$

and observe that the following diagram does not commute.

$$\begin{array}{ccc} \mathbf{K}(\mathrm{Inj} \Lambda) & \xrightarrow{\hat{\mathbf{R}}F} & \mathbf{K}(\mathrm{Inj} k) \\ \downarrow Q & & \downarrow Q \\ \mathbf{D}(\mathrm{Mod} \Lambda) & \xrightarrow{\mathbf{R}F} & \mathbf{D}(\mathrm{Mod} k) \end{array}$$

For instance, we have $QX = 0$ and $(\hat{\mathbf{R}}F)X \neq 0$ if we take for X the acyclic complex

$$\cdots \xrightarrow{t} \Lambda \xrightarrow{t} \Lambda \xrightarrow{t} \Lambda \xrightarrow{t} \cdots$$

in $\mathbf{K}(\mathrm{Inj} \Lambda)$.

Proof of Theorem 1.5. We need to show that both squares in (6.1) commute. Then we use the localization sequence

$$\mathbf{S}(\mathrm{Qcoh} \mathbb{X}) \xrightarrow{I} \mathbf{K}(\mathrm{Inj} \mathbb{X}) \xrightarrow{Q} \mathbf{D}(\mathrm{Qcoh} \mathbb{X})$$

from Proposition 3.6 and obtain from $\hat{\mathbf{R}}f_*$ and $(\hat{\mathbf{R}}f_*)_\rho$ an adjoint pair of functors between $\mathbf{S}(\mathrm{Qcoh} \mathbb{X})$ and $\mathbf{S}(\mathrm{Qcoh} \mathbb{Y})$.

In order to show the commutativity of (6.1), we apply Lemma 6.3 and need to show that f_* sends an acyclic complex X of injective objects to an acyclic complex. The question is local in \mathbb{Y} and we may assume \mathbb{Y} affine. Cover \mathbb{X} by a finite number of

affines. Then f_* can be computed using the Čech cohomology of the cover. If there are n open sets in the cover, then for any quasi-coherent sheaf A we have

$$\mathbf{R}^{n+1}f_*A = 0.$$

Now take our acyclic complex X of injective sheaves on \mathbb{X} . Then the sequence

$$0 \longrightarrow X^0 \longrightarrow X^1 \longrightarrow X^2 \longrightarrow \dots$$

is an injective resolution of the kernel A of the map $X^0 \rightarrow X^1$. Applying f_* , the sequence computes for us $\mathbf{R}^i f_* A$, which vanishes if $i \geq n + 1$. Thus $f_* X$ is acyclic above degree n , but by shifting we conclude that it is acyclic everywhere.

Having shown the commutativity of the left hand square, the commutativity of the right hand square follows, because it is obtained by taking right adjoints. Thus the proof is complete. \square

Next we investigate for an exact functor $F: \mathcal{A} \rightarrow \mathcal{B}$ an extension $\hat{\mathbf{L}}F$ of the derived functor $\mathbf{L}F: \mathbf{D}(\mathcal{A}) \rightarrow \mathbf{D}(\mathcal{B})$. For this we need some assumptions, and it is convenient to introduce the following notation. As before, \mathcal{A} and \mathcal{B} denote locally noetherian Grothendieck categories. Let $f: \text{noeth } \mathcal{A} \rightarrow \text{noeth } \mathcal{B}$ be an additive functor. Then there is, up to isomorphism, a unique functor $f^*: \mathcal{A} \rightarrow \mathcal{B}$ which extends f and preserves filtered colimits. This has a right adjoint $f_*: \mathcal{B} \rightarrow \mathcal{A}$ if and only if f is right exact. Note that f is exact iff f^* is exact iff f_* sends injective objects to injective objects. Here is an example.

Example 6.5. Let $f: \mathbb{X} \rightarrow \mathbb{Y}$ be a morphism between noetherian schemes. Then the inverse image functor $f^*: \text{Qcoh } \mathbb{Y} \rightarrow \text{Qcoh } \mathbb{X}$ sends coherent sheaves to coherent sheaves and preserves filtered colimits. Moreover, the direct image functor f_* is a right adjoint of f^* . Our notation is therefore consistent if we identify the morphism $f: \mathbb{X} \rightarrow \mathbb{Y}$ with the functor $\text{coh } \mathbb{Y} \rightarrow \text{coh } \mathbb{X}$.

Theorem 6.6. *Let \mathcal{A} and \mathcal{B} locally noetherian Grothendieck categories such that $\mathbf{D}(\mathcal{A})$ and $\mathbf{D}(\mathcal{B})$ are compactly generated. Let $f: \text{noeth } \mathcal{A} \rightarrow \text{noeth } \mathcal{B}$ be an exact functor. Then $\hat{\mathbf{R}}f_*$ has a left adjoint $\hat{\mathbf{L}}f^*$ which induces a functor $\mathbf{S}f^*$ making the following diagram commutative.*

$$(6.2) \quad \begin{array}{ccccc} & & \mathbf{K}^c(\text{Inj } \mathcal{A}) & \xrightarrow{\sim} & \mathbf{D}^b(\text{noeth } \mathcal{A}) \\ & & \downarrow & & \downarrow \mathbf{D}^b(f) \\ \mathbf{S}(\mathcal{A}) & \xrightarrow{I} & \mathbf{K}(\text{Inj } \mathcal{A}) & \xrightarrow{Q} & \mathbf{D}(\mathcal{A}) \\ \downarrow \mathbf{S}f^* & & \downarrow \hat{\mathbf{L}}f^* & & \downarrow \mathbf{L}f^* \\ & & \mathbf{K}^c(\text{Inj } \mathcal{B}) & \xrightarrow{\sim} & \mathbf{D}^b(\text{noeth } \mathcal{B}) \\ \mathbf{S}(\mathcal{B}) & \xrightarrow{I} & \mathbf{K}(\text{Inj } \mathcal{B}) & \xrightarrow{Q} & \mathbf{D}(\mathcal{B}) \end{array}$$

If $\mathbf{R}f_*$ preserves coproducts, then in addition the following diagram commutes.

$$\begin{array}{ccccc}
 & \mathbf{D}^b(\text{noeth } \mathcal{A})/\mathbf{D}^c(\mathcal{A}) & \longleftarrow & \mathbf{K}^c(\text{Inj } \mathcal{A}) & \longleftarrow & \mathbf{D}^c(\mathcal{A}) \\
 & \swarrow F_{\mathcal{A}} & \downarrow & \swarrow & \downarrow & \swarrow \\
 \mathbf{S}(\mathcal{A}) & \xleftarrow{I_{\lambda}} & \mathbf{K}(\text{Inj } \mathcal{A}) & \xleftarrow{Q_{\lambda}} & \mathbf{D}(\mathcal{A}) & \xleftarrow{\quad} & \mathbf{D}^c(\mathcal{A}) \\
 & \downarrow \overline{\mathbf{D}^b(f)} & \downarrow \hat{\mathbf{L}}f^* & \downarrow & \downarrow \mathbf{L}f^* & \downarrow & \downarrow \\
 & \mathbf{D}^b(\text{noeth } \mathcal{B})/\mathbf{D}^c(\mathcal{B}) & \longleftarrow & \mathbf{K}^c(\text{Inj } \mathcal{B}) & \longleftarrow & \mathbf{D}^c(\mathcal{B}) \\
 & \swarrow F_{\mathcal{B}} & \downarrow & \swarrow & \downarrow & \swarrow \\
 \mathbf{S}(\mathcal{B}) & \xleftarrow{I_{\lambda}} & \mathbf{K}(\text{Inj } \mathcal{B}) & \xleftarrow{Q_{\lambda}} & \mathbf{D}(\mathcal{B}) & \xleftarrow{\quad} & \mathbf{D}^c(\mathcal{B}) \\
 & \downarrow \mathbf{S}f^* & & & & & \\
 & & & & & &
 \end{array}$$

Note that $F_{\mathcal{A}}$ and $F_{\mathcal{B}}$ induce, up to direct factors, equivalences onto the full subcategories of compact objects in $\mathbf{S}(\mathcal{A})$ and $\mathbf{S}(\mathcal{B})$ respectively. Thus $\overline{\mathbf{D}^b(f)}$ determines the functor $\mathbf{S}f^*$.

Proof. The exactness of f implies the exactness of f^* . Thus f_* sends injective objects to injective objects and we have the following commutative diagram.

$$(6.3) \quad \begin{array}{ccc}
 \mathbf{K}(\text{Inj } \mathcal{B}) & \xrightarrow{J} & \mathbf{K}(\mathcal{B}) \\
 \downarrow \hat{\mathbf{R}}f_* & & \downarrow \mathbf{K}(f_*) \\
 \mathbf{K}(\text{Inj } \mathcal{B}) & \xrightarrow{J} & \mathbf{K}(\mathcal{B})
 \end{array}$$

The right adjoint f_* preserves products and we have therefore a left adjoint for $\hat{\mathbf{R}}f_*$, by Proposition 3.3, which we denote by $\hat{\mathbf{L}}f^*$. We obtain the following diagram

$$\begin{array}{ccccc}
 \mathbf{K}(\mathcal{A}) & \xrightarrow{J_{\lambda}} & \mathbf{K}(\text{Inj } \mathcal{A}) & \xrightarrow{Q} & \mathbf{D}(\mathcal{A}) \\
 \downarrow \mathbf{K}(f^*) & & \downarrow \hat{\mathbf{L}}f^* & & \downarrow \mathbf{L}f^* \\
 \mathbf{K}(\mathcal{B}) & \xrightarrow{J_{\lambda}} & \mathbf{K}(\text{Inj } \mathcal{B}) & \xrightarrow{Q} & \mathbf{D}(\mathcal{B})
 \end{array}$$

and claim it is commutative. The left hand square commutes because it is obtained from (6.3) by taking left adjoints. The outer square commutes because the composite $Q \circ J_{\lambda}$ is naturally isomorphic to the canonical functor $\mathbf{K}(\mathcal{A}) \rightarrow \mathbf{D}(\mathcal{A})$; see Remark 3.7. We conclude the commutativity of the right hand square, using that $J_{\lambda} \circ J \cong \text{Id}_{\mathbf{K}(\text{Inj } \mathcal{A})}$. Clearly, $\hat{\mathbf{L}}f^*$ sends acyclic complexes to acyclic complexes and we obtain the functor $\mathbf{S}f^*$ making the diagram (6.2) commutative.

Now assume in addition that $\mathbf{R}f_*$ preserves coproducts. We use the fact that a left adjoint preserves compactness if the right adjoint preserves coproducts; see Lemma 3.5. The functor $\hat{\mathbf{R}}f_*$ preserves coproducts since f_* preserves coproducts. Thus $\mathbf{L}f^*$ and $\hat{\mathbf{L}}f^*$ preserve compactness. Note that Q_{λ} identifies $\mathbf{D}(\mathcal{A})$ with the localizing subcategory of $\mathbf{K}(\text{Inj } \mathcal{A})$ which is generated by

$$\mathbf{D}^c(\mathcal{A}) \subseteq \mathbf{D}^b(\text{noeth } \mathcal{A}) \cong \mathbf{K}^c(\text{Inj } \mathcal{A}) \subseteq \mathbf{K}(\text{Inj } \mathcal{A}).$$

Of course, the same applies for \mathcal{B} . We obtain the following diagram

$$\begin{array}{ccccc} \mathbf{S}(\mathcal{A}) & \xleftarrow{I_\lambda} & \mathbf{K}(\text{Inj } \mathcal{A}) & \xleftarrow{Q_\lambda} & \mathbf{D}(\mathcal{A}) \\ \downarrow S & & \downarrow \hat{\mathbf{L}}f^* & & \downarrow \mathbf{L}f^* \\ \mathbf{S}(\mathcal{B}) & \xleftarrow{I_\lambda} & \mathbf{K}(\text{Inj } \mathcal{B}) & \xleftarrow{Q_\lambda} & \mathbf{D}(\mathcal{B}) \end{array}$$

where the right hand square commutes. The horizontal sequences are localization sequences by Theorem 4.2, and $\hat{\mathbf{L}}f^*$ induces a functor $S: \mathbf{S}(\mathcal{A}) \rightarrow \mathbf{S}(\mathcal{B})$ making the left hand square commutative. Moreover, we have

$$S = S \circ I_\lambda \circ I = I_\lambda \circ \hat{\mathbf{L}}f^* \circ I = I_\lambda \circ I \circ \mathbf{S}f^* = \mathbf{S}f^*.$$

The functors $F_{\mathcal{A}}$ and $F_{\mathcal{B}}$ are both induced by I_λ , and the commutativity

$$\mathbf{S}f^* \circ F_{\mathcal{A}} = F_{\mathcal{B}} \circ \overline{\mathbf{D}^b(f)}$$

is easily checked; see Corollary 5.4. This completes the proof. \square

Remark 6.7. Let $f: \mathbb{X} \rightarrow \mathbb{Y}$ be a morphism between noetherian schemes and suppose f^* is exact. Then $\mathbf{S}f^*$ is the left adjoint of $\mathbf{S}f_*$ which appears in Theorem 1.5.

Next we investigate the inclusion $f: \mathbb{X} \rightarrow \mathbb{Y}$ of an open subscheme. In this case, the adjoint pair of functors f_* and f^* between $\text{Qcoh } \mathbb{X}$ and $\text{Qcoh } \mathbb{Y}$ restricts to an adjoint pair of functors between $\text{Inj } \mathbb{X}$ and $\text{Inj } \mathbb{Y}$; see [15, VI]. Moreover, $f^* \circ f_* \cong \text{Id}_{\text{Qcoh } \mathbb{X}}$. Thus we can identify $\hat{\mathbf{R}}f_* = f_*$ and $\hat{\mathbf{L}}f^* = f^*$. Note that both functors send acyclic complexes with injective components to acyclic complexes. This is clear for f^* because it is exact, and follows for f_* from Theorem 1.5. We denote for each sheaf A in $\text{Qcoh } \mathbb{Y}$ by $\text{Supp } A$ the support of A and observe that f^* annihilates A if and only if $\text{Supp } A$ is contained in $\mathbb{Y} \setminus \mathbb{X}$. In fact, the natural map $A \rightarrow (f_* \circ f^*)A$ induces a split exact sequence

$$0 \rightarrow A' \rightarrow A \rightarrow (f_* \circ f^*)A \rightarrow 0$$

if A is injective. In particular, the support of A' is contained in $\mathbb{Y} \setminus \mathbb{X}$.

Now fix a complex X in $\mathbf{K}(\text{Inj } \mathbb{Y})$. The *support* of X is by definition

$$\text{Supp } X = \bigcup_{n \in \mathbb{Z}} \text{Supp } X^n.$$

We write $X_{\mathbb{X}} = (f_* \circ f^*)X$, and the natural map $X \rightarrow X_{\mathbb{X}}$ induces an exact triangle

$$(6.4) \quad X_{\mathbb{Y} \setminus \mathbb{X}} \rightarrow X \rightarrow X_{\mathbb{X}} \rightarrow \Sigma(X_{\mathbb{Y} \setminus \mathbb{X}})$$

in $\mathbf{K}(\text{Inj } \mathbb{Y})$ where the support of $X_{\mathbb{Y} \setminus \mathbb{X}}$ is contained in $\mathbb{Y} \setminus \mathbb{X}$.

Lemma 6.8. *Let \mathbb{Y} be a separated noetherian scheme and $f: \mathbb{X} \rightarrow \mathbb{Y}$ be the inclusion of an open subscheme. If X is a complex in $\mathbf{K}(\text{Inj } \mathbb{Y})$, then $f^*X = 0$ if and only if X is isomorphic to a complex with support contained in $\mathbb{Y} \setminus \mathbb{X}$.*

Proof. We have $f^*X = 0$ if and only if the first map in the triangle (6.4) is an isomorphism. \square

It is well known that f^* induces an equivalence

$$\mathbf{D}(\text{Qcoh } \mathbb{Y}) / \mathbf{D}_{\mathbb{Y} \setminus \mathbb{X}}(\text{Qcoh } \mathbb{Y}) \xrightarrow{\sim} \mathbf{D}(\text{Qcoh } \mathbb{X}),$$

where $\mathbf{D}_{\mathbb{Y} \setminus \mathbb{X}}(\mathrm{Qcoh} \mathbb{Y})$ denotes the full subcategory of all complexes in $\mathbf{D}(\mathrm{Qcoh} \mathbb{Y})$ such that the support of the cohomology is contained in $\mathbb{Y} \setminus \mathbb{X}$. We obtain an analogue for $\mathbf{K}(\mathrm{Inj} \mathbb{Y})$ and $\mathbf{S}(\mathrm{Qcoh} \mathbb{Y})$ if we define

$$\begin{aligned} \mathbf{K}_{\mathbb{Y} \setminus \mathbb{X}}(\mathrm{Inj} \mathbb{Y}) &= \{X \in \mathbf{K}(\mathrm{Inj} \mathbb{Y}) \mid \mathrm{Supp} X \subseteq \mathbb{Y} \setminus \mathbb{X}\}, \\ \mathbf{S}_{\mathbb{Y} \setminus \mathbb{X}}(\mathrm{Qcoh} \mathbb{Y}) &= \{X \in \mathbf{S}(\mathrm{Qcoh} \mathbb{Y}) \mid \mathrm{Supp} X \subseteq \mathbb{Y} \setminus \mathbb{X}\}. \end{aligned}$$

Proposition 6.9. *Let \mathbb{Y} be a seperated noetherian scheme and $f: \mathbb{X} \rightarrow \mathbb{Y}$ be the inclusion of an open subscheme. Then f^* induces equivalences*

$$\begin{aligned} \mathbf{K}(\mathrm{Inj} \mathbb{Y}) / \mathbf{K}_{\mathbb{Y} \setminus \mathbb{X}}(\mathrm{Inj} \mathbb{Y}) &\xrightarrow{\sim} \mathbf{K}(\mathrm{Inj} \mathbb{X}), \\ \mathbf{S}(\mathrm{Qcoh} \mathbb{Y}) / \mathbf{S}_{\mathbb{Y} \setminus \mathbb{X}}(\mathrm{Qcoh} \mathbb{Y}) &\xrightarrow{\sim} \mathbf{S}(\mathrm{Qcoh} \mathbb{X}). \end{aligned}$$

Proof. We have $f^* \circ f_* \cong \mathrm{Id}_{\mathrm{Qcoh} \mathbb{X}}$, and this carries over to complexes of injectives. On the other hand, we have for X in $\mathbf{K}(\mathrm{Inj} \mathbb{Y})$ a natural map $X \rightarrow (f_* \circ f^*)X$ which induces an isomorphism in $\mathbf{K}(\mathrm{Inj} \mathbb{Y}) / \mathbf{K}_{\mathbb{Y} \setminus \mathbb{X}}(\mathrm{Inj} \mathbb{Y})$, by Lemma 6.8. \square

Let us give a more elaborate formulation of Proposition 6.9. The functor $\hat{\mathbf{R}}f_* = f_*: \mathbf{K}(\mathrm{Inj} \mathbb{X}) \rightarrow \mathbf{K}(\mathrm{Inj} \mathbb{Y})$ admits a left and a right adjoint. Therefore $\hat{\mathbf{R}}f_*$ induces a recollement

$$\mathbf{K}(\mathrm{Inj} \mathbb{X}) \rightleftarrows \mathbf{K}(\mathrm{Inj} \mathbb{Y}) \rightleftarrows \mathbf{K}_{\mathbb{Y} \setminus \mathbb{X}}(\mathrm{Inj} \mathbb{Y}).$$

This recollement is compatible with the recollement

$$\mathbf{S}(\mathrm{Qcoh} \mathbb{Y}) \rightleftarrows \mathbf{K}(\mathrm{Inj} \mathbb{Y}) \rightleftarrows \mathbf{D}(\mathrm{Qcoh} \mathbb{Y}),$$

and we obtain the following diagram.

$$\begin{array}{ccccc} \mathbf{S}(\mathrm{Qcoh} \mathbb{X}) & \rightleftarrows & \mathbf{K}(\mathrm{Inj} \mathbb{X}) & \rightleftarrows & \mathbf{D}(\mathrm{Qcoh} \mathbb{X}) \\ \updownarrow & & \updownarrow & & \updownarrow \\ \mathbf{S}(\mathrm{Qcoh} \mathbb{Y}) & \rightleftarrows & \mathbf{K}(\mathrm{Inj} \mathbb{Y}) & \rightleftarrows & \mathbf{D}(\mathrm{Qcoh} \mathbb{Y}) \\ \updownarrow & & \updownarrow & & \updownarrow \\ \mathbf{S}_{\mathbb{Y} \setminus \mathbb{X}}(\mathrm{Qcoh} \mathbb{Y}) & \rightleftarrows & \mathbf{K}_{\mathbb{Y} \setminus \mathbb{X}}(\mathrm{Inj} \mathbb{Y}) & \rightleftarrows & \mathbf{D}_{\mathbb{Y} \setminus \mathbb{X}}(\mathrm{Qcoh} \mathbb{Y}) \end{array}$$

In this diagram, each row and each column is a recollement. Moreover, the diagram is commutative if one restricts to arrows in south and east direction. All other commutativity relations follow by taking left adjoints or right adjoints.

Proposition 6.9 tells us precisely when the inclusion of a subscheme induces an equivalence for the stable derived category. In [28], Orlov observed that the bounded stable derived category of a noetherian scheme depends only on the singular points. We extend this result to the unbounded stable derived category, using a completely different proof.

Corollary 6.10. *Let \mathbb{Y} be a seperated noetherian scheme of finite Krull dimension. If $f: \mathbb{X} \rightarrow \mathbb{Y}$ denotes the inclusion of an open subscheme which contains all singular points of \mathbb{Y} , then $\mathbf{S}f^*: \mathbf{S}(\mathrm{Qcoh} \mathbb{Y}) \rightarrow \mathbf{S}(\mathrm{Qcoh} \mathbb{X})$ is an equivalence.*

Proof. We apply Proposition 6.9 and need to show that $\mathbf{S}_{\mathbb{Y} \setminus \mathbb{X}}(\mathrm{Qcoh} \mathbb{Y}) = 0$. But this is clear from our assumptions on \mathbb{X} and \mathbb{Y} . \square

Orlov's result [28, Proposition 1.14] is an immediate consequence if one restricts the equivalence $\mathbf{S}f^*$ to compact objects; see Theorem 6.6.

Corollary 6.11. *Let \mathbb{Y} be a separated noetherian scheme of finite Krull dimension. If $f: \mathbb{X} \rightarrow \mathbb{Y}$ denotes the inclusion of an open subscheme which contains all singular points of \mathbb{Y} , then f^* induces (up to direct factors) an equivalence*

$$\mathbf{D}^b(\mathrm{coh} \mathbb{Y})/\mathbf{D}^{\mathrm{perf}}(\mathrm{coh} \mathbb{Y}) \longrightarrow \mathbf{D}^b(\mathrm{coh} \mathbb{X})/\mathbf{D}^{\mathrm{perf}}(\mathrm{coh} \mathbb{X}).$$

7. GORENSTEIN INJECTIVE APPROXIMATIONS AND TATE COHOMOLOGY

Let \mathcal{A} be a locally noetherian Grothendieck category and suppose that the derived category $\mathbf{D}(\mathcal{A})$ is compactly generated. In this section we study the category of complete injective resolutions. We assign functorially to each complex of injectives a complete resolution. This yields Gorenstein injective approximations and Tate cohomology groups for objects in \mathcal{A} . The classical definition of Tate cohomology is based on complete projective resolutions. Our approach is essentially the same, using however resolutions with injective instead of projective components. Another aspect in this section is the interplay between the stable derived category $\mathbf{S}(\mathcal{A})$ and the *stable category* $\underline{\mathcal{A}}$ modulo injective objects, which is obtained from \mathcal{A} by identifying two maps if their difference factors through some injective object. Given objects A, B in \mathcal{A} , we write

$$\underline{\mathrm{Hom}}_{\mathcal{A}}(A, B) = \mathrm{Hom}_{\underline{\mathcal{A}}}(A, B).$$

The functor

$$\mathbf{K}(\mathrm{Inj} \mathcal{A}) \longrightarrow \underline{\mathcal{A}}, \quad X \mapsto Z^0 X = \mathrm{Ker}(X^0 \rightarrow X^1)$$

provides a link between the stable categories $\mathbf{S}(\mathcal{A})$ and $\underline{\mathcal{A}}$. In particular, we obtain an explicit description of the stabilization functor

$$S: \mathcal{A} \xrightarrow{\mathrm{can}} \mathbf{D}(\mathcal{A}) \xrightarrow{I_\lambda \circ Q_\rho} \mathbf{S}(\mathcal{A})$$

provided that \mathcal{A} has some appropriate Gorenstein property.

Most of the concepts in this section are classical, but seem to be new in this setting and this generality. We refer to the end of this section for historical remarks and references to the literature.

Let us start with the relevant definitions. A complex X in $\mathrm{Inj} \mathcal{A}$ is called *totally acyclic* if $\mathrm{Hom}_{\mathcal{A}}(A, X)$ and $\mathrm{Hom}_{\mathcal{A}}(X, A)$ are acyclic complexes of abelian groups for all A in $\mathrm{Inj} \mathcal{A}$. We denote by $\mathbf{K}_{\mathrm{tac}}(\mathcal{A})$ the full subcategory of all totally acyclic complexes in $\mathbf{K}(\mathrm{Inj} \mathcal{A})$. Following [13], we call an object A in \mathcal{A} *Gorenstein injective* if it is of the form $Z^0 X$ for some X in $\mathbf{K}_{\mathrm{tac}}(\mathcal{A})$. We write $\mathrm{GInj} \mathcal{A}$ for the full subcategory formed by all Gorenstein injective objects.

Lemma 7.1. *Let \mathcal{A} be an abelian category and let X, Y be objects in $\mathbf{K}(\mathrm{Inj} \mathcal{A})$. Suppose $H^n X = 0$ for all $n > 0$ and Y is totally acyclic. Then the canonical map*

$$\sigma_{X,Y}: \mathrm{Hom}_{\mathbf{K}(\mathrm{Inj} \mathcal{A})}(X, Y) \longrightarrow \underline{\mathrm{Hom}}_{\mathcal{A}}(Z^0 X, Z^0 Y)$$

is bijective.

Proof. Fix a map $\alpha: Z^0 X \rightarrow Z^0 Y$ in \mathcal{A} . We need to extend α to a chain map $\bar{\alpha}: X \rightarrow Y$ such that $Z^0 \bar{\alpha} = \alpha$. We use the assumption on X to extend α in non-negative degrees, and the assumption on Y allows to extend α in negative degrees. Thus $\sigma_{X,Y}$ is surjective. To show that $\sigma_{X,Y}$ is injective, let $\phi: X \rightarrow Y$ be a chain map such that $Z^0 \phi$ factors through some injective object. A similar argument as before yields a chain homotopy $X \rightarrow Y$ which shows that ϕ is null-homotopic. Thus the proof is complete. \square

Let us denote by $\text{GInj } \underline{\mathcal{A}}$ the full subcategory of $\underline{\mathcal{A}}$ formed by the objects in $\text{GInj } \mathcal{A}$. Observe that $\text{GInj } \mathcal{A}$ is a Frobenius category with respect to the class of exact sequences from \mathcal{A} . With respect to this exact structure, an object A in $\text{GInj } \mathcal{A}$ is projective iff A is injective iff A belongs to $\text{Inj } \mathcal{A}$. Thus the category $\text{GInj } \underline{\mathcal{A}}$ carries a triangulated structure. The shift takes an object A to the cokernel ΣA of a monomorphism $A \rightarrow E$ into an injective object E . The exact triangles are induced from short exact sequences in \mathcal{A} .

Proposition 7.2. *Let \mathcal{A} be an abelian category. Then the functor*

$$\mathbf{K}_{\text{tac}}(\text{Inj } \mathcal{A}) \longrightarrow \text{GInj } \underline{\mathcal{A}}, \quad X \mapsto Z^0 X,$$

is an equivalence of triangulated categories.

Proof. We need to show that the functor is fully faithful and surjective on isomorphism classes of objects. The last property is clear from the definition of $\text{GInj } \mathcal{A}$. The functor is fully faithful by Lemma 7.1. Finally, observe that an exact triangle of complexes comes, up to isomorphism, from a sequence of complexes which is split exact in each degree. Thus we obtain an exact sequence in \mathcal{A} and an exact triangle in $\underline{\mathcal{A}}$ if we apply Z^0 . \square

The following lemma is crucial because it provides the existence of complete injective resolutions.

Lemma 7.3. *Let \mathcal{A} be a locally noetherian Grothendieck category and suppose that $\mathbf{D}(\mathcal{A})$ is compactly generated. Then the inclusion $J: \mathbf{K}_{\text{tac}}(\text{Inj } \mathcal{A}) \rightarrow \mathbf{K}(\text{Inj } \mathcal{A})$ has a left adjoint*

$$J_\lambda: \mathbf{K}(\text{Inj } \mathcal{A}) \longrightarrow \mathbf{K}_{\text{tac}}(\text{Inj } \mathcal{A}).$$

Proof. The inclusion $I: \mathbf{K}_{\text{ac}}(\text{Inj } \mathcal{A}) \rightarrow \mathbf{K}(\text{Inj } \mathcal{A})$ has a left adjoint I_λ by Theorem 4.2. Thus it is sufficient to show that the inclusion $\mathbf{K}_{\text{tac}}(\text{Inj } \mathcal{A}) \rightarrow \mathbf{K}_{\text{ac}}(\text{Inj } \mathcal{A})$ has a left adjoint. Let us fix an injective cogenerator E in \mathcal{A} . By definition, $\mathbf{K}_{\text{tac}}(\text{Inj } \mathcal{A})$ consists of all objects X in $\mathbf{K}_{\text{ac}}(\text{Inj } \mathcal{A})$ such that

$$\text{Hom}_{\mathbf{K}(\text{Inj } \mathcal{A})}(\Sigma^n E, X) \cong \text{Hom}_{\mathbf{K}_{\text{ac}}(\text{Inj } \mathcal{A})}(I_\lambda(\Sigma^n E), X)$$

vanishes for all $n \in \mathbb{Z}$. The category $\mathbf{K}_{\text{ac}}(\text{Inj } \mathcal{A})$ is compactly generated by Corollary 5.4, and we can apply Proposition 3.4 to obtain a left adjoint for the inclusion $\mathbf{K}_{\text{tac}}(\text{Inj } \mathcal{A}) \rightarrow \mathbf{K}_{\text{ac}}(\text{Inj } \mathcal{A})$. \square

Given an object A in \mathcal{A} with injective resolution iA , we call the natural map

$$iA \longrightarrow J_\lambda iA$$

a *complete injective resolution* of A . If we apply the functor Z^0 to this map, we obtain a Gorenstein injective approximation of A .

Theorem 7.4. *Let \mathcal{A} be a locally noetherian Grothendieck category and suppose that $\mathbf{D}(\mathcal{A})$ is compactly generated. Then the inclusion $\text{GInj } \underline{\mathcal{A}} \rightarrow \underline{\mathcal{A}}$ has a left adjoint*

$$T: \underline{\mathcal{A}} \longrightarrow \text{GInj } \underline{\mathcal{A}}.$$

Thus we have for each object A in \mathcal{A} a natural map $A \rightarrow TA$ which induces a bijection

$$\underline{\text{Hom}}_{\underline{\mathcal{A}}}(TA, B) \xrightarrow{\sim} \underline{\text{Hom}}_{\underline{\mathcal{A}}}(A, B) \quad \text{for all } B \in \text{GInj } \underline{\mathcal{A}}.$$

Proof. Fix an object A in \mathcal{A} and choose an injective resolution iA . We put

$$TA = Z^0(J_\lambda iA),$$

and this induces a functor $T: \underline{\mathcal{A}} \rightarrow \text{GInj } \underline{\mathcal{A}}$. Let B in $\text{GInj } \mathcal{A}$ and fix a totally acyclic complex tB such that $Z^0 tB = B$. The natural map $iA \rightarrow J_\lambda iA$ induces a map $A \rightarrow TA$ in $\underline{\mathcal{A}}$ which makes the following square commutative.

$$\begin{array}{ccc} \text{Hom}_{\mathbf{K}(\text{Inj } \mathcal{A})}(J_\lambda iA, tB) & \xrightarrow{\sim} & \text{Hom}_{\mathbf{K}(\text{Inj } \mathcal{A})}(iA, tB) \\ \downarrow Z^0 & & \downarrow Z^0 \\ \underline{\text{Hom}}_{\mathcal{A}}(TA, B) & \longrightarrow & \underline{\text{Hom}}_{\mathcal{A}}(A, B) \end{array}$$

The vertical maps are bijective by Lemma 7.1, and we conclude that T is a left adjoint for the inclusion $\text{GInj } \underline{\mathcal{A}} \rightarrow \underline{\mathcal{A}}$. \square

Next we use complete injective resolutions to define Tate cohomology groups for objects in \mathcal{A} .

Definition 7.5. Given objects A, B in \mathcal{A} and $n \in \mathbb{Z}$, the *Tate cohomology group* is

$$\widehat{\text{Ext}}_{\mathcal{A}}^n(A, B) = H^n \text{Hom}_{\mathcal{A}}(A, J_\lambda iB)$$

Remark 7.6. The correct term for this cohomology theory would be ‘injective Tate cohomology’ in order to distinguish it from the usual ‘projective Tate cohomology’ which is defined via complete projective resolutions. For simplicity, we drop the extra adjective ‘injective’. Note that confusion is not possible because we do not consider projective Tate cohomology in this paper.

Tate cohomology is natural in both arguments because the formation of complete injective resolutions is functorial. In addition, we have a *comparison map*

$$\text{Ext}_{\mathcal{A}}^n(A, B) \longrightarrow \widehat{\text{Ext}}_{\mathcal{A}}^n(A, B),$$

which is induced by the map $iB \rightarrow J_\lambda iB$. There is an alternative description of Tate cohomology which is based on the left adjoint $T: \underline{\mathcal{A}} \rightarrow \text{GInj } \underline{\mathcal{A}}$.

Proposition 7.7. *Given objects A, B in \mathcal{A} and $n \in \mathbb{Z}$, there is a natural isomorphism*

$$\widehat{\text{Ext}}_{\mathcal{A}}^n(A, B) \cong \underline{\text{Hom}}_{\mathcal{A}}(A, \Sigma^n(TB)).$$

Proof. Using Lemmas 2.1 and 7.1, we have the following sequence of isomorphisms

$$\begin{aligned} H^n \text{Hom}_{\mathcal{A}}(A, J_\lambda iB) &\cong \text{Hom}_{\mathbf{K}(\mathcal{A})}(A, \Sigma^n(J_\lambda iB)) \\ &\cong \text{Hom}_{\mathbf{K}(\mathcal{A})}(iA, \Sigma^n(J_\lambda iB)) \\ &\cong \underline{\text{Hom}}_{\mathcal{A}}(A, \Sigma^n(TB)). \end{aligned}$$

\square

We have a more conceptual definition of Tate cohomology for objects in $\mathbf{D}(\mathcal{A})$ which uses the composite

$$U: \mathbf{D}(\mathcal{A}) \xrightarrow{J_\lambda \circ Q_\rho} \mathbf{K}_{\text{tac}}(\text{Inj } \mathcal{A}).$$

Thus we define for objects X, Y in $\mathbf{D}(\mathcal{A})$ and $n \in \mathbb{Z}$

$$\widehat{\text{Ext}}_{\mathcal{A}}^n(X, Y) = \text{Hom}_{\mathbf{K}(\mathcal{A})}(UX, \Sigma^n(UY)).$$

Note that this definition is consistent with the original definition of Tate cohomology if we take objects in \mathcal{A} and view them as complexes concentrated in degree 0. This follows from the fact that $Q_\rho X$ is nothing but an injective resolution of X^0 when X is concentrated in degree 0. From now on, we will use one of the alternative descriptions of Tate cohomology whenever this is convenient.

Next we show that each exact sequence in \mathcal{A} induces a long exact sequence in Tate cohomology. This is based on the following simple lemma.

Lemma 7.8. *The left adjoint $T: \underline{\mathcal{A}} \rightarrow \text{GInj } \underline{\mathcal{A}}$ has the following properties.*

- (1) *An exact sequence $0 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0$ in \mathcal{A} induces an exact triangle*

$$TA' \longrightarrow TA \longrightarrow TA'' \longrightarrow \Sigma(TA') \quad \text{in } \text{GInj } \underline{\mathcal{A}}.$$

- (2) *Let A, B be in \mathcal{A} and $n \in \mathbb{Z}$. The natural map $A \rightarrow TA$ induces an isomorphism*

$$\widehat{\text{Ext}}_{\mathcal{A}}^n(TA, B) \cong \widehat{\text{Ext}}_{\mathcal{A}}^n(A, B).$$

Proof. (1) We have an exact triangle $A' \rightarrow A \rightarrow A'' \rightarrow \Sigma(A')$ in $\mathbf{D}(\mathcal{A})$. Now use that the exact functor $Z^0 \circ J_\lambda \circ Q_\rho$ computes T .

- (2) The adjointness property of T implies $\underline{\text{Hom}}_{\mathcal{A}}(TA, TB) \cong \underline{\text{Hom}}_{\mathcal{A}}(A, TB)$. \square

Proposition 7.9. *Let $0 \rightarrow B' \rightarrow B \rightarrow B'' \rightarrow 0$ be an exact sequence in \mathcal{A} . Then we have for A and C in \mathcal{A} the following long exact sequences.*

$$\begin{aligned} \cdots \longrightarrow \widehat{\text{Ext}}_{\mathcal{A}}^n(A, B') &\longrightarrow \widehat{\text{Ext}}_{\mathcal{A}}^n(A, B) \longrightarrow \widehat{\text{Ext}}_{\mathcal{A}}^n(A, B'') \longrightarrow \\ &\longrightarrow \widehat{\text{Ext}}_{\mathcal{A}}^{n+1}(A, B') \longrightarrow \widehat{\text{Ext}}_{\mathcal{A}}^{n+1}(A, B) \longrightarrow \widehat{\text{Ext}}_{\mathcal{A}}^{n+1}(A, B'') \longrightarrow \cdots \\ \cdots \longrightarrow \widehat{\text{Ext}}_{\mathcal{A}}^n(B'', C) &\longrightarrow \widehat{\text{Ext}}_{\mathcal{A}}^n(B, C) \longrightarrow \widehat{\text{Ext}}_{\mathcal{A}}^n(B', C) \longrightarrow \\ &\longrightarrow \widehat{\text{Ext}}_{\mathcal{A}}^{n+1}(B'', C) \longrightarrow \widehat{\text{Ext}}_{\mathcal{A}}^{n+1}(B, C) \longrightarrow \widehat{\text{Ext}}_{\mathcal{A}}^{n+1}(B', C) \longrightarrow \cdots \end{aligned}$$

Proof. We apply Lemma 7.8 and use the fact that $\underline{\text{Hom}}_{\mathcal{A}}(TA, -)$ and $\underline{\text{Hom}}_{\mathcal{A}}(-, TC)$ are cohomological functors. \square

We compute Tate cohomology for Gorenstein injective objects.

Proposition 7.10. *Let A, B be objects in \mathcal{A} and suppose B is Gorenstein injective. Then the comparison map*

$$\text{Ext}_{\mathcal{A}}^n(A, B) \longrightarrow \widehat{\text{Ext}}_{\mathcal{A}}^n(A, B)$$

is an isomorphism for $n > 0$ and induces an isomorphism

$$\underline{\text{Hom}}_{\mathcal{A}}(A, B) \xrightarrow{\sim} \widehat{\text{Ext}}_{\mathcal{A}}^0(A, B) \quad \text{for } n = 0.$$

Proof. Our assumption implies $TB = B$. The case $n = 0$ is clear. For $n = 1$, choose an exact sequence $0 \rightarrow B \rightarrow E \rightarrow \Sigma B \rightarrow 0$ with E injective and apply $\underline{\text{Hom}}_{\mathcal{A}}(A, -)$. The cokernel of

$$\text{Hom}_{\mathcal{A}}(A, E) \longrightarrow \text{Hom}_{\mathcal{A}}(A, \Sigma B)$$

is isomorphic $\text{Ext}_{\mathcal{A}}^1(A, B)$; it is isomorphic to $\underline{\text{Hom}}_{\mathcal{A}}(A, \Sigma B)$ since B is Gorenstein injective. For $n > 1$, use dimension shift. \square

Next we describe those objects A in \mathcal{A} such that $\widehat{\text{Ext}}_{\mathcal{A}}^*(A, -)$ vanishes. For instance, Tate cohomology vanishes for all objects having finite projective or finite injective dimension.

Proposition 7.11. *For an object A in \mathcal{A} , the following are equivalent.*

- (1) $\widehat{\text{Ext}}_{\mathcal{A}}^*(A, -) = 0$.
- (2) $\widehat{\text{Ext}}_{\mathcal{A}}^0(A, A) = 0$.
- (3) $\widehat{\text{Ext}}_{\mathcal{A}}^*(-, A) = 0$.
- (4) $\text{Ext}_{\mathcal{A}}^1(A, B) = 0$ for all $B \in \text{GInj } \mathcal{A}$.
- (5) $\underline{\text{Hom}}_{\mathcal{A}}(A, B) = 0$ for all $B \in \text{GInj } \mathcal{A}$.

Proof. Use the isomorphism $\widehat{\text{Ext}}_{\mathcal{A}}^*(-, -) \cong \underline{\text{Hom}}_{\mathcal{A}}^*(T-, T-)$ and the fact that $\widehat{\text{Ext}}_{\mathcal{A}}^1(-, B)$ can be computed via $\text{Ext}_{\mathcal{A}}^1(-, TB)$. \square

The following result formulates our analogue of the maximal Cohen-Macaulay approximation in the sense of Auslander and Buchweitz [2]. Note that a Gorenstein injective object is the ‘dual’ of a maximal Cohen-Macaulay object which one defines in a category with enough projectives. Let

$$\mathcal{X} = \{A \in \mathcal{A} \mid \text{Ext}_{\mathcal{A}}^1(A, B) = 0 \text{ for all } B \in \text{GInj } \mathcal{A}\} \quad \text{and} \quad \mathcal{Y} = \text{GInj } \mathcal{A}.$$

Theorem 7.12. *Let \mathcal{A} be a locally noetherian Grothendieck category and suppose that $\mathbf{D}(\mathcal{A})$ is compactly generated.*

- (1) *Every object A in \mathcal{A} fits into exact sequences*

$$0 \rightarrow Y_A \rightarrow X_A \rightarrow A \rightarrow 0 \quad \text{and} \quad 0 \rightarrow A \rightarrow Y^A \rightarrow X^A \rightarrow 0$$

in \mathcal{A} with X_A, X^A in \mathcal{X} and Y_A, Y^A in \mathcal{Y} .

- (2) *The map $A \mapsto X_A$ induces a right adjoint for the inclusion $\underline{\mathcal{X}} \rightarrow \underline{\mathcal{A}}$.*
- (3) *The map $A \mapsto Y^A$ induces a left adjoint for the inclusion $\underline{\mathcal{Y}} \rightarrow \underline{\mathcal{A}}$.*
- (4) $\mathcal{X} \cap \mathcal{Y} = \text{Inj } \mathcal{A}$.

Note that this is essentially the statement of Theorem 1.3 from the introduction, since \mathcal{X} is precisely the subcategory of objects A in \mathcal{A} such that the Tate cohomology functor $\widehat{\text{Ext}}_{\mathcal{A}}^*(A, -)$ vanishes.

Proof. We use the basic properties of Tate cohomology.

- (1) Fix an object A in \mathcal{A} and a complete injective resolution

$$iA \longrightarrow J_{\lambda}iA = yA.$$

We complete this map to an exact triangle

$$(7.1) \quad iA \longrightarrow yA \longrightarrow xA \longrightarrow \Sigma(iA).$$

in $\mathbf{K}(\text{Inj } \mathcal{A})$ and have therefore a sequence $0 \rightarrow iA \rightarrow yA \rightarrow xA \rightarrow 0$ of complexes which is split exact in each degree. Applying $Z^0: \mathbf{K}(\text{Inj } \mathcal{A}) \rightarrow \underline{\mathcal{A}}$ produces an exact sequence

$$0 \longrightarrow A \xrightarrow{\alpha} Y^A \xrightarrow{\beta} X^A \longrightarrow 0$$

in \mathcal{A} . Clearly, Y^A belongs to \mathcal{Y} . On the other hand, $\widehat{\text{Ext}}_{\mathcal{A}}^*(\alpha, -)$ is an isomorphism. Thus $\widehat{\text{Ext}}_{\mathcal{A}}^*(X^A, -)$ vanishes and X^A belongs to \mathcal{X} . The second sequence ending in A is obtained by rotating the triangle (7.1).

(2) We consider the exact sequence

$$0 \longrightarrow Y_A \xrightarrow{\mu} X_A \xrightarrow{\nu} A \longrightarrow 0$$

and need to show that $\underline{\text{Hom}}_{\mathcal{A}}(X, \nu)$ is bijective for all X in \mathcal{X} . To see this, let $\phi: X \rightarrow A$ be a map with X in \mathcal{X} . The map ϕ factors through ν since $\text{Ext}_{\mathcal{A}}^1(X, Y_A) = 0$. Therefore $\underline{\text{Hom}}_{\mathcal{A}}(X, \nu)$ is surjective. To show that $\underline{\text{Hom}}_{\mathcal{A}}(X, \nu)$ is injective, let $\psi: X \rightarrow X_A$ be a map such that $\nu \circ \psi$ has a factorization $X \xrightarrow{\phi'} E \xrightarrow{\phi''} A$ with injective E . We obtain a factorization $\phi'' = \nu \circ \chi$, since E belongs to \mathcal{X} . We have $\nu \circ (\psi - \chi \circ \phi') = 0$, and $\psi - \chi \circ \phi'$ needs to factor through μ . Therefore $\psi - \chi \circ \phi'$ factors through some injective object, since $\underline{\text{Hom}}_{\mathcal{A}}(X, Y_A) = 0$. We conclude that ψ factors through an injective object. Thus the map $\underline{\text{Hom}}_{\mathcal{A}}(X, \nu)$ is bijective.

(3) We consider the exact sequence

$$0 \longrightarrow A \xrightarrow{\alpha} Y^A \xrightarrow{\beta} X^A \longrightarrow 0$$

and need to show that $\underline{\text{Hom}}_{\mathcal{A}}(\alpha, Y)$ is bijective for all Y in \mathcal{Y} . But this is clear from the long exact sequence for Tate cohomology, since $\underline{\text{Hom}}_{\mathcal{A}}(-, Y) \cong \widehat{\text{Ext}}_{\mathcal{A}}^0(-, Y)$ and $\widehat{\text{Ext}}_{\mathcal{A}}^*(X^A, -)$ vanishes.

(4) Clearly, $\text{Inj } \mathcal{A}$ is contained in $\mathcal{X} \cap \mathcal{Y}$. Now let A be in $\mathcal{X} \cap \mathcal{Y}$. Thus

$$\underline{\text{Hom}}_{\mathcal{A}}(A, A) \cong \widehat{\text{Ext}}_{\mathcal{A}}^0(A, A) = 0.$$

Therefore the identity map $A \rightarrow A$ factors through an injective object. We conclude that A is injective. \square

Let us comment on the interplay between the stable category $\underline{\mathcal{A}}$ and the stable derived category $\mathbf{S}(\mathcal{A})$. We have already seen that the definition of Tate cohomology is possible in both settings. It is more elementary in $\underline{\mathcal{A}}$, but more conceptual using the category of complete injective resolutions $\mathbf{K}_{\text{tac}}(\text{Inj } \mathcal{A})$ which is a subcategory of $\mathbf{S}(\mathcal{A})$. The same phenomenon appears when one studies Gorenstein injective approximations. The proof of Theorem 7.12 we have given uses the category of complexes $\mathbf{K}(\text{Inj } \mathcal{A})$. There is an alternative proof which avoids complexes and uses instead the left adjoint $T: \underline{\mathcal{A}} \rightarrow \text{GInj } \underline{\mathcal{A}}$.

Gorenstein rings and schemes play an important role in applications and have a number of interesting homological properties. It is therefore important to formulate a Gorenstein property for a locally noetherian Grothendieck category \mathcal{A} . Let us denote by $\Sigma^\infty \mathcal{A}$ the full subcategory of objects A in \mathcal{A} which fit into an exact sequence

$$\cdots \longrightarrow E_2 \longrightarrow E_1 \longrightarrow E_0 \longrightarrow A \longrightarrow 0$$

with E_n injective for all n . We say that \mathcal{A} has the *injective Gorenstein property* if the equivalent conditions in the following proposition are satisfied. This property has been studied by Beligiannis in [4].

Proposition 7.13. *Let \mathcal{A} be a locally noetherian Grothendieck category and suppose that $\mathbf{D}(\mathcal{A})$ is compactly generated. Then the following are equivalent.*

- (1) $\text{Ext}_{\mathcal{A}}^1(A, B) = 0$ for all $A \in \text{Inj } \mathcal{A}$ and $B \in \Sigma^\infty \mathcal{A}$.
- (2) Every acyclic complex in $\text{Inj } \mathcal{A}$ is totally acyclic.
- (3) $\text{GInj } \mathcal{A} = \Sigma^\infty \mathcal{A}$.
- (4) $S: \mathcal{A} \xrightarrow{\text{can}} \mathbf{D}(\mathcal{A}) \xrightarrow{I_\lambda \circ Q_\rho} \mathbf{S}(\mathcal{A})$ annihilates all injective objects.

(5) S induces an equivalence $\mathrm{GInj} \underline{\mathcal{A}} \rightarrow \mathbf{S}(\mathcal{A})$.

Proof. The conditions (1) - (3) are pairwise equivalent. This follows from the formula

$$\mathrm{Ext}_{\mathcal{A}}^n(A, Z^0 Y) \cong \mathrm{Hom}_{\mathbf{K}(\mathcal{A})}(A, \Sigma^n Y) \cong H^n \mathrm{Hom}_{\mathcal{A}}(A, Y)$$

where A is any object in \mathcal{A} and Y is an acyclic complex in $\mathrm{Inj} \mathcal{A}$. The first isomorphism is valid for all $n \geq 1$, and the second for all $n \in \mathbb{Z}$.

Now observe that I_λ annihilates precisely those objects X in $\mathbf{K}(\mathrm{Inj} \mathcal{A})$ such that $\mathrm{Hom}_{\mathbf{K}(\mathrm{Inj} \mathcal{A})}(X, Y) = 0$ for every acyclic complex Y in $\mathrm{Inj} \mathcal{A}$. On the other hand, $\mathcal{A} \rightarrow \mathbf{D}(\mathcal{A})$ and Q_ρ are faithful. Thus (1) - (3) are equivalent to (4). Also, (5) implies (4). So, it remains to show that (1) - (4) imply (5).

Suppose (2) and (4) hold. We have already seen in Proposition 7.2 that

$$\mathbf{S}(\mathcal{A}) \longrightarrow \mathrm{GInj} \underline{\mathcal{A}}, \quad X \mapsto Z^0 X,$$

is an equivalence, since every acyclic complex is totally acyclic. On the other hand, S annihilates all injective objects and induces therefore a functor $\mathrm{GInj} \underline{\mathcal{A}} \rightarrow \mathbf{S}(\mathcal{A})$. It is easily checked that both functors are quasi-inverse to each other. \square

We are now in the position that we can describe the stabilization functor $S: \mathcal{A} \rightarrow \mathbf{S}(\mathcal{A})$, provided that \mathcal{A} has the injective Gorenstein property. We use the left adjoint $T: \underline{\mathcal{A}} \rightarrow \mathrm{GInj} \underline{\mathcal{A}}$ of the inclusion $\mathrm{GInj} \underline{\mathcal{A}} \rightarrow \underline{\mathcal{A}}$. For A in \mathcal{A} , choose any acyclic complex X of injective objects such that $Z^0 X \cong TA$. Then $SA \cong X$.

Example 7.14. Let Λ be a ring and suppose Λ is *Gorenstein*, that is, Λ is two-sided noetherian and Λ has finite injective dimension as left and right Λ -module. In this case, the category $\mathrm{Mod} \Lambda$ has the injective Gorenstein property. This follows from the fact that every injective Λ -module has finite projective dimension if Λ is Gorenstein; see Example 5.6. Given a Λ -module A , Tate cohomology $\widehat{\mathrm{Ext}}_\Lambda^*(A, -)$ vanishes iff A has finite injective dimension iff A has finite projective dimension. Note that for Gorenstein rings, the classical Tate cohomology defined via complete projective resolutions coincides with our Tate cohomology, which is defined via complete injective resolutions; see [5].

Example 7.15. Let Λ be a ring and suppose that projective and injective Λ -modules coincide. Then every Λ -module is Gorenstein injective. In particular, $\mathbf{S}(\mathrm{Mod} \Lambda)$ is equivalent to the stable category $\underline{\mathrm{Mod}} \Lambda$. Given a Λ -module A , there is an exact triangle

$$pA \longrightarrow iA \longrightarrow tA \longrightarrow \Sigma(pA)$$

in $\mathbf{K}(\mathrm{Inj} \mathcal{A})$ where pA denotes a projective, iA an injective, and tA a Tate resolution of A . This triangle is isomorphic to the canonical triangle

$$(Q_\lambda \circ Q) \bar{A} \longrightarrow \bar{A} \longrightarrow (I \circ I_\lambda) \bar{A} \longrightarrow \Sigma(Q_\lambda \circ Q) \bar{A}$$

where $\bar{A} = Q_\rho A$. For details in case Λ is a finite dimensional cocommutative Hopf algebra, see [18].

Example 7.16. Let \mathbb{X} be a noetherian scheme and suppose that every injective object E in $\mathrm{Qcoh} \mathbb{X}$ admits a finite resolution

$$0 \longrightarrow L_r \longrightarrow \cdots \longrightarrow L_2 \longrightarrow L_1 \longrightarrow L_0 \longrightarrow E \longrightarrow 0$$

with L_n locally free for each n . Then the category $\mathrm{Qcoh} \mathbb{X}$ has the injective Gorenstein property.

Historical remarks. Gorenstein injective approximations and Tate cohomology have a long history. Auslander and Bridger [1] introduce the stable module category and assign to each module a G-dimension. Over Gorenstein rings, the modules of G-dimension 0 are precisely the maximal Cohen-Macaulay or Gorenstein projective modules. Auslander and Buchweitz establish maximal Cohen-Macaulay approximations in [2], and there is an alternative unpublished approach by Buchweitz [12] which involves the derived category. Enochs and his collaborators drop finiteness conditions on modules and prove the existence of Gorenstein projective and Gorenstein injective approximations for arbitrary modules, for instance over Gorenstein rings [14]. Further generalizations can be found in work of Beligiannis [4]. Jørgensen [19] constructs Gorenstein projective approximations for artin algebras via Bousfield localization, using the category of complete projective resolutions. Papers of Hovey [17] and Beligiannis and Reiten [5] employ the formalism of model category structures and cotorsion pairs.

The exposition of Buchweitz [12] discusses the close connection between maximal Cohen-Macaulay approximations and Tate cohomology over Gorenstein rings. For more general settings, we refer to the work of Beligiannis and Reiten [5]. Another exposition of Tate cohomology over noetherian rings can be found in a paper of Avramov and Martsinkovsky [3].

APPENDIX A. THE DG CATEGORY OF NOETHERIAN OBJECTS

Let \mathcal{A} be a locally noetherian Grothendieck category. We give an alternative description of the homotopy category $\mathbf{K}(\text{Inj } \mathcal{A})$ as the derived category of some DG category. Here, we follow closely Keller's exposition in [22].

Let \mathcal{C} be a small DG category. We recall the definition of the derived category $\mathbf{D}_{\text{dg}}(\mathcal{C})$ of \mathcal{C} . The category $\mathbf{C}_{\text{dg}}(\mathcal{C})$ of *cochain complexes* has by definition as objects all DG \mathcal{C} -modules. A map in $\mathbf{C}_{\text{dg}}(\mathcal{C})$ is a map of DG \mathcal{C} -modules which is homogeneous of degree zero and commutes with the differential. The *homotopy category* $\mathbf{K}_{\text{dg}}(\mathcal{C})$ is obtained from $\mathbf{C}_{\text{dg}}(\mathcal{C})$ by identifying homotopy equivalent maps, where $f, g: X \rightarrow Y$ are homotopy equivalent if there exists a map $s: X \rightarrow Y$ of graded modules which is homogeneous of degree -1 and satisfies

$$(f - g)^n = s^{n+1} \circ d + d \circ s^n \quad \text{for all } n \in \mathbb{Z}.$$

Finally, the *derived category* of \mathcal{C} is obtained from $\mathbf{K}_{\text{dg}}(\mathcal{C})$ as the localization

$$\mathbf{D}_{\text{dg}}(\mathcal{C}) = \mathbf{K}_{\text{dg}}(\mathcal{C})[Q^{-1}]$$

with respect to the class Q of all maps f which induce an isomorphism H^*f .

Given two cochain complexes X and Y in \mathcal{A} , we define the cochain complex $\mathcal{H}om_{\mathcal{A}}(X, Y)$. The n th component is

$$\prod_{p \in \mathbb{Z}} \text{Hom}_{\mathcal{A}}(X^p, Y^{n+p})$$

and the differential is given by

$$d(f^p) = d \circ f^p - (-1)^n f^{p+1} \circ d.$$

Now fix a class \mathcal{C} of objects in \mathcal{A} . We obtain a DG category $\bar{\mathcal{C}}$ by taking as objects for each A in \mathcal{C} an injective resolution \bar{A} , and as maps

$$\text{Hom}_{\bar{\mathcal{C}}}(\bar{A}, \bar{B}) = \mathcal{H}om_{\mathcal{A}}(\bar{A}, \bar{B}).$$

Proposition A.1. *Let \mathcal{A} be a locally noetherian Grothendieck category, and let \mathcal{C} be a class of noetherian objects which generate $\mathbf{D}^b(\text{noeth } \mathcal{A})$, that is, there is no proper thick subcategory containing \mathcal{C} . Then the functor*

$$\mathbf{K}(\text{Inj } \mathcal{A}) \longrightarrow \mathbf{D}_{\text{dg}}(\bar{\mathcal{C}}), \quad X \mapsto \mathcal{H}om_{\mathcal{A}}(-, X)|_{\bar{\mathcal{C}}},$$

is an equivalence of triangulated categories.

Proof. The functor is exact. To see that it preserves coproducts, fix an object A in \mathcal{C} and a family of objects X_i in $\mathbf{K}(\text{Inj } \mathcal{A})$. Then we have for every $n \in \mathbb{Z}$

$$\begin{aligned} H^n \coprod_i \mathcal{H}om_{\mathcal{A}}(\bar{A}, X_i) &\cong \coprod_i H^n \mathcal{H}om_{\mathcal{A}}(\bar{A}, X_i) \cong \coprod_i \text{Hom}_{\mathbf{K}(\text{Inj } \mathcal{A})}(\Sigma^{-n} \bar{A}, X_i) \\ &\cong \text{Hom}_{\mathbf{K}(\text{Inj } \mathcal{A})}(\Sigma^{-n} \bar{A}, \coprod_i X_i) \cong H^n \mathcal{H}om_{\mathcal{A}}(\bar{A}, \coprod_i X_i) \end{aligned}$$

since \bar{A} is compact in $\mathbf{K}(\text{Inj } \mathcal{A})$ by Lemma 2.1. Thus the canonical map

$$\coprod_i \mathcal{H}om_{\mathcal{A}}(-, X_i)|_{\bar{\mathcal{C}}} \longrightarrow \mathcal{H}om_{\mathcal{A}}(-, \coprod_i X_i)|_{\bar{\mathcal{C}}}$$

is an isomorphism. Furthermore, the functor induces for objects A and B in \mathcal{C} bijections

$$\text{Hom}_{\mathbf{K}(\text{Inj } \mathcal{A})}(\bar{A}, \Sigma^n \bar{B}) \cong H^n \mathcal{H}om_{\mathcal{A}}(\bar{A}, \bar{B}) \cong H^n \text{Hom}_{\bar{\mathcal{C}}}(\bar{A}, \bar{B}) \cong \text{Hom}_{\mathbf{D}_{\text{dg}}(\bar{\mathcal{C}})}(\bar{A}^\wedge, \Sigma^n \bar{B}^\wedge),$$

where \bar{A}^\wedge denotes the free module $\text{Hom}_{\bar{\mathcal{C}}}(-, \bar{A})$. Using infinite dévissage, we conclude that the functor is fully faithful since \mathcal{C} generates $\mathbf{K}(\text{Inj } \mathcal{A})$. The functor is, up to isomorphism, surjective on objects since the image contains the free $\bar{\mathcal{C}}$ -modules which generate $\mathbf{D}_{\text{dg}}(\bar{\mathcal{C}})$. \square

Corollary A.2. *Viewing $\text{noeth } \mathcal{A}$ as DG category, we have an equivalence*

$$\mathbf{K}(\text{Inj } \mathcal{A}) \xrightarrow{\sim} \mathbf{D}_{\text{dg}}(\text{noeth } \mathcal{A}).$$

We remark that the proof of Proposition A.1 works for any homotopy category. To be precise, let \mathcal{X} be an additive category with arbitrary coproducts and let \mathcal{C} be a set of objects in $\mathbf{K}(\mathcal{X})$ which are compact (when viewed as objects in the localizing subcategory generated by \mathcal{C}). Define $\bar{\mathcal{C}}$ as before by

$$\text{Hom}_{\bar{\mathcal{C}}}(A, B) = \mathcal{H}om_{\mathcal{X}}(A, B)$$

for A and B in \mathcal{C} . Then the functor

$$\mathbf{K}(\mathcal{X}) \longrightarrow \mathbf{D}_{\text{dg}}(\bar{\mathcal{C}}), \quad X \mapsto \mathcal{H}om_{\mathcal{X}}(-, X)|_{\bar{\mathcal{C}}},$$

induces an equivalence between the localizing subcategory which is generated by \mathcal{C} , and $\mathbf{D}_{\text{dg}}(\bar{\mathcal{C}})$.

Acknowledgements. It is a pleasure to thank a number of colleagues for their interest in the topic of this paper: Øyvind Solberg helped to prove the crucial Proposition 2.3. Peter Jørgensen pointed out the relevance of the stable derived category in algebraic geometry. Amnon Neeman communicated an alternative proof of Corollary 4.4 in the algebraic geometric context. Ragnar Buchweitz explained the connection with his earlier unpublished work. Finally, Apostolos Beligiannis provided numerous comments on a preliminary version of this manuscript.

REFERENCES

- [1] M. AUSLANDER AND M. BRIDGER: Stable module theory. *Mem. Amer. Math. Soc.* **94** (1969).
- [2] M. AUSLANDER AND R.-O. BUCHWEITZ: Maximal Cohen-Macaulay approximations. *Mem. Soc. Math. France* **38** (1989), 5–37.
- [3] L. L. AVRAMOV AND A. MARTSINKOVSKY: Absolute, relative, and Tate cohomology of modules of finite Gorenstein dimension. *Proc. London Math. Soc.* **85** (2002), 393–440.
- [4] A. BELIGIANNIS: The homological theory of contravariantly finite subcategories: Auslander-Buchweitz contexts, Gorenstein categories and (co-)stabilization. *Comm. Algebra* **28** (2000), 4547–4596.
- [5] A. BELIGIANNIS AND I. REITEN: Homological and Homotopical Aspects of Torsion Theories. Preprint (2002), 200 pp.
- [6] A. A. BEILINSON, J. BERNSTEIN, AND P. DELIGNE: Faisceaux pervers. *Astérisque* **100** (1982).
- [7] A. BEILINSON, V. GINZBURG, AND W. SOERGEL: Koszul duality patterns in representation theory. *J. Amer. Math. Soc.* **9** (1996), 473–527.
- [8] J. N. BERNSTEIN, I. M. GELFAND, AND S. I. GELFAND: Algebraic bundles over \mathbb{P}^n and problems of linear algebra. *Funktional. Anal. i Prilozhen.* **12** (1978), 66–67.
- [9] A. BONDAL AND M. VAN DEN BERGH: Generators and representability of functors in commutative and noncommutative geometry. *Mosc. Math. J.* **3** (2003), 1–36.
- [10] M. BÖKSTEDT AND A. NEEMAN: Homotopy limits in triangulated categories. *Compositio Math.* **86** (1993), 209–234.
- [11] R.-O. BUCHWEITZ, D. EISENBUD, AND J. HERZOG: Cohen-Macaulay modules on quadrics. In: *Singularities, representation of algebras, and vector bundles*. Springer Lecture Notes in Math. **1273** (1987), 58–116.
- [12] R.-O. BUCHWEITZ: Maximal Cohen-Macaulay modules and Tate-cohomology over Gorenstein rings. Unpublished manuscript (1987), 155 pp.
- [13] E. E. ENOCHS AND O. M. G. JENDA: Gorenstein injective and projective modules. *Math. Z.* **220** (1995), 611–633.
- [14] E. E. ENOCHS AND O. M. G. JENDA: *Relative homological algebra*. Walter de Gruyter, Berlin (2000).
- [15] P. GABRIEL: Des catégories abéliennes. *Bull. Soc. Math. France* **90** (1962), 323–448.
- [16] R. HARTSHORNE: *Residues and Duality*, Springer Lecture Notes in Math. **20** (1966).
- [17] M. HOVEY: Cotorsion pairs, model category structures, and representation theory. *Math. Z.* **241** (2002), 553–592.
- [18] M. HOVEY, J. H. PALMIERI, AND N. P. STRICKLAND: Axiomatic stable homotopy theory. *Mem. Amer. Math. Soc.* **610** (1997).
- [19] P. JØRGENSEN: Spectra of modules. *J. Algebra* **244** (2001), 744–784.
- [20] P. JØRGENSEN: The homotopy category of complexes of projective modules. Preprint (2003).
- [21] P. JØRGENSEN: A non-commutative BGG correspondence. Preprint (2003).
- [22] B. KELLER: Deriving DG categories. *Ann. Sci. Ecole. Norm. Sup.* **27** (1994), 63–102.
- [23] B. KELLER AND D. VOSSIECK: Sous les catégories dérivées. *C. R. Acad. Sci. Paris* **305** (1987), 225–228.
- [24] H. KRAUSE: A Brown representability theorem via coherent functors. *Topology* **41** (2002), 853–861.
- [25] A. NEEMAN: The Connection between the K -theory localization theorem of Thomason, Trobaugh and Yao and the smashing subcategories of Bousfield and Ravenel. *Ann. Sci. École Norm. Sup.* **25** (1992), 547–566.
- [26] A. NEEMAN: The Grothendieck duality theorem via Bousfield’s techniques and Brown representability. *J. Amer. Math. Soc.* **9** (1996), 205–236.
- [27] A. NEEMAN: *Triangulated categories*. Annals of Mathematics Studies **148**, Princeton University Press (2001).
- [28] D. ORLOV: Triangulated categories of singularities and D-branes in Landau-Ginzburg models. arXiv (2003).
- [29] R. RENTSCHLER: Sur les modules M tels que $\text{Hom}(M, -)$ commute avec les sommes directes. *C. R. Acad. Sc. Paris Ser. A* **268** (1969), 930–933.
- [30] J. RICKARD: Derived categories and stable equivalence. *J. Pure Appl. Algebra* **61** (1989), 303–317.

- [31] N. SPALTENSTEIN: Resolutions of unbounded complexes. *Compositio Math.* **65** (1988), 121–154.
- [32] J. L. VERDIER: Des catégories dérivées des catégories abéliennes. *Astérisque* **239** (1996).

HENNING KRAUSE, INSTITUT FÜR MATHEMATIK, UNIVERSITÄT PADERBORN, 33095 PADERBORN, GERMANY.

E-mail address: `hkrause@math.uni-paderborn.de`