

MODERN FOUNDATIONS FOR STABLE HOMOTOPY THEORY

A. D. ELMENDORF, I. KRIZ, AND J. P. MAY

CONTENTS

Introduction	1
1. Spectra and the stable homotopy category	6
2. Smash products and twisted half-smash products	11
3. The category of S-modules and its derived category	13
4. The smash product of S-modules	15
5. A_∞ and E_∞ ring spectra and their categories of modules	19
6. The smash product of R-modules and function R-modules	22
7. Tor and Ext in topology and algebra	26
8. Universal coefficient and Künneth spectral sequences	29
9. Algebraic constructions in the derived category of R-modules	31
10. Algebra structures on localizations and on quotients by ideals	34
11. The specialization to MU-modules and algebras	37
References	39

INTRODUCTION

It is a truism that algebraic topology is a very young subject. In some of its most fundamental branches, the foundations have not yet reached a state of shared consensus. Our theme will be stable homotopy theory and an emerging consensus on what its foundations should be. The consensus is different than would have been the case as recently as a decade ago. We shall illustrate the force of the change of paradigm with new constructions of some of the most basic objects in modern algebraic topology, namely the various spectra and cohomology theories that can be derived from complex cobordism. The two following articles will give introductions to completions

in stable homotopy theory and to equivariant stable homotopy theory. The three papers have a common theme: the relationship between commutative algebra and stable homotopy theory, both relations of analogy and relations of application.

Stable homotopy theory began around 1937 with the Freudenthal suspension theorem. In simplest terms, it states that, if q is small relative to n , then $\pi_{n+q}(S^n)$ is independent of n . Stable phenomena had of course appeared earlier, at least implicitly: reduced homology and cohomology are examples of functors that are invariant under suspension without limitation on dimension. Stable homotopy theory emerged as a distinct branch of algebraic topology with Adams' introduction of his eponymous spectral sequence and his spectacular conceptual use of the notion of stable phenomena in his solution to the Hopf invariant one problem. Its centrality was reinforced by two related developments that occurred at very nearly the same time, in the late 1950's. One was the introduction of generalized homology and cohomology theories and especially K -theory, by Atiyah and Hirzebruch. The other was the work of Thom which showed how to reduce the problem of classifying manifolds up to cobordism to a problem, more importantly, a solvable problem, in stable homotopy theory. The reduction of geometric phenomena to solvable problems in stable homotopy theory has remained an important mathematical theme, the most recent major success being Stolz's use of Spin cobordism to study the classification of manifolds with positive scalar curvature. In an entirely different direction, the early 1970's saw Quillen's introduction of higher algebraic K -theory and the recognition by Segal and others that it could be viewed as a construction in stable homotopy theory. With algebraic K -theory as an intermediary, there has been a growing volume of work that relates algebraic geometry to stable homotopy theory. With Waldhausen's introduction of the algebraic K -theory of spaces in the late 1970's, stable homotopy became a bridge between algebraic K -theory and the study of diffeomorphisms of manifolds. Within algebraic topology, the study of stable homotopy theory has been and remains the focus of much of the best work in the subject. The study of nilpotence and periodic phenomena by Hopkins, Mahowald, Ravenel, and many others has been especially successful.

We shall focus on the study of structured ring, module, and algebra spectra. This study plays a significant role in all of the directions of work that we have just mentioned and would have been technically impossible within the foundational consensus that existed a decade ago.

Stable homotopy theory demands a category in which to work. One could set up the ordinary Adams spectral sequence ad hoc, as Adams did, but it would be ugly at best to set up the Adams spectral sequence based on a generalized homology theory that way. One wants objects – called spectra – that play the role of spaces in unstable

homotopy theory, and one wants a category in which all of the usual constructions on spaces are present and, up to homotopy, the suspension functor is an equivalence. At this point, we introduce a sharp distinction: there is a category of point-set level objects, and there is an associated homotopy category. There has been consensus on what the latter should be, up to equivalence of categories, since the fundamental work of Boardman in the 1960's. The change in paradigm concerns the point-set level category that underlies the stable homotopy category.

There is an analogy with algebra that is fundamental to an understanding of this area of mathematics. Suppose given a (discrete) commutative ring R . It has an associated category \mathcal{M}_R of (\mathbb{Z} -graded) chain complexes, there is a notion of homotopy between maps of chain complexes, and there is a resulting homotopy category $h\mathcal{M}_R$. However, this is not the category that algebraists are interested in. For example, if R -modules M and N are regarded as chain complexes concentrated in degree zero, then, in the derived category, the homology of their tensor product should be their torsion product $\mathrm{Tor}_*^R(M, N)$. Formally, the fundamental invariants of chain complexes are their homology groups, and one constructs a category that reflects this. A map of chain complexes is said to be a quasi-isomorphism if it induces an isomorphism of homology groups. The derived category \mathcal{D}_R is obtained by adjoining formal inverses to the quasi-isomorphisms. The best way to make this rigorous is to introduce a notion of cell R -module such that every quasi-isomorphism between cell R -modules is a chain homotopy equivalence (Whitehead theorem) and every chain complex is quasi-isomorphic to a cell R -module. Then \mathcal{D}_R is equivalent to the ordinary homotopy category of cell R -modules. See [15, 21]. This is a topologist's way of thinking about the appropriate generalization to chain complexes of projective resolutions of modules.

We think of the sphere spectrum S as the analog of R . We think of spectra as analogs of chain complexes, or rather as a first approximation to the definitive analogs, which will be S -modules. We let \mathcal{S} denote the category of spectra. There is a notion of homotopy of maps between spectra, and there is a resulting homotopy category $h\mathcal{S}$. The fundamental invariants of spectra are their homotopy groups, and a map of spectra is a weak equivalence if it induces an isomorphism of homotopy groups. The stable homotopy category, which we denote by $\bar{h}\mathcal{S}$, is obtained by formally inverting the weak equivalences. This is made rigorous by introducing CW spectra. A weak equivalence between CW spectra is a homotopy equivalence and every spectrum is weakly equivalent to a CW spectrum. Then $\bar{h}\mathcal{S}$ is equivalent to the ordinary homotopy category of CW spectra.

Now the category \mathcal{M}_R has an associative and commutative tensor product. If we regard R as a chain complex concentrated in degree zero, then R is a unit for the

tensor product. A differential R -algebra A is a chain complex with a unit $R \rightarrow A$ and product $A \otimes_R A \rightarrow A$ such that the evident associativity and unity diagrams commute. It is commutative if the evident commutativity diagram also commutes. These are, obviously enough, point-set level structures. Algebraists would have trouble taking seriously the idea of an algebra defined in \mathcal{D}_R , with unit and product only defined in that category.

The category \mathcal{S} has a smash product but, in contrast with the tensor product, it is not associative, commutative, or unital. The induced smash product on the stable homotopy category $\bar{h}\mathcal{S}$ is associative and commutative, and it has S as unit. Topologists routinely study ring spectra, which are objects E of $\bar{h}\mathcal{S}$ with a unit $\eta : S \rightarrow E$ and product $\phi : E \wedge E \rightarrow E$ such that the evident unit diagrams commute; that is, $\phi \circ (\eta \wedge \text{id}) = \text{id} = \phi \circ (\text{id} \wedge \eta)$ in $\bar{h}\mathcal{S}$. Similarly, E is associative or commutative if the appropriate diagrams commute in $\bar{h}\mathcal{S}$. Given that the point-set level smash product is not associative or commutative, it would seem at first sight that these up to homotopy notions are the only ones possible.

It is a recent discovery that there is a category \mathcal{M}_S of S -modules that has an associative and commutative smash product \wedge_S [11]. Its objects are spectra with additional structure, and we say that a map of S -modules is a weak equivalence if it is a weak equivalence as a map of spectra. The derived category \mathcal{D}_S is obtained from \mathcal{M}_S by formally inverting the weak equivalences, and \mathcal{D}_S is equivalent to the stable homotopy category $\bar{h}\mathcal{S}$. Again, this is made rigorous by a theory of CW S -modules that is just like the theory of CW spectra. There is a natural unit map $\lambda : S \wedge_S M \rightarrow M$; it is not an isomorphism but it induces an isomorphism on passage to \mathcal{D}_S . There is a modification \star_S of \wedge_S that is defined on S -modules equipped with a map $S \rightarrow M$ and that has S as a strict unit; we will not go into that in the present account, but it is vital to some of the more sophisticated applications.

In the category \mathcal{M}_S , we have a point-set level notion of a ring spectrum R that is defined in terms of maps $\eta : S \rightarrow R$ and $\phi : R \wedge_S R \rightarrow R$ in \mathcal{M}_S . We require that the unit diagrams commute in \mathcal{M}_S , in the sense that $\phi \circ (\eta \wedge_S \text{id}) = \lambda = \phi \circ (\text{id} \wedge_S \eta)$, and we require that the associativity diagram commutes. Such an R is called an A_∞ ring spectrum; if the commutativity diagram also commutes, then R is called an E_∞ ring spectrum. Here “ A_∞ ” stands historically for “associative up to an infinite sequence of higher homotopies”; similarly, “ E_∞ ” stands for “homotopy everything”, meaning that the product is associative and commutative up to all higher coherence homotopies. With the definitions just given the higher homotopies are hidden in the definition of the associative and commutative smash product in \mathcal{M}_S , but the definitions are in fact precisely equivalent to earlier definitions in which the higher homotopies were exhibited in terms of an “operad action”. It is tempting to simply call these objects

associative and commutative ring spectra, but that would be a mistake. These terms have long established meanings, as associative and commutative rings in the stable homotopy category, and the more precise point-set level notions do not make the older notions obsolete: there are plenty of examples of associative or commutative ring spectra that do not admit structures of A_∞ or E_∞ ring spectra. It is part of the new paradigm that one must always be aware of when one is working in the derived category and when one is working on the point-set level.

Now fix an E_∞ ring spectrum R . An R -module M is an S -module together with a map $\mu : R \wedge_S M \rightarrow M$ such that the evident unit and transitivity diagrams commute, the former stating that $\mu \circ (\eta \wedge_S \text{id}) = \lambda$. Let \mathcal{M}_R be the category of R -modules. Again we have a homotopy category $h\mathcal{M}_R$ and a derived category \mathcal{D}_R that is obtained from it by inverting the weak equivalences, by which we mean maps of R -modules that are weak equivalences of underlying spectra. The construction of \mathcal{D}_R is made rigorous by a theory of cell R -modules, the one slight catch being that, unless R is connective, in the sense that its homotopy groups are zero in negative degrees, we cannot insist that cells be attached only to cells of lower dimension, so that our cell R -modules cannot be restricted to be CW R -modules. These categories enjoy all of the good properties that we have described in the special case $R = S$. There is an associative and commutative smash product over R , and it has a unit map $\lambda : R \wedge_R M \rightarrow M$ that becomes an isomorphism on passage to the derived category. We can therefore go on to define A_∞ and E_∞ R -algebras A in terms of point-set level associative and commutative multiplications $A \wedge_R A \rightarrow A$, and we can also define derived category level associative and commutative R -algebras A exactly like the classical associative and commutative ring spectra.

It is the derived category \mathcal{D}_R that we wish to focus on in describing the current state of the art in stable homotopy theory. We can mimic classical commutative algebra in this category. In particular, for an ideal I and multiplicatively closed subset Y in the coefficient ring $R_* = \pi_*(R)$, we will show how to construct quotients M/IM and localizations $M[Y^{-1}]$. When applied with R taken to be the representing spectrum MU for complex cobordism, these constructions specialize to give simple constructions of various spectra that are central to modern stable homotopy theory, such as the Morava K -theory spectra. Moreover, we shall see that these spectra are MU -algebra spectra.

This account is largely a summary of the more complete and technical paper [11], to which the reader is referred for further background and detailed proofs.

1. SPECTRA AND THE STABLE HOMOTOPY CATEGORY

We here give a bare bones summary of the construction of the stable homotopy category, referring to [16] and [11] for details and to [22] for a more leisurely exposition. We aim to give just enough of the basic definitional framework that the reader can feel comfortable with the ideas.

By Brown's representability theorem [6], if E^* is a reduced cohomology theory on based spaces, then there are CW complexes E_n such that, for CW complexes X , $E^n(X)$ is naturally isomorphic to the set $[X, E_n]$ of homotopy classes of based maps $X \rightarrow E_n$. The suspension isomorphism $E^n(X) \cong E^{n+1}(\Sigma X)$ gives rise to a homotopy equivalence $\tilde{\sigma}_n : E_n \rightarrow \Omega E_{n+1}$. The object $E = \{E_n, \tilde{\sigma}_n\}$ is called an Ω -spectrum. A map $f : E \rightarrow E'$ of Ω -spectra is a sequence of homotopy classes of maps $f_n : E_n \rightarrow E'_n$ that are compatible up to homotopy with the equivalences $\tilde{\sigma}_n$ and $\tilde{\sigma}'_n$. The category of Ω -spectra is equivalent to the category of cohomology theories on based spaces and can be thought of as an intuitive first approximation to the stable homotopy category. However, this category does not have a usable theory of cofibration sequences and is not suitable for either point-set level or homotopical work. For that, one needs more precise objects and morphisms that are defined without use of homotopies but that still represent cohomology theories and their maps. More subtly, one needs a coordinate-free setting in order to define smash products sensibly. The n th space E_n relates to the n -sphere and thus to \mathbb{R}^n . Restricting to spaces E_n is very much like restricting to the standard basis of \mathbb{R}^∞ when doing linear algebra.

A coordinate-free spectrum is indexed on the set of finite dimensional subspaces V of a "universe" U , namely a real inner product space isomorphic to the sum \mathbb{R}^∞ of countably many copies of \mathbb{R} . In detail, writing $W - V$ for the orthogonal complement of V in W , a spectrum E assigns a based space EV to each finite dimensional subspace V of U , with structure maps

$$\tilde{\sigma}_{V,W} : EV \xrightarrow{\cong} \Omega^{W-V} EW$$

when $V \subset W$; these maps are required to be *homeomorphisms*. Here $\Omega^W X$ is the function space $F(S^W, X)$ of based maps $S^W \rightarrow X$, where S^W is the one-point compactification of W . The structure maps are required to satisfy an evident transitivity relation when $V \subset W \subset Z$. A map of spectra $f : E \rightarrow E'$ is a collection of maps of based spaces $f_V : EV \rightarrow E'V$ for which each of the following diagrams commutes:

$$\begin{array}{ccc} EV & \xrightarrow{f_V} & E'V \\ \tilde{\sigma}_{V,W} \downarrow & & \downarrow \tilde{\sigma}'_{V,W} \\ \Omega^{W-V} EW & \xrightarrow{\Omega^{W-V} f_W} & \Omega^{W-V} E'W. \end{array}$$

We obtain the category $\mathcal{S} = \mathcal{S}U$ of spectra indexed on U . We obtain an equivalent category if we restrict to any cofinal family of indexing spaces. If we drop the requirement that the maps $\tilde{\sigma}_{V,W}$ be homeomorphisms, we obtain the notion of a prespectrum and the category $\mathcal{P} = \mathcal{P}U$ of prespectra indexed on U . The forgetful functor $\ell : \mathcal{S} \rightarrow \mathcal{P}$ has a left adjoint L . When the structure maps $\tilde{\sigma}$ are inclusions, $(LE)(V)$ is just the union of the spaces $\Omega^{W-V}EW$ for $V \subset W$. We write $\sigma : \Sigma^{W-V}EV \rightarrow EW$ for the adjoint structure maps, where $\Sigma^V X = X \wedge S^V$.

Examples 1.1. Let X be a based space. The suspension prespectrum $\Pi^\infty X$ is the prespectrum whose V th space is $\Sigma^V X$; the structure maps σ are the evident identifications $\Sigma^{W-V}\Sigma^V X \cong \Sigma^W X$. The suspension spectrum of X is $\Sigma^\infty X = L\Pi^\infty X$. Let $QX = \cup \Omega^V \Sigma^V X$, where the union is taken over the inclusions obtained from the adjoints of the cited identifications. Then $(\Sigma^\infty X)(V) = Q(\Sigma^V X)$. The functor Σ^∞ from based spaces to spectra is left adjoint to the functor that assigns the zeroth space $E_0 = E(\{0\})$ to a spectrum E . More generally, for a fixed subspace $Z \subset U$, define $\Pi_Z^\infty X$ to be the analogous prespectrum whose V th space is $\Sigma^{V-Z} X$ if $Z \subset V$ and a point otherwise and define $\Sigma_Z^\infty X = L\Pi_Z^\infty X$. Then Σ_Z^∞ is left adjoint to the functor that sends a spectrum to its Z th space EZ ; these functors are generally called “shift desuspensions”.

Functors on prespectra that do not preserve spectra are extended to spectra by applying the functor L . For example, for a based space X and a prespectrum E , we have the prespectrum $E \wedge X$ specified by $(E \wedge X)(V) = EV \wedge X$. When E is a spectrum, the structure maps for this prespectrum level smash product are not homeomorphisms, and we understand the smash product $E \wedge X$ to be the spectrum $L(\ell E \wedge X)$. Function spectra are easier. We set $F(X, E)(V) = F(X, EV)$ and find that this functor on prespectra preserves spectra. If we topologize the set $\mathcal{S}(E, E')$ as a subspace of the product over V of the function spaces $F(EV, E'V)$ and let \mathcal{T} be the category of based spaces with sets of maps topologized as function spaces, then there result homeomorphisms

$$\mathcal{S}(E \wedge X, E') \cong \mathcal{T}(X, \mathcal{S}(E, E')) \cong \mathcal{S}(E, F(X, E')).$$

Recall that a category is said to be cocomplete if it has all colimits and complete if it has all limits.

Proposition 1.2. The category \mathcal{S} is complete and cocomplete.

Proof. Limits and colimits are defined on prespectra spacewise. Limits preserve spectra, and colimits of spectra are obtained by use of the left adjoint L . \square

We write Y_+ for the union of a space Y and a disjoint basepoint. A homotopy in the category of spectra is a map $E \wedge I_+ \rightarrow E'$. We have cofibration and fibration sequences that are defined exactly as on the space level (e.g. [29]) and enjoy the same homotopical properties. Let $[E, E']$ denote the set of homotopy classes of maps $E \rightarrow E'$; we shall later understand that, when using this notation, E must be of the homotopy type of a CW spectrum. For based spaces X and Y with X compact, we have

$$[\Sigma^\infty X, \Sigma^\infty Y] \cong \operatorname{colim}_n [\Sigma^n X, \Sigma^n Y].$$

Fix a copy of \mathbb{R}^∞ in U . In the equivariant generalization of the present theory, it is essential not to insist that \mathbb{R}^∞ be all of U , but the reader may take $U = \mathbb{R}^\infty$ here. We write $\Sigma_n^\infty = \Sigma_{\mathbb{R}^n}^\infty$. For $n \geq 0$, the sphere spectrum S^n is $\Sigma^\infty S^n$. For $n > 0$, the sphere spectrum S^{-n} is $\Sigma_n^\infty S^0$. We write S for the zero sphere spectrum. The n th homotopy group of a spectrum E is the set $[S^n, E]$ of homotopy classes of maps $S^n \rightarrow E$, and this fixes the notion of a weak equivalence of spectra. The adjunctions of Examples 1.1 make it clear that a map f of spectra is a weak equivalence if and only if each of its component maps f_Z is a weak equivalence of spaces. The stable homotopy category $\bar{h}\mathcal{S}$ is constructed from the homotopy category of spectra by adjoining formal inverses to the weak equivalences, a process that is made rigorous by CW approximation.

The theory of CW spectra is developed by taking sphere spectra as the domains of attaching maps of cells $CS^n = S^n \wedge I$ [16, I§5]. The one major difference from the space level theory of CW complexes is that we have to construct CW spectra as unions $E = \cup E_n$, where E_0 is the trivial spectrum and where we are allowed to attach cells of arbitrary dimension when constructing E_{n+1} from E_n . There results a notion of a cell spectrum. We define a CW spectrum to be a cell spectrum whose cells are attached only to cells of lower dimension. Thus CW spectra have two filtrations, the sequential filtration $\{E_n\}$ that gives the order in which cells are attached, and the skeletal filtration $\{E^q\}$, where E^q is the union of the cells of dimension at most q . We say that a map between CW spectra is cellular if it preserves both filtrations. In fact, by redefining the sequential filtration appropriately, we can always arrange that the sequential filtration is preserved. We have three basic results, whose proofs are very little different from their space level counterparts.

Theorem 1.3 (Whitehead). If E is a CW spectrum and $f : F \rightarrow F'$ is a weak equivalence of spectra, then $f_* : [E, F] \rightarrow [E, F']$ is an isomorphism. Therefore a weak equivalence between CW spectra is a homotopy equivalence.

Theorem 1.4 (Cellular approximation). Let A be a subcomplex of a CW spectrum E , let F be a CW spectrum, and let $f : E \rightarrow F$ be a map whose restriction

to A is cellular. Then f is homotopic relative to A to a cellular map. Therefore any map $E \rightarrow F$ is homotopic to a cellular map, and any two homotopic cellular maps are cellularly homotopic.

Theorem 1.5 (Approximation by CW spectra). For a spectrum E , there is a CW spectrum ΓE and a weak equivalence $\gamma : \Gamma E \rightarrow E$. On the homotopy category $h\mathcal{S}$, Γ is a functor such that γ is natural.

It follows that the stable category $\bar{h}\mathcal{S}$ is equivalent to the homotopy category of CW spectra. Homotopy-preserving functors on spectra that do not preserve weak equivalences are transported to the stable category by first replacing their variables by weakly equivalent CW spectra.

Observe that there has been no mention of space level CW complexes in our development so far. The total lack of hypotheses on the spaces and structural maps of our prespectra allows considerable point-set level pathology, even if, as usual in modern algebraic topology, we restrict attention to compactly generated weak Hausdorff spaces. Recall that a space X is weak Hausdorff if the diagonal subspace is closed in the compactly generated product $X \times X$. More restrictively, a space X is said to be LEC (locally equiconnected) if the inclusion of the diagonal subspace is a cofibration. We record the following list of special kinds of prespectra both to prepare for our discussion of smash products and to compare our definitions with those adopted in the original treatments of the stable homotopy category.

Definition 1.6. A prespectrum D is said to be

- (i) Σ -cofibrant if each $\sigma : \Sigma^{W-V}DV \rightarrow DW$ is a based cofibration (that is, satisfies the based homotopy extension property).
- (ii) CW if each DV is LEC and has the homotopy type of a CW complex.
- (iii) strictly CW if each DV is a based CW complex and the structure maps σ are the inclusions of subcomplexes.

A spectrum E is said to be Σ -cofibrant if it is isomorphic to LD for some Σ -cofibrant prespectrum D ; E is said to be tame if it is of the homotopy type of a Σ -cofibrant spectrum.

If E is a spectrum, then the maps $\tilde{\sigma}$ are homeomorphisms. Therefore the underlying prespectrum ℓE is not Σ -cofibrant unless it is trivial. However, it is a very weak condition on a spectrum that it be tame. We shall see that this weak condition is enough to avoid serious point-set topological problems. If D is a Σ -cofibrant prespectrum, then the maps $\tilde{\sigma}$ are inclusions and therefore $LD(V)$ is just the union of the spaces $\Omega^{W-V}DW$. We have the following relations between CW prespectra

and CW spectra. Remember that CW spectra are defined in terms of spectrum level attaching maps.

Theorem 1.7. If D is a CW prespectrum, then LD has the homotopy type of a CW spectrum. If E is a CW spectrum, then each space EV has the homotopy type of a CW complex and E is homotopy equivalent to LD for some CW prespectrum D . Thus a spectrum has the homotopy type of a CW spectrum if and only if it has the homotopy type of LD for some CW prespectrum D .

In particular, spectra of the homotopy types of CW spectra are tame.

Implicitly or explicitly, early constructions of the stable homotopy category restricted attention to the spectra arising from strict CW prespectra. This is far too restrictive for serious point-set level work, and it is also too restrictive to admit a sensible equivariant analogue. Note that such a category cannot possibly be complete or have well-behaved point-set level function spectra.

One reason for focusing on Σ -cofibrant spectra is that they are built up out of their component spaces in a simple fashion.

Proposition 1.8. If $E = LD$, where D is a Σ -cofibrant prespectrum, then

$$E \cong \operatorname{colim}_V \Sigma_V^\infty DV,$$

where the colimit is computed as the prespectrum level colimit of the maps

$$\Sigma_W^\infty \sigma : \Sigma_V^\infty DV \cong \Sigma_W^\infty \Sigma^{W-V} DV \longrightarrow \Sigma_W^\infty DW.$$

That is, the prespectrum level colimit is a spectrum that is isomorphic to E . The maps of the colimit system are shift desuspensions of based cofibrations.

Another reason is that general spectra can be replaced functorially by weakly equivalent Σ -cofibrant spectra.

Proposition 1.9. There is a functor $K : \mathcal{P}U \longrightarrow \mathcal{P}U$, called the cylinder functor, such that KD is Σ -cofibrant for any prespectrum D , and there is a natural spacewise weak equivalence of prespectra $KD \longrightarrow D$. On spectra E , define $KE = LK\ell E$. Then there is a natural weak equivalence of spectra $KE \longrightarrow E$.

In practice, if one is given a prespectrum D , perhaps indexed only on integers, and one wishes to construct a spectrum from it that retains homotopical information, one forms $E = LKD$. Then

$$\pi_n(E) = \operatorname{colim}_q \pi_{n+q} D_q.$$

If D is an Ω -spectrum that represents a given cohomology theory on spaces, then $E = LKD$ is a genuine spectrum that represents the same theory.

2. SMASH PRODUCTS AND TWISTED HALF-SMASH PRODUCTS

The construction of the smash product of spectra proceeds by internalization of an “external smash product”. The latter is an associative and commutative pairing

$$\mathcal{S}U \times \mathcal{S}U' \rightarrow \mathcal{S}(U \oplus U')$$

for any pair of universes U and U' . It is constructed by starting with the prespectrum level definition

$$(E \wedge E')(V \oplus V') = EV \wedge E'V'.$$

The structure maps fail to be homeomorphisms when E and E' are spectra, and we apply the spectrification functor L to obtain the desired spectrum level smash product.

In order to obtain smash products internal to a single universe U , we exploit the “twisted half-smash product”. The input data for this functor consist of two universes U and U' , an unbased space A with a given map $\alpha : A \rightarrow \mathcal{S}(U, U')$, and a spectrum E indexed on U . The output is the spectrum $A \times E$, which is indexed on U' . It must be remembered that the construction depends on α and not just on A , although different choices of α lead to equivalent functors on the level of stable categories. When A is a point, α is a choice of a linear isometry $f : U \rightarrow U'$ and we write f_* for the twisted half-smash product. For a prespectrum D ,

$$(f_*D)(V') = D(V) \wedge S^{V'-f(V)}, \quad \text{where } V = f^{-1}(V' \cap \text{im } f).$$

For a spectrum E , f_*E is obtained by application of L to the prespectrum level construction. The functor f_* is left adjoint to the more elementary functor f^* specified by $(f^*E')(V) = E'(f(V))$. For general A and α , the intuition is that $A \times E$ is obtained by suitably topologizing the union of the $\alpha(a)_*(E)$. Another intuition is that the twisted half-smash product is a generalization to spectra of the “untwisted” functor $A_+ \wedge X$ on based spaces X . This intuition is made precise by the following “untwisting formula” relating twisted half-smash products and shift desuspensions.

Proposition 2.1. For a map $A \rightarrow \mathcal{S}(U, U')$ and an isomorphism $V \cong V'$, where $V \subset U$ and $V' \subset U'$, there is an isomorphism of spectra

$$A \times \Sigma_V^\infty X \cong A_+ \wedge \Sigma_{V'}^\infty X$$

that is natural in spaces A over $\mathcal{S}(U, U')$ and based spaces X .

The twisted-half smash product functor enjoys essentially the same formal properties as the space level functor $A_+ \wedge X$. The functor $A \times E$ is homotopy-preserving in E , and it therefore preserves homotopy equivalences in the variable E . However, it only preserves homotopies over $\mathcal{S}(U, U')$ in A . Nevertheless, it very often preserves

homotopy equivalences in the variable A . The following central technical result is an easy consequence of Propositions 1.8 and 2.1.

Theorem 2.2. Let $E \in \mathcal{S}U$ be tame and let A be a space over $\mathcal{S}(U, U')$. If $\phi : A' \rightarrow A$ is a homotopy equivalence, then $\phi \times \text{id} : A' \times E \rightarrow A \times E$ is a homotopy equivalence.

Since $A \times E$ is a CW spectrum if A is a CW complex and E is a CW spectrum, this has the following consequence.

Corollary 2.3. Let $E \in \mathcal{S}U$ be a spectrum that has the homotopy type of a CW spectrum and let A be a space over $\mathcal{S}(U, U')$ that has the homotopy type of a CW complex. Then $A \times E$ has the homotopy type of a CW spectrum.

Now, as before, restrict attention to a particular universe U and write $\mathcal{S} = \mathcal{S}U$; again, the reader may think of U as \mathbb{R}^∞ . We are especially interested in twisted half-smash products defined in terms of the following spaces of linear isometries.

Notations 2.4. Let U^j be the direct sum of j copies of U and let $\mathcal{L}(j) = \mathcal{S}(U^j, U)$. The space $\mathcal{L}(0)$ is the point i , where $i : \{0\} \rightarrow U$, and $\mathcal{L}(1)$ contains the identity map $1 = \text{id}_U : U \rightarrow U$. The left action of Σ_j on U^j by permutations induces a free right action of Σ_j on the contractible space $\mathcal{L}(j)$. Define maps

$$\gamma : \mathcal{L}(k) \times \mathcal{L}(j_1) \times \cdots \times \mathcal{L}(j_k) \rightarrow \mathcal{L}(j_1 + \cdots + j_k)$$

by

$$\gamma(g; f_1, \dots, f_k) = g \circ (f_1 \oplus \cdots \oplus f_k).$$

The spaces $\mathcal{L}(j)$ form an operad [18, p.1] with structural maps γ , called the linear isometries operad. Points $f \in \mathcal{L}(j)$ give functors f_* that send spectra indexed on U^j to spectra indexed on U . Applied to a j -fold external smash product $E_1 \wedge \cdots \wedge E_j$, there results an internal smash product $f_*(E_1 \wedge \cdots \wedge E_j)$. All of these smash products become equivalent in the stable category $\bar{h}\mathcal{S}$, but none of them are associative or commutative on the point set level. More precisely, the following result holds.

Theorem 2.5. Let $\mathcal{S}_t \subset \mathcal{S}$ be the full subcategory of tame spectra and let $h\mathcal{S}_t$ be its homotopy category. On \mathcal{S}_t , the internal smash products $f_*(E \wedge E')$ determined by varying $f \in \mathcal{L}(2)$ are canonically homotopy equivalent, and $h\mathcal{S}_t$ is symmetric monoidal under the internal smash product. For based spaces X and tame spectra E , there is a natural homotopy equivalence $E \wedge X \simeq f_*(E \wedge \Sigma^\infty X)$.

This implies formally that we have arrived at a stable situation. As for spaces, the suspension functor Σ is given by $\Sigma E = E \wedge S^1$ and is left adjoint to the loop functor Ω given by $\Omega E = F(S^1, E)$. The cofibre Cf of a map $f : E \rightarrow E'$ of spectra is the pushout $E' \cup_f CE$.

Theorem 2.6. The suspension functor $\Sigma : h\mathcal{S}_t \rightarrow h\mathcal{S}_t$ is an equivalence of categories. A cofibre sequence $E \xrightarrow{f} E' \rightarrow Cf$ in \mathcal{S}_t gives rise to a long exact sequence of homotopy groups

$$\cdots \rightarrow \pi_q(E) \rightarrow \pi_q(E') \rightarrow \pi_q(Cf) \rightarrow \pi_{q-1}(E) \rightarrow \cdots .$$

Proof. For based spaces X , $\Sigma^\infty X$ is naturally isomorphic to $(\Sigma_1^\infty X) \wedge S^1$ because both functors are left adjoint to the zeroth space functor. Thus, for tame spectra E , the previous theorem gives a natural homotopy equivalence

$$E = E \wedge S^0 \simeq f_*(E \wedge \Sigma^\infty S^0) \cong f_*(E \wedge \Sigma_1^\infty S^0) \wedge S^1.$$

Therefore Σ is an equivalence of categories with inverse obtained by smashing with the (-1) -sphere spectrum $S^{-1} = \Sigma_1^\infty S^0$. It follows categorically that $\Omega E \simeq f_*(E \wedge S^{-1})$ and that the unit and counit

$$\eta : E \rightarrow \Omega \Sigma E \quad \text{and} \quad \varepsilon : \Sigma \Omega E \rightarrow E$$

of the adjunction are homotopy equivalences. The last statement is a standard consequence of the fact that maps can now be desuspended. \square

Note that only actual homotopy equivalences, not weak ones, are relevant to the last two results. For this reason among others, $h\mathcal{S}_t$ is a technically convenient halfway house between the homotopy category of spectra and the stable homotopy category.

3. THE CATEGORY OF S -MODULES AND ITS DERIVED CATEGORY

For $f \in \mathcal{L}(j)$ and $E_i \in \mathcal{S}_t$, Theorem 2.2 implies that the inclusion $\{f\} \subset \mathcal{L}(j)$ induces a homotopy equivalence

$$f_*(E_1 \wedge \cdots \wedge E_j) \rightarrow \mathcal{L}(j) \times (E_1 \wedge \cdots \wedge E_j).$$

The proof of Theorem 2.5 above is entirely based on the use of such equivalences. It therefore seems natural to think of

$$\mathcal{L}(j) \times (E_1 \wedge \cdots \wedge E_j)$$

as a canonical j -fold smash product. It is still not associative, but it seems closer to being so. However, to take this idea seriously, we must take note of the difference between E and its “1-fold smash product” $\mathcal{L}(1) \times E$. The space $\mathcal{L}(1)$ is a monoid

under composition, and the formal properties of twisted half-smash products imply a natural isomorphism

$$\mathcal{L}(1) \times (\mathcal{L}(1) \times E) \cong (\mathcal{L}(1) \times \mathcal{L}(1)) \times E,$$

where, on the right, $\mathcal{L}(1) \times \mathcal{L}(1)$ is regarded as a space over $\mathcal{L}(1)$ via the composition product. This product induces a map $\mu : (\mathcal{L}(1) \times \mathcal{L}(1)) \times E \rightarrow \mathcal{L}(1) \times E$, and the inclusion $\{1\} \rightarrow \mathcal{L}(1)$ induces a map $\eta : E \rightarrow \mathcal{L}(1) \times E$. Thus it makes sense to consider spectra E with an action $\xi : \mathcal{L}(1) \times E \rightarrow E$ of the monoid $\mathcal{L}(1)$. It is required that the following diagrams commute:

$$\begin{array}{ccc} (\mathcal{L}(1) \times \mathcal{L}(1)) \times E & \xrightarrow{\mu} & \mathcal{L}(1) \times E & \text{and} & E & \xrightarrow{\eta} & \mathcal{L}(1) \times E \\ \mathcal{L}(1) \times \xi \downarrow & & \downarrow \xi & & \searrow = & & \downarrow \xi \\ \mathcal{L}(1) \times E & \xrightarrow{\xi} & E & & & & E. \end{array}$$

Definition 3.1. An S -module is a spectrum E together with an action ξ of $\mathcal{L}(1)$. A map $f : E \rightarrow E'$ of S -modules is a map of spectra such that the following diagram commutes:

$$\begin{array}{ccc} \mathcal{L}(1) \times E & \xrightarrow{\mathcal{L}(1) \times f} & \mathcal{L}(1) \times E' \\ \xi_E \downarrow & & \downarrow \xi_{E'} \\ E & \xrightarrow{f} & E'. \end{array}$$

We let \mathcal{M}_S denote the category of S -modules.

A number of basic properties of the category of spectra are directly inherited by the category of S -modules.

Theorem 3.2. The category of S -modules is complete and cocomplete, with both limits and colimits created in the underlying category \mathcal{S} . If X is a based space and M is an S -module, then $M \wedge X$ and $F(X, M)$ are S -modules, and the spectrum level fibre and cofibre of a map of S -modules are S -modules.

A homotopy in the category of S -modules is a map $M \wedge I_+ \rightarrow M'$. A map of S -modules is a weak equivalence if it is a weak equivalence as a map of spectra. The derived category \mathcal{D}_S is constructed from the homotopy category of S -modules by adjoining formal inverses to the weak equivalences. There is a theory of CW S -modules that is exactly like the theory of CW spectra, and, again, the construction of \mathcal{D}_S is made rigorous by CW approximation. We have a free functor \mathbb{L} from spectra to S -modules specified by $\mathbb{L}E = \mathcal{L}(1) \times E$. The ‘‘sphere S -modules’’ that we take as

the domains of attaching maps when defining CW S -modules are the free S -modules $\mathbb{L}S^n$. Using the freeness adjunction

$$\mathcal{M}_S(\mathbb{L}E, M) \cong \mathcal{S}(E, M),$$

it is easy to prove Whitehead, cellular approximation, and approximation by CW S -modules theorems exactly like those stated for spectra in Section 1, and \mathcal{D}_S is equivalent to the homotopy category of CW S -modules. There is one catch: although S and all other suspension spectra are S -modules in a natural way, using the untwisting isomorphism of Proposition 2.1 and the projection $\mathcal{L}(1) \rightarrow \{*\}$, S does not have the homotopy type of a CW S -module. However, it is not hard to see that the categories $\bar{h}\mathcal{S}$ and \mathcal{D}_S are equivalent.

Theorem 3.3. The following conclusions hold.

- (i) The free functor $\mathbb{L} : \mathcal{S} \rightarrow \mathcal{M}_S$ carries CW spectra to CW S -modules.
- (ii) The forgetful functor $\mathcal{M}_S \rightarrow \mathcal{S}$ carries S -modules of the homotopy types of CW S -modules to spectra of the homotopy types of CW spectra.
- (iii) Every CW S -module M is homotopy equivalent as an S -module to $\mathbb{L}E$ for some CW spectrum E .
- (iv) If $E \in \mathcal{S}_t$, for example if E is a CW spectrum, then $\eta : E \rightarrow \mathbb{L}E$ is a homotopy equivalence of spectra.
- (v) If M has the homotopy type of a CW S -module, then $\xi : \mathbb{L}M \rightarrow M$ is a homotopy equivalence of S -modules.

Therefore the free and forgetful functors establish an adjoint equivalence between the stable homotopy category $\bar{h}\mathcal{S}$ and the derived category \mathcal{D}_S .

4. THE SMASH PRODUCT OF S -MODULES

One of the most surprising developments of recent years is the discovery of an associative and commutative smash product in the category of S -modules. We proceed to define it. To begin with, observe that the monoid $\mathcal{L}(1) \times \mathcal{L}(1)$ acts from the right on $\mathcal{L}(2)$ and acts from the left on $\mathcal{L}(i) \times \mathcal{L}(j)$, via instances of the structural maps γ of the linear isometries operad. Another instance of γ gives rise to a map

$$(4.1) \quad \bar{\gamma} : \mathcal{L}(2) \times_{\mathcal{L}(1) \times \mathcal{L}(1)} \mathcal{L}(i) \times \mathcal{L}(j) \rightarrow \mathcal{L}(i+j).$$

The space on the left is the balanced product (formally a coequalizer) of the two specified actions by $\mathcal{L}(1) \times \mathcal{L}(1)$. The essential, elementary, point is that this map is a homeomorphism if $i \geq 1$ and $j \geq 1$. To see this, choose linear isometric isomorphisms $s : U \rightarrow U^i$ and $t : U \rightarrow U^j$. Composition on the right with $s \oplus t$

gives vertical homeomorphisms in the commutative diagram

$$\begin{array}{ccc} \mathcal{L}(2) \times_{\mathcal{L}(1) \times \mathcal{L}(1)} \mathcal{L}(i) \times \mathcal{L}(j) & \xrightarrow{\bar{\gamma}} & \mathcal{L}(i+j) \\ \downarrow & & \downarrow \\ \mathcal{L}(2) \times_{\mathcal{L}(1) \times \mathcal{L}(1)} \mathcal{L}(1) \times \mathcal{L}(1) & \xrightarrow{\bar{\gamma}} & \mathcal{L}(2), \end{array}$$

and the lower map $\bar{\gamma}$ is clearly a homeomorphism. Note also that $\mathcal{L}(1)$ acts from the left on $\mathcal{L}(2)$ and that this action commutes with the right action of $\mathcal{L}(1) \times \mathcal{L}(1)$.

Regard $\mathcal{L}(1) \times \mathcal{L}(1)$ as a space over $\mathcal{S}(U^2, U^2)$ via the direct sum of isometries map. If M and N are S -modules, then $\mathcal{L}(1) \times \mathcal{L}(1)$ acts from the left on the external smash product $M \wedge N$ via the map

$$\xi : (\mathcal{L}(1) \times \mathcal{L}(1)) \times (M \wedge N) \cong (\mathcal{L}(1) \times M) \wedge (\mathcal{L}(1) \times N) \xrightarrow{\xi \wedge \xi} M \wedge N.$$

The smash product over S of M and N is simply the balanced product (again, formally a coequalizer)

$$(4.2) \quad M \wedge_S N = \mathcal{L}(2) \times_{\mathcal{L}(1) \times \mathcal{L}(1)} (M \wedge N).$$

The left action of $\mathcal{L}(1)$ on $\mathcal{L}(2)$ induces a left action of $\mathcal{L}(1)$ on $M \wedge_S N$ that gives it a structure of S -module. Use of the transposition $\sigma \in \Sigma_2$ and the commutativity of the external smash product easily gives a commutativity isomorphism

$$\tau : M \wedge_S N \longrightarrow N \wedge_S M.$$

More substantially, there is a natural associativity isomorphism

$$(M \wedge_S N) \wedge_S P \cong M \wedge_S (N \wedge_S P).$$

In fact, using the case $i = 2$ and $j = 1$ of the homeomorphism $\bar{\gamma}$, we obtain isomorphisms

$$\begin{aligned} (M \wedge_S N) \wedge_S P &\cong \mathcal{L}(2) \times_{\mathcal{L}(1)^2} (\mathcal{L}(2) \times_{\mathcal{L}(1)^2} (M \wedge N)) \wedge (\mathcal{L}(1) \times_{\mathcal{L}(1)} P) \\ &\cong (\mathcal{L}(2) \times_{\mathcal{L}(1)^2} \mathcal{L}(2) \times \mathcal{L}(1)) \times_{\mathcal{L}(1)^3} (M \wedge N \wedge P) \\ &\cong \mathcal{L}(3) \times_{\mathcal{L}(1)^3} M \wedge N \wedge P. \end{aligned}$$

The symmetric argument shows that this is also isomorphic to $M \wedge_S (N \wedge_S P)$. In view of the generality of the homeomorphisms (4.1), the argument iterates to give

$$(4.3) \quad M_1 \wedge_S \cdots \wedge_S M_j \cong \mathcal{L}(j) \times_{\mathcal{L}(1)^j} (M_1 \wedge \cdots \wedge M_j),$$

where the iterated smash product on the left is associated in any fashion.

On passage to the derived category \mathcal{D}_S , the smash product of S -modules just constructed can be used interchangeably with the internal smash product on the

stable category $\bar{h}\mathcal{S}$. To see this, one defines the latter by use of a linear isometric isomorphism $f : U^2 \rightarrow U$ (not just an isometry). With this choice, it is not hard to check the following result.

Proposition 4.4. For spectra E and F , there are isomorphisms of S -modules

$$\mathbb{L}E \wedge_S \mathbb{L}F \cong \mathcal{L}(2) \times E \wedge F \cong \mathbb{L}f_*(E \wedge F).$$

For CW S -modules M and N , $M \wedge_S N$ is a CW S -module with one $(p+q)$ -cell for each p -cell of M and q -cell of N .

Together with Theorem 2.6, this gives the starting point towards the proof of the following fundamental theorem.

Theorem 4.5. Let E be a CW spectrum and M be a CW S -module.

(i) For spectra F in \mathcal{S}_t , there is a natural spectral sequence

$$\mathrm{Tor}^{\pi_* S}(\pi_* E, \pi_* F) \Longrightarrow \pi_*((f_*(E \wedge F))).$$

(ii) If $\phi : F \rightarrow F'$ is a weak equivalence between spectra in \mathcal{S}_t , then

$$f_*(\mathrm{id} \wedge \phi) : f_*(E \wedge F) \rightarrow f_*(E \wedge F')$$

is a weak equivalence of spectra.

(iii) For S -modules N in \mathcal{S}_t , there is a natural spectral sequence

$$\mathrm{Tor}^{\pi_* S}(\pi_* M, \pi_* N) \Longrightarrow \pi_*(M \wedge_S N).$$

(iv) If $\phi : N \rightarrow N'$ is a weak equivalence of S -modules in \mathcal{S}_t , then

$$\mathrm{id} \wedge_S \phi : M \wedge_S N \rightarrow M \wedge_S N'$$

is a weak equivalence of S -modules.

(v) The composite natural map of spectra

$$f_*(M \wedge N) \rightarrow \mathcal{L}(2) \times M \wedge N \rightarrow M \wedge_S N$$

is a weak equivalence when the S -module N is in \mathcal{S}_t .

Therefore the equivalence of categories $\mathcal{D}_S \rightarrow \bar{h}\mathcal{S}$ carries the smash product over S to the internal smash product of spectra.

We must still address the question of units.

Proposition 4.6. There is a natural map of S -modules $\lambda : S \wedge_S N \rightarrow N$, and λ is a homotopy equivalence of spectra if N is a CW S -module. Therefore λ induces a natural isomorphism of functors on \mathcal{D}_S .

Proof. When N is the free S -module $\mathbb{L}E = \mathcal{L}(1) \rtimes E$ generated by a spectrum E , λ is given by the map

$$\begin{aligned} S \wedge_S \mathbb{L}E &= \mathcal{L}(2) \rtimes_{\mathcal{L}(1) \times \mathcal{L}(1)} (\mathcal{L}(0) \rtimes S^0) \wedge (\mathcal{L}(1) \rtimes E) \\ &\cong (\mathcal{L}(2) \times_{\mathcal{L}(1) \times \mathcal{L}(1)} \mathcal{L}(0) \times \mathcal{L}(1)) \rtimes (S^0 \wedge E) \\ &\xrightarrow{\bar{\gamma} \times \text{id}} \mathcal{L}(1) \rtimes E = \mathbb{L}E. \end{aligned}$$

Here the map $\bar{\gamma}$, although not a homeomorphism, is a homotopy equivalence. Therefore, by Theorem 2.2, λ is a homotopy equivalence when $E \in \mathcal{S}_t$. For general N , the map just constructed for $\mathbb{L}N$ induces the required map for N by a comparison of coequalizer diagrams. By Theorem 3.3(iii), we conclude directly that λ is a homotopy equivalence when N is a CW S -module. \square

There is one case when λ is an isomorphism. It turns out that the map $\bar{\gamma}$ of (4.1) is a homeomorphism when $i = j = 0$; that is, non-obviously since $\mathcal{L}(1)$ is a monoid but not a group, the domain of (4.1) is then a point. This implies that $S \wedge_S S \cong S$. More generally, it implies that the smash product over S precisely generalizes the smash product of based spaces.

Proposition 4.7. For based spaces X and Y ,

$$\Sigma^\infty(X \wedge Y) \cong \Sigma^\infty X \wedge_S \Sigma^\infty Y.$$

We shall not display the constructions here, but the twisted half-smash product functor $A \rtimes E$ has a right adjoint twisted function spectrum functor $F[A, E']$ and the external smash product has a right adjoint function spectrum functor. Using these functors and appropriate equalizer diagrams, dual to the coequalizer diagrams that were implicit in the definition of \wedge_S , we can construct function S -modules.

Theorem 4.8. There is a function S -module functor $F_S(M, N)$ such that

$$\mathcal{M}_S(L \wedge_S M, N) \cong \mathcal{M}_S(L, F_S(M, N)).$$

Theorem 4.9. Let M be a CW S -module.

(i) If $\phi : N \rightarrow N'$ is a weak equivalence of S -modules, then

$$F_S(\text{id}, \phi) : F_S(M, N) \longrightarrow F_S(M, N')$$

is a weak equivalence of S -modules.

(ii) There is an induced adjunction isomorphism

$$\mathcal{D}_S(L \wedge_S M, N) \cong \mathcal{D}_S(L, F_S(M, N)).$$

(iii) There is a natural weak equivalence of spectra $F_S(M, N) \longrightarrow F(M, f^*N)$.

Therefore the equivalence of categories $\mathcal{D}_S \rightarrow \bar{h}\mathcal{S}$ induced by the forgetful functor from S -modules to spectra carries the function S -module functor F_S to the internal function spectrum functor F .

Proposition 4.10. The adjoint $N \rightarrow F_S(S, N)$ of the unit map $N \wedge_S S \rightarrow N$ becomes a natural equivalence on passage to the derived category \mathcal{D}_S .

5. A_∞ AND E_∞ RING SPECTRA AND THEIR CATEGORIES OF MODULES

Intuitively, A_∞ ring spectra are as near to associative rings with unit as one can get in stable homotopy theory, and E_∞ ring spectra are as near as one can get to commutative rings.

Definition 5.1. An A_∞ ring spectrum is an S -module R together with maps of S -modules $\eta : S \rightarrow R$ and $\phi : R \wedge_S R \rightarrow R$ such that the following diagrams commute:

$$\begin{array}{ccc}
 S \wedge_S R & \xrightarrow{\eta \wedge_S \text{id}} & R \wedge_S R & \xleftarrow{\text{id} \wedge_S \eta} & R \wedge_S S \\
 & \searrow \lambda & \downarrow \phi & & \swarrow \lambda \tau \\
 & & R & &
 \end{array}
 \quad \text{and} \quad
 \begin{array}{ccc}
 R \wedge_S R \wedge_S R & \xrightarrow{\text{id} \wedge_S \phi} & R \wedge_S R \\
 \phi \wedge_S \text{id} \downarrow & & \downarrow \phi \\
 R \wedge_S R & \xrightarrow{\phi} & R;
 \end{array}$$

R is an E_∞ ring spectrum if the following diagram also commutes:

$$\begin{array}{ccc}
 R \wedge_S R & \xrightarrow{\tau} & R \wedge_S R \\
 & \searrow \phi & \swarrow \phi \\
 & & R.
 \end{array}$$

Definition 5.2. Let R be an A_∞ ring spectrum. A (left) R -module M is an S -module together with a map $\mu : R \wedge_S M \rightarrow M$ of S -modules such that the following diagrams commute:

$$\begin{array}{ccc}
 S \wedge_S M & \xrightarrow{\eta \wedge_S \text{id}} & R \wedge_S M \\
 & \searrow \lambda & \downarrow \mu \\
 & & M
 \end{array}
 \quad \text{and} \quad
 \begin{array}{ccc}
 R \wedge_S R \wedge_S M & \xrightarrow{\text{id} \wedge_S \mu} & R \wedge_S M \\
 \phi \wedge_S \text{id} \downarrow & & \downarrow \mu \\
 R \wedge_S M & \xrightarrow{\mu} & M.
 \end{array}$$

A map $f : M \rightarrow M'$ of R -modules is a map of S -modules such that the following diagram commutes:

$$\begin{array}{ccc}
 R \wedge_S M & \xrightarrow{\text{id} \wedge f} & R \wedge_S M' \\
 \mu \downarrow & & \downarrow \mu' \\
 M & \xrightarrow{f} & M'.
 \end{array}$$

We let \mathcal{M}_R denote the category of R -modules.

If R is an E_∞ ring spectrum, then an R -module is the same thing as a left module over R regarded as an A_∞ ring spectrum, exactly as in algebra. From here, we can mimic vast areas of algebra, one particularly striking direction being the development of topological Hochschild homology. However, we shall concentrate on the generalized analog of stable homotopy theory that we obtain by studying the homotopy theory of R -modules for a fixed E_∞ ring spectrum R . Everything that makes sense is also true for A_∞ ring spectra and their left and right modules. The following observation is exactly the same as for S -modules in Theorem 3.2.

Theorem 5.3. The category of R -modules is complete and cocomplete, with both limits and colimits created in the underlying category \mathcal{S} . If X is a based space and M is an R -module, then $M \wedge X$ and $F(X, M)$ are R -modules, and the spectrum level fibre and cofibre of a map of R -modules are R -modules.

Here we have the following complement.

Proposition 5.4. Let M and N be R -modules and let K be an S -module. Then $M \wedge_S K$ and $F_S(K, N)$ are R -modules such that

$$\mathcal{M}_R(M \wedge_S K, N) \cong \mathcal{M}_R(M, F_S(K, N)),$$

and $F_S(M, K)$ is also an R -module.

A homotopy in the category of R -modules is a map $M \wedge I_+ \rightarrow M'$. A map of R -modules is a weak equivalence if it is a weak equivalence as a map of spectra. The derived category \mathcal{D}_R is constructed from the homotopy category $h\mathcal{M}_R$ by adjoining formal inverses to the weak equivalences; again, the process is made rigorous by the approximation of general R -modules by cell R -modules.

Cell theory is based on the free R -module functor \mathbb{F} from spectra to R -modules. It is the composite of the free S -module functor \mathbb{L} and a free R -module functor $\mathcal{M}_S \rightarrow \mathcal{M}_R$, and, as usual, it is characterized by an adjunction

$$\mathcal{M}_R(\mathbb{F}E, M) \cong \mathcal{S}(E, M).$$

This formal property is vital, but, for calculational utility, it is also essential that \mathbb{F} enjoys the following homotopical property. To ensure this, we must assume that R is tame and the unit $\eta : S \rightarrow R$ is a cofibration of S -modules. Fortunately, the cylinder functor K of Proposition 1.9 carries E_∞ ring spectra to weakly equivalent Σ -cofibrant E_∞ ring spectra, and we can also arrange the cofibration hypothesis without loss of generality. We do not assume that R has the homotopy type of a CW spectrum.

Proposition 5.5. If E is a CW spectrum, there is a natural weak homotopy equivalence from the internal smash product $R \wedge E$ to $\mathbb{F}E$. If E is a wedge of sphere spectra, then $\pi_*(\mathbb{F}E)$ is the free $\pi_*(R)$ -module with one generator of degree n for each wedge summand S^n .

Part of the point is that $\mathbb{F}S$ is only weakly equivalent to R , not equal to it. We think of the free R -modules $\mathbb{F}S^n$ as “sphere R -modules”. For cells, we note that the cone functor $CE = E \wedge I$ commutes with \mathbb{F} , so that $C\mathbb{F}S^n \cong \mathbb{F}CS^n$. Thus, via the adjunction, maps out of sphere R -modules and their cones are induced by maps on the spectrum level. Using this, we can simply parrot the theory of cell spectra in the context of R -modules, reducing proofs to the spectrum level via adjunction. We easily obtain the analogs of the Whitehead theorem and of the approximation by cell R -modules theorem, and \mathcal{D}_R is equivalent to the homotopy category of cell R -modules. If R is connective, but not otherwise, we obtain the cellular approximation theorem when we restrict attention to CW R -modules, namely cell R -modules such that cells are only attached to cells of lower dimension.

The category \mathcal{D}_R has all homotopy limits and colimits; they are created as the corresponding constructions on the underlying diagrams of spectra. Thus we have enough information to quote the categorical form of Brown’s representability theorem given in [6]. Adams’ analog [3] for functors defined only on finite CW spectra also applies in our context, with the same proof.

Theorem 5.6 (Brown). A contravariant functor $k : \mathcal{D}_R \rightarrow \text{Sets}$ is representable in the form $k(M) \cong \mathcal{D}_R(M, N)$ for some R -module N if and only if k converts wedges to products and converts homotopy pushouts to weak pullbacks.

Theorem 5.7 (Adams). A contravariant group-valued functor k on the homotopy category of finite cell R -modules is representable in the form $k(M) \cong \mathcal{D}_R(M, N)$ for some R -module N if and only if k converts finite wedges to direct products and converts homotopy pushouts to weak pullbacks of underlying sets.

In fact, Brown’s theorem is the kind of formal result that can be derived in any (closed) model category in the sense of Quillen (see [8] for a good exposition), and we have the following result. Serre fibrations of spectra are maps that satisfy the covering homotopy property with respect to the set of cone spectra

$$\left\{ \Sigma_q^\infty CS^n \mid q \geq 0 \text{ and } n \geq 0 \right\}.$$

Relative cell R -modules $M \rightarrow N$ are constructed exactly like cell R -modules, except that one starts the inductive construction of $N = \cup N_n$ with $N_0 = M$.

We write Cofibrations and Fibrations here to avoid confusion with cofibrations (HEP) and fibrations (CHP); the ambiguous use of the same term for both the classical and the model theoretic concepts is one of the banes of the literature.

Theorem 5.8. The category of R -modules is a model category. Its weak equivalences and Fibrations are those maps of R -modules that are weak equivalences or Serre fibrations of spectra. Its Cofibrations are the retracts of relative cell R -modules.

6. THE SMASH PRODUCT OF R -MODULES AND FUNCTION R -MODULES

Continuing to work with our fixed E_∞ ring spectrum R , we mimic the definition of tensor products of modules over algebras.

Definition 6.1. For R -modules M and N , define $M \wedge_R N$ to be the coequalizer displayed in the following diagram of S -modules:

$$M \wedge_S R \wedge_S N \begin{array}{c} \xrightarrow{\mu \wedge_S \text{id}} \\ \xrightarrow{\text{id} \wedge_S \nu} \end{array} M \wedge_S N \longrightarrow M \wedge_R N.$$

where μ and ν are the given actions of R on M and N . Then $M \wedge_R N$ has a canonical R -module structure induced from the R -module structure of M or, equivalently, N .

Of course, S is an E_∞ ring spectrum and our new $M \wedge_S N$ coincides with our old $M \wedge_S N$. The functor \wedge_R preserves colimits in each of its variables, and smash products with spaces commute with \wedge_R , in the sense that

$$X \wedge (M \wedge_R N) \cong (X \wedge M) \wedge_R N \text{ and } (M \wedge_R N) \wedge X \cong M \wedge_R (N \wedge X).$$

Therefore the functor \wedge_R commutes with cofibre sequences in each of its variables. We have analogous relations with smash products over S and an adjunction that can be thought of as completing Proposition 5.4.

Proposition 6.2. For an S -module K ,

$$K \wedge_S (M \wedge_R N) \cong (K \wedge_S M) \wedge_R N \text{ and } (M \wedge_R N) \wedge_S K \cong M \wedge_R (N \wedge_S K).$$

Moreover,

$$\mathcal{M}_S(M \wedge_R N, K) \cong \mathcal{M}_R(N, F_S(M, K)).$$

The associativity and commutativity of the smash product over S is inherited by the smash product over R .

Theorem 6.3. There is a natural commutativity isomorphism of R -modules

$$\tau : M \wedge_R N \longrightarrow N \wedge_R M.$$

There is also a natural associativity isomorphism of R -modules

$$(M \wedge_R N) \wedge_R P \cong M \wedge_R (N \wedge_R P).$$

The action $\nu : R \wedge_S N \longrightarrow N$ of an R -module N factors through a natural unit map of R -modules $\lambda : R \wedge_R N \longrightarrow N$. If N is a cell R -module, then $\lambda : R \wedge_R N \rightarrow N$ is a homotopy equivalence of spectra and thus a weak equivalence of R -modules.

We can deduce not only formal but also homotopical properties of \wedge_R from corresponding properties of \wedge_S . For example, the proof of the unit equivalence just stated reduces to the case of sphere R -modules, where the conclusion is a consequence of the following result. As in Section 4, we use an isomorphism of universes $f : U \oplus U \rightarrow U$ to define the internal smash product $f_*(E \wedge F)$.

Proposition 6.4. Let E and F be spectra and let N be an R -module. There is a natural isomorphism of R -modules

$$\mathbb{F}E \wedge_R N \cong \mathbb{L}E \wedge_S N.$$

There is also a natural isomorphism of R -modules

$$\mathbb{F}E \wedge_R \mathbb{F}F \cong \mathbb{F}f_*(E \wedge F).$$

If M and N are cell R -modules, then $M \wedge_R N$ is a cell R -module with one $(p+q)$ -cell for each p -cell of M and q -cell of N .

Similarly, this proposition and Theorem 4.5 imply the following analog of the latter.

Theorem 6.5. If M is a cell R -module and $\phi : N \longrightarrow N'$ is a weak equivalence of R -modules in \mathcal{S}_t , then

$$\text{id} \wedge_R \phi : M \wedge_R N \longrightarrow M \wedge_R N'$$

is a weak equivalence of R -modules.

We construct \wedge_R as a functor $\mathcal{D}_R \times \mathcal{D}_R \rightarrow \mathcal{D}_R$ by approximating the variables by cell R -modules, and we need only approximate one of the variables provided that the other is tame. That is, if $N \in \mathcal{S}_t$, then the derived smash product of M and N is represented by $\Gamma M \wedge_R N$.

We have a function spectrum functor F_R to go with our smash product. It is defined as the equalizer of a certain pair of maps $F_S(M, N) \longrightarrow F_S(R \wedge_S M, N)$. The details are dictated by the expected adjunction. Again, $F_R(M, N)$ inherits a structure of R -module from M or, equivalently, N .

Proposition 6.6. For R -modules N and P and an S -module K ,

$$\mathcal{M}_R(K \wedge_S N, P) \cong \mathcal{M}_S(K, F_R(N, P)).$$

If M is an R -module, then

$$\mathcal{M}_R(M \wedge_R N, P) \cong \mathcal{M}_R(M, F_R(N, P)).$$

Therefore

$$F_R(M \wedge_R N, P) \cong F_R(M, F_R(N, P)).$$

Formal arguments from the adjunction, as in algebra, give a natural associative and unital composition pairing

$$(6.7) \quad \pi : F_R(N, P) \wedge_R F_R(M, N) \longrightarrow F_R(M, P).$$

Parenthetically, we note that this gives rise to a host of examples of A_∞ ring spectra; in fact, R itself need only be an A_∞ ring spectrum in the following result.

Proposition 6.8. For R -modules M and N , $F_R(M, M)$ is an A_∞ ring spectrum and $F_R(M, N)$ is an $(F_R(N, N), F_R(M, M))$ -bimodule.

Proposition 6.9. Let E be a spectrum and M be an R -module. There is a natural isomorphism of R -modules

$$F_R(\mathbb{F}E, M) \cong F_S(\mathbb{L}E, M).$$

The functor $F_R(M, N)$ converts colimits and cofibre sequences in M to limits and fibre sequences. It preserves fibre sequences in N . Using the previous result to deal with sphere R -modules, we obtain the analog of Theorem 6.5.

Proposition 6.10. If M is a cell R -module and $\phi : N \longrightarrow N'$ is a weak equivalence of R -modules, then

$$F_R(\text{id}, \phi) : F_R(M, N) \longrightarrow F_R(M, N')$$

is a weak equivalence of R -modules.

In the derived category \mathcal{D}_R , $F_R(M, N)$ means $F_R(\Gamma M, N)$, where ΓM is a cell approximation of M . The unit equivalence for the smash product implies its function module analog.

Corollary 6.11. The adjoint $N \rightarrow F_R(R, N)$ of the unit map $R \wedge_R N \rightarrow N$ induces a natural isomorphism of functors on the derived category \mathcal{D}_R .

Summarizing, we obtain the following derived category level conclusion.

Theorem 6.12. The derived category \mathcal{D}_R is symmetric monoidal under the product derived from \wedge_R , and

$$\mathcal{D}_R(M \wedge_R N, P) \cong \mathcal{D}_R(M, F_R(N, P)).$$

There is a formal theory of duality (explained in [16, Ch. III]) that now applies to \mathcal{D}_R . We define the dual of an R -module M to be $D_R M = F_R(M, R)$. We have an evaluation map $\varepsilon : D_R M \wedge_R M \rightarrow R$ and a map $\eta : R \rightarrow F_R(M, M)$, namely the adjoint of $\lambda : R \wedge_R M \rightarrow M$. There is also a natural map

$$(6.13) \quad \nu : F_R(L, M) \wedge_R N \rightarrow F_R(L, M \wedge_R N).$$

By composition with $F_R(\text{id}, \lambda)$, ν specializes to a map

$$(6.14) \quad \nu : D_R M \wedge_R M \rightarrow F_R(M, M).$$

We say that M is “finite”, or “strongly dualizable”, if it has a coevaluation map $\eta : R \rightarrow M \wedge_R D_R M$ such that the following diagram commutes in \mathcal{D}_R :

$$(6.15) \quad \begin{array}{ccc} R & \xrightarrow{\eta} & M \wedge_R D_R M \\ \eta \downarrow & & \downarrow \tau \\ F_R(M, M) & \xleftarrow{\nu} & D_R M \wedge_R M. \end{array}$$

The definition has many purely formal implications. The map ν of (6.14) is an isomorphism in \mathcal{D}_R if either L or N is finite. The map ν of (6.15) is an isomorphism if and only if M is finite, and the coevaluation map η is then the composite $\tau\nu^{-1}\eta$ in (6.16). The natural map

$$\rho : M \rightarrow D_R D_R M$$

is an isomorphism if M is finite. The natural map

$$\wedge : F_R(M, N) \wedge_R F_R(M', N') \rightarrow F_R(M \wedge_R M', N \wedge_R N')$$

is an isomorphism if M and M' are finite or if M is finite and $N = R$.

Say that a cell R -module N is a wedge summand up to homotopy of a cell R -module M if there is a homotopy equivalence of R -modules between M and $N \vee N'$ for some cell R -module N' . In contrast with the usual stable homotopy category, when M is a finite cell R -module it does not follow that N has the homotopy type of a finite cell R -module.

Theorem 6.16. A cell R -module is finite in the sense just defined if and only if it is a wedge summand up to homotopy of a finite cell R -module.

The analogy with finitely generated projective modules in algebra should be clear.

7. TOR AND EXT IN TOPOLOGY AND ALGEBRA

Still restricting for definiteness to an E_∞ ring spectrum R and its modules, we define Tor and Ext groups as the homotopy groups of derived smash product and function modules.

Definition 7.1. For R -modules M and N , define

$$\mathrm{Tor}_n^R(M, N) = \pi_n(M \wedge_R N)$$

and

$$\mathrm{Ext}_R^n(M, N) = \pi_{-n}(F_R(M, N)).$$

Note that $\mathrm{Tor}_*^R(M, N)$ and $\mathrm{Ext}_R^*(M, N)$ are $\pi_*(R)$ -modules.

We emphasize that the smash product and function spectra are understood to be taken in the derived category \mathcal{D}_R . At this point in our exposition, we act as traditional topologists, taking it for granted that all spectra and modules are to be approximated as cell modules, without change of notation, whenever necessary. Various properties reminiscent of those of the classical Tor and Ext functors follow directly from the definition and the results of the previous sections. The intuition is that the definition gives an analogue of the differential Tor and Ext functors (alias hyperhomology and cohomology functors) in the context of differential graded modules over differential graded algebras. In particular, the grading should not be thought of as the resolution grading of the classical torsion product, but rather as a total grading that sums a resolution degree and an internal degree; this idea will be made precise by the grading of the spectral sequences that we shall describe for the calculation of these functors.

Proposition 7.2. $\mathrm{Tor}_*^R(M, N)$ satisfies the following properties.

- (i) If R , M , and N are connective, then $\mathrm{Tor}_n^R(M, N) = 0$ for $n < 0$.
- (ii) A cofibre sequence $N' \rightarrow N \rightarrow N''$ gives rise to a long exact sequence

$$\cdots \rightarrow \mathrm{Tor}_n^R(M, N') \rightarrow \mathrm{Tor}_n^R(M, N) \rightarrow \mathrm{Tor}_n^R(M, N'') \rightarrow \mathrm{Tor}_{n-1}^R(M, N') \rightarrow \cdots$$
- (iii) $\mathrm{Tor}_*^R(M, R) \cong \pi_*(M)$ and, for a spectrum X ,

$$\mathrm{Tor}_*^R(M, \mathbb{F}X) \cong \pi_*(M \wedge X).$$

- (iv) The functor $\mathrm{Tor}_*^R(M, ?)$ carries wedges to direct sums.

The commutativity and associativity relations for the smash product imply various further properties. We content ourselves with the following examples:

$$\mathrm{Tor}_*^R(M, N) \cong \mathrm{Tor}_*^R(N, M)$$

and

$$\mathrm{Tor}_*^R(M \wedge_R N, P) \cong \mathrm{Tor}_*^R(M, N \wedge_R P).$$

Say that a spectrum N is coconnective if $N_n = 0$ for $n > 0$.

Proposition 7.3. $\text{Ext}_R^*(M, N)$ satisfies the following properties.

- (i) If R and M are connective and N is coconnective, then $\text{Ext}_R^n(M, N) = 0$ for $n > 0$.
- (ii) Cofibre sequences $N' \rightarrow N \rightarrow N''$ and $M' \rightarrow M \rightarrow M''$ give rise to long exact sequences

$$\cdots \rightarrow \text{Ext}_R^n(M, N') \rightarrow \text{Ext}_R^n(M, N) \rightarrow \text{Ext}_R^n(M, N'') \rightarrow \text{Ext}_R^{n+1}(M, N') \rightarrow \cdots$$

and

$$\cdots \rightarrow \text{Ext}_R^n(M'', N) \rightarrow \text{Ext}_R^n(M, N) \rightarrow \text{Ext}_R^n(M', N) \rightarrow \text{Ext}_R^{n+1}(M'', N) \rightarrow \cdots .$$

- (iii) $\text{Ext}_R^*(R, N) \cong \pi_{-*}(N)$ and, for a spectrum X ,

$$\text{Ext}_R^*(\mathbb{F}X, N) \cong \pi_{-*}(F(X, N)).$$

- (iv) The functor $\text{Ext}_R^*(?, N)$ carries wedges to products.

Passing to homotopy groups from the pairings (6.7), we obtain the following further property. As usual, for a spectrum E , abbreviate

$$E_n = \pi_n(E) = E^{-n}.$$

Proposition 7.4. There is a natural, associative, and unital system of pairings of R^* -modules

$$\pi^* : \text{Ext}_R^*(M, N) \otimes_{R^*} \text{Ext}_R^*(L, M) \longrightarrow \text{Ext}_R^*(L, N).$$

The formal duality theory of the previous section implies the following result, together with various other such isomorphisms.

Proposition 7.5. For a finite cell R -module M and any R -module N ,

$$\text{Tor}_n^R(D_R M, N) \cong \text{Ext}_R^{-n}(M, N).$$

Thinking of the derived category \mathcal{D}_R as a stable homotopy category, we may change notations and reinterpret the functors Tor and Ext as prescribing homology and cohomology theories in this category.

Definition 7.6. Let M and E be R -modules. Define

$$E_n^R(M) = \pi_n(E \wedge_R M) \text{ and } E_R^n(M) = \pi_{-n}(F_R(M, E)).$$

The properties of Tor and Ext translate directly to statements about homology and cohomology. All of the standard homotopical machinery is available to us, and the previous result now takes the form of Spanier-Whitehead duality.

Corollary 7.7. For a finite cell R -module M and any R -module E ,

$$E_n^R(D_R M) \cong E_R^{-n}(M).$$

Since the equivalence between the classical stable homotopy category and the derived category of S -modules preserves smash products and function spectra, we obtain all of the usual homology and cohomology theories by taking $R = S$.

We also obtain the classical algebraic Tor and Ext groups as special cases, by specialization to Eilenberg-MacLane spectra. Thus let R be a discrete commutative ring for a moment. Recall that HR denotes a spectrum whose zeroth homotopy group is R and whose remaining homotopy groups are zero. It follows from multiplicative infinite loop space theory [20] that the Eilenberg-MacLane spectrum $HR = K(R, 0)$ is an E_∞ ring spectrum. Analogously, if M is an R -module, then HM can be constructed as an HR -module. We shall see a quick and easy construction shortly. Granting this, we have the following result.

Theorem 7.8. For a discrete commutative ring R and R -modules M and N ,

$$\mathrm{Tor}_*^R(M, N) \cong \mathrm{Tor}_*^{HR}(HM, HN)$$

and

$$\mathrm{Ext}_R^*(M, N) \cong \mathrm{Ext}_{HR}^*(HM, HN).$$

Under the second isomorphism, the topologically defined pairing

$$\mathrm{Ext}_{HR}^*(HM, HN) \otimes_R \mathrm{Ext}_{HR}^*(HL, HM) \longrightarrow \mathrm{Ext}_{HR}^*(HL, HN)$$

coincides with the algebraic Yoneda product.

The proof is clear enough: we just check the axioms for Tor and Ext.

We can elaborate this result to an equivalence of derived categories. Recall from [28] or [15, Ch.III] that the derived category \mathcal{D}_R is obtained from the homotopy category of chain complexes over R by formally inverting the quasi-isomorphisms, exactly as we obtained the category \mathcal{D}_{HR} from the homotopy category of HR -modules by inverting the weak equivalences. The algebraic theory of cell and CW chain complexes over R in the latter source makes the analogy precise and gives a treatment of tensor products and Hom functors in \mathcal{D}_R that exactly parallels our treatment of \wedge_{HR} and F_{HR} . The proof of the equivalence is quite easy. The category \mathcal{D}_{HR} is equivalent to the homotopy category of CW HR -modules and cellular maps. It is a simple matter to see that CW HR -modules have associated chain complexes. This gives a functor $\mathcal{D}_{HR} \rightarrow \mathcal{D}_R$. An inverse functor Φ is obtained by applying Brown's representability theorem. Indeed, for a given chain complex X , the functor k on \mathcal{D}_{HR} specified by $k(M) = \mathcal{D}_R(C_*(M), X)$ satisfies the hypotheses of that result, and we let $\Phi(X)$ represent this functor. Specialization to R -modules regarded as chain complexes

concentrated in degree zero gives the promised construction of Eilenberg-MacLane HR -modules from R -modules.

Theorem 7.9. The cellular chain functor C_* on HR -modules induces an equivalence of categories $\mathcal{D}_{HR} \rightarrow \mathcal{D}_R$. The functor C_* satisfies $H_*(C_*(M)) \cong \pi_*(M)$ and carries the functors \wedge_{HR} and F_{HR} to the functors \otimes_R and Hom_R . The inverse equivalence Φ satisfies $\pi_*(\Phi(X)) \cong H_*(X)$ and carries the functors \otimes_R and Hom_R to the functors \wedge_{HR} and F_{HR} .

Proof. By construction, we have an adjunction

$$\mathcal{D}_R(C_*(M), X) \cong \mathcal{D}_{HR}(M, \Phi(X)),$$

and one checks that its unit and counit are isomorphisms. The statements relating \wedge_{HR} and F_{HR} to \otimes_R and Hom_R are all consequences of the fact that if M and N are CW HR -modules, then $M \wedge_{HR} N$ is a CW HR -module such that

$$C_*(M \wedge_{HR} N) \cong C_*(M) \otimes_R C_*(N). \quad \square$$

8. UNIVERSAL COEFFICIENT AND KÜNNETH SPECTRAL SEQUENCES

Returning to our general E_∞ ring spectrum R , we find spectral sequences for the calculation of our Tor and Ext groups that are analogous to the Eilenberg-Moore (or hyperhomology) spectral sequences in differential homological algebra. Compare [9, 13, 15]. They may be viewed as giving universal coefficient and Künneth spectral sequences for homology and cohomology theories on R -modules, and they specialize to give such spectral sequences for homology and cohomology theories on spectra.

Theorem 8.1. For R -modules M and N , there are natural spectral sequences of differential R_* -modules

$$E_{p,q}^2 = \text{Tor}_{p,q}^{R_*}(M_*, N_*) \implies \text{Tor}_{p+q}^R(M, N)$$

and

$$E_2^{p,q} = \text{Ext}_{R_*}^{p,q}(M^*, N^*) \implies \text{Ext}_R^{p+q}(M, N).$$

Moreover, the pairing $F_R(M, N) \wedge_R F_R(L, M) \rightarrow F_R(L, N)$ induces a pairing of spectral sequences that coincides with the algebraic Yoneda pairing

$$\text{Ext}_{R_*}^{*,*}(M^*, N^*) \otimes_{R_*} \text{Ext}_{R_*}^{*,*}(L^*, M^*) \longrightarrow \text{Ext}_{R_*}^{*,*}(L^*, N^*)$$

on the E_2 -level and that converges to the induced pairing of Ext groups.

The Tor spectral sequence is of standard homological type, with

$$d_{p,q}^r : E_{p,q}^r \longrightarrow E_{p-r,q+r-1}^r.$$

It lies in the right half-plane, and it converges strongly. The Ext spectral sequence is of standard cohomological type, with

$$d_r : E_r^{p,q} \rightarrow E_r^{p+r,q-r+1}.$$

It lies in the right half plane. In the language of Boardman [5] (see also [12, App B]), it is conditionally convergent. It therefore converges strongly if, for each fixed (p, q) , there are only finitely many r such that d_r is non-zero on $E_r^{p,q}$.

Setting $M = \mathbb{F}X$ in the two spectral sequences of Theorem 8.1, we obtain a universal coefficient spectral sequence. We have written the stars to indicate the way the grading is usually thought of in cohomology.

Theorem 8.2 (Universal coefficient). For an R -module N and any spectrum X , there are spectral sequences of the form

$$\mathrm{Tor}_{*,*}^{R_*}(R_*(X), N_*) \Longrightarrow N_*(X)$$

and

$$\mathrm{Ext}_{R_*}^{*,*}(R_{-*}(X), N^*) \Longrightarrow N^*(X).$$

Of course, replacing R and N by Eilenberg-Mac Lane spectra HR and HN for a ring R and R -module N , we obtain the classical universal coefficient theorems. Replacing N by $\mathbb{F}Y$ and by $F_R(\mathbb{F}Y, R)$ in the two universal coefficient spectral sequences, we arrive at Künneth spectral sequences.

Theorem 8.3 (Künneth). For any spectra X and Y , there are spectral sequences of the form

$$\mathrm{Tor}_{*,*}^{R_*}(R_*(X), R_*(Y)) \Longrightarrow R_*(X \wedge Y)$$

and

$$\mathrm{Ext}_{R_*}^{*,*}(R_{-*}(X), R^*(Y)) \Longrightarrow R^*(X \wedge Y).$$

Adams [1] first observed that one can derive Künneth spectral sequences from universal coefficient spectral sequences, and he observed that, by duality, the four spectral sequences of Theorems 8.2 and 8.3 imply two more universal coefficient and two more Künneth spectral sequences. He derived spectral sequences of this sort under the hypothesis that his given ring spectrum E is the colimit of finite subspectra E_α such that $H^*(E_\alpha; E^*)$ is E^* -projective and the Atiyah-Hirzebruch spectral sequence converging from $H^*(E_\alpha; E^*)$ to $E^*(E_\alpha)$ satisfies $E_2 = E_\infty$. Of course, this is an ad hoc calculational hypothesis that requires case-by-case verification. It covers some cases that are not covered by the results above, and conversely. The cited

paper of Adams, and his later book [2], are prime sources for the first flowering of stable homotopy theory. While some of their foundational parts may be obsolete, their applications and calculational parts certainly are not.

The following generalized Künneth theorem admits a number of variants; see [11].

Theorem 8.4. Let E and R be E_∞ ring spectra and M and N be R -modules. Assume that E or R is of the homotopy type of a CW spectrum. Then there is a spectral sequence of differential $E_*(R)$ -modules of the form

$$\mathrm{Tor}_{p,q}^{E_*(R)}(E_*(M), E_*(N)) \implies E_{p+q}(M \wedge_R N).$$

9. ALGEBRAIC CONSTRUCTIONS IN THE DERIVED CATEGORY OF R -MODULES

If we replace the pair (S, R) by a pair (R, A) in Definition 5.1, we arrive at the notion of an A_∞ or E_∞ algebra A over an E_∞ ring spectrum R . For example, the A_∞ ring spectra $F_R(M, M)$ of Proposition 6.8 are actually A_∞ R -algebras. Again, if A is an algebra over a discrete commutative ring R , then HA is an A_∞ HR -algebra. Proceeding in this line, we can, for instance, construct R -modules whose homotopy groups realize the Hochschild homology of A with coefficients in (A, A) -bimodules.

However, we now proceed in a more homotopical direction, thinking of the derived category of R -modules as an analog of the stable homotopy category. From this point of view, we have the notion of an R -algebra up to homotopy, which is just like the classical notion of a ring spectrum in the stable homotopy category.

Definition 9.1. An R -algebra A is an R -module A with unit $\eta : R \rightarrow A$ and product $\phi : A \wedge_R A \rightarrow A$ in \mathcal{D}_R such that the following left and right unit diagram commutes in \mathcal{D}_R .

$$\begin{array}{ccccc} R \wedge_R A & \xrightarrow{\eta \wedge \mathrm{id}} & A \wedge_R A & \xleftarrow{\mathrm{id} \wedge \eta} & A \wedge_R R \\ & \searrow \simeq & \downarrow \phi & \swarrow \simeq & \\ & \lambda & A & \lambda \tau & \end{array}$$

A is associative or commutative if the appropriate diagram commutes in \mathcal{D}_R .

Lemma 9.2. If A and B are R -algebras, then so is $A \wedge_R B$. If A and B are associative or commutative, then so is $A \wedge_R B$.

By neglect of structure, an R -algebra A is a ring spectrum in the sense of classical stable homotopy theory; its unit is the composite of the unit of R and the unit of A and its product is the composite of the product of A and the canonical map

$$A \wedge A \simeq A \wedge_S A \rightarrow A \wedge_R A.$$

Similarly, for an R -algebra A , we have the evident homotopical notion of an A -module. These structures play a role in the study of \mathcal{D}_R analogous to the role played by ring spectra and their modules in classical stable homotopy theory. When $R = S$, S -algebras and their modules are equivalent to classical ring spectra and their modules.

We show in this section how to construct quotients M/IM and localizations $M[Y^{-1}]$ of modules over an E_∞ ring spectrum R and indicate in the next section when these constructions inherit a structure of R -algebra from an R -algebra structure on M . When specialized to MU , these results give highly structured versions of spectra that in the past were constructed by means of the Baas-Sullivan theory of manifolds with singularities or the Landweber exact functor theorem. At least at odd primes, the results give an entirely satisfactory, and very simple, treatment of algebra structures on the resulting MU -modules.

We are interested in homotopy groups, and we make use of the isomorphisms

$$(9.3) \quad M_n = h\mathcal{S}(S^n, M) \cong h\mathcal{M}_S(\mathbb{L}S^n, M) \cong h\mathcal{M}_R(\mathbb{F}S^n, M)$$

to represent elements as maps of R -modules. For $x \in R_n$, the composite map of R -modules

$$(9.4) \quad \mathbb{F}S^n \wedge_R M \xrightarrow{x \wedge \text{id}} R \wedge_R M \xrightarrow{\lambda} M$$

is a module theoretic version of the map $x \cdot : \Sigma^n M \rightarrow M$, and we agree to write $\Sigma^n M$ for $\mathbb{F}S^n \wedge_R M$ in this section. By Proposition 6.4, $\mathbb{F}S^n \wedge_R M$ is isomorphic as an R -module to $\mathbb{L}S^n \wedge_S M$ and, by Theorem 4.5, $\mathbb{L}S^n \wedge_S M$ is weakly equivalent as a spectrum to $S^n \wedge M$. Therefore, the R -module $\Sigma^n M$ is a model for the spectrum level suspension of M .

Definition 9.5. Define M/xM to be the cofibre of the map (9.4) and let $\iota : M \rightarrow M/xM$ be the canonical map. Inductively, for a finite sequence $\{x_1, \dots, x_n\}$ of elements of R_* , define

$$M/(x_1, \dots, x_n)M = N/x_n N, \quad \text{where } N = M/(x_1, \dots, x_{n-1})M.$$

For a (countably) infinite sequence $X = \{x_i\}$, define M/XM to be the telescope of the $M/(x_1, \dots, x_n)M$, where the telescope is taken with respect to the successive canonical maps ι .

Clearly we have a long exact sequence

$$(9.6) \quad \cdots \rightarrow \pi_{q-n}(M) \xrightarrow{x \cdot} \pi_q(M) \xrightarrow{\iota_*} \pi_q(M/xM) \rightarrow \pi_{q-n-1}(M) \rightarrow \cdots$$

If x is not a zero divisor for $\pi_*(M)$, then ι_* induces an isomorphism of R_* -modules

$$(9.7) \quad \pi_*(M)/x \cdot \pi_*(M) \cong \pi_*(M/xM).$$

If $\{x_1, \dots, x_n\}$ is a regular sequence for $\pi_*(M)$, in the sense that x_i is not a zero divisor for $\pi_*(M)/(x_1, \dots, x_{i-1})\pi_*(M)$ for $1 \leq i \leq n$, then

$$(9.8) \quad \pi_*(M)/(x_1, \dots, x_n)\pi_*(M) \cong \pi_*(M/(x_1, \dots, x_n)M),$$

and similarly for a possibly infinite regular sequence $X = \{x_i\}$. The following result implies that M/XM is independent of the ordering of the elements of the set X . We write R/X instead of R/XR .

Lemma 9.9. For a set X of elements of R_* , there is a natural weak equivalence

$$(R/X) \wedge_R M \longrightarrow M/XM.$$

In particular, for a finite set $X = \{x_1, \dots, x_n\}$,

$$R/(x_1, \dots, x_n) \simeq (R/x_1) \wedge_R \cdots \wedge_R (R/x_n).$$

If I denotes the ideal generated by X , then it is reasonable to define

$$(9.10) \quad M/IM = M/XM.$$

However, this notation must be used with caution since, if we fail to restrict attention to regular sequences X , the homotopy type of M/XM will depend on the set X and not just on the ideal it generates. For example, quite different modules are obtained if we repeat a generator x_i of I in our construction.

We next construct localizations of R -modules at countable multiplicatively closed subsets Y of R_* . Let $\{y_i\}$ be any cofinal sequence of Y , with $y_i \in R_{n_i}$, so that every $y \in Y$ divides some y_i . We may represent y_i by an R -map $\mathbb{F}S^0 \longrightarrow \mathbb{F}S^{-n_i}$, which we also denote by y_i . Let $q_0 = 0$ and, inductively, $q_i = q_{i-1} + n_i$. The R -map

$$\mathbb{F}S^0 \wedge_R M \xrightarrow{y_i \wedge \text{id}} \mathbb{F}S^{-n_i} \wedge_R M$$

represents y_i . Smashing over R with $\mathbb{F}S^{-q_{i-1}}$, we obtain a sequence of R -maps

$$(9.11) \quad \mathbb{F}S^{-q_{i-1}} \wedge_R M \longrightarrow \mathbb{F}S^{-q_i} \wedge_R M.$$

Definition 9.12. Define the localization of M at Y , denoted $M[Y^{-1}]$, to be the telescope of the sequence of maps (9.11). Since $M \cong \mathbb{F}S^0 \wedge_R M$ in \mathcal{D}_R , we may regard the inclusion of the initial stage $\mathbb{F}S^0 \wedge_R M$ of the telescope as a natural map $\lambda : M \longrightarrow M[Y^{-1}]$.

Since homotopy groups commute with localization, we see immediately that λ induces an isomorphism of R_* -modules

$$(9.13) \quad \pi_*(M[Y^{-1}]) \cong \pi_*(M)[Y^{-1}].$$

As in Lemma 9.9, the localization of M is the smash product of M with the localization of R .

Lemma 9.14. For a multiplicatively closed set Y of elements of R_* , there is a natural equivalence

$$R[Y^{-1}] \wedge_R M \longrightarrow M[Y^{-1}].$$

Moreover, $R[Y^{-1}]$ is independent of the ordering of the elements of Y . For sets X and Y , $R[(X \cup Y)^{-1}]$ is equivalent to the composite localization $R[X^{-1}][Y^{-1}]$.

10. ALGEBRA STRUCTURES ON LOCALIZATIONS AND ON QUOTIENTS BY IDEALS

The behavior of localizations with respect to algebra structures is immediate.

Proposition 10.1. Let Y be a multiplicatively closed set of elements of R_* . If A is an R -algebra, then $A[Y^{-1}]$ is an R -algebra such that $\lambda : A \longrightarrow A[Y^{-1}]$ is a map of R -algebras. If A is associative or commutative, then so is $A[Y^{-1}]$.

Proof. By Lemmas 9.2 and 9.14, it suffices to observe that $R[Y^{-1}]$ is an associative and commutative R -algebra with unit λ and product the equivalence

$$R[Y^{-1}] \wedge_R R[Y^{-1}] \simeq R[Y^{-1}][Y^{-1}] \simeq R[Y^{-1}]. \quad \square$$

This doesn't work for quotients since $(R/X)/X$ is not equivalent to R/X . However, we can analyze the problem by analyzing the deviation, and, by Lemma 9.9, we may as well work one element at a time. We have a necessary condition for R/x to be an R -algebra that is familiar from classical stable homotopy theory.

Lemma 10.2. Let A be an R -algebra. If A/xA admits an R -algebra structure such that $\iota : A \longrightarrow A/xA$ is a map of R -algebras, then $x : A/xA \longrightarrow A/xA$ is null homotopic as a map of R -modules.

Thus, for example, the Moore spectrum $S/2$ is not an S -algebra since the map $2 : S/2 \longrightarrow S/2$ is not null homotopic. To give a criterion for when R/x does have an R -algebra structure, we first note an easy formal lemma.

Lemma 10.3. Let $\iota : R \longrightarrow M$ be any map of R -modules. Then

$$(\iota \wedge \text{id}) \circ \iota \simeq (\text{id} \wedge \iota) \circ \iota : R \longrightarrow M \wedge_R M.$$

Theorem 10.4. Let $x \in R_m$ and assume that $\pi_{m+1}(R/x) = 0$ and $\pi_{2m+1}(R/x) = 0$. Then R/x admits a structure of R -algebra with unit $\iota : R \rightarrow R/x$. Therefore A/XA admits a structure of R -algebra such that $\iota : A \rightarrow A/XA$ is a map of R -algebras for every R -algebra A and every sequence X of elements of R_* such that $\pi_{m+1}(R/x) = 0$ and $\pi_{2m+1}(R/x) = 0$ if $x \in X$ has degree m .

Proof. Consider the following diagram in the derived category \mathcal{D}_R :

(10.5)

$$\begin{array}{ccccc}
 & & \Sigma^{2m+1}R & & \\
 & & \downarrow x & & \\
 & & \Sigma^{m+1}R & \xrightarrow{x} & \Sigma R \\
 & \swarrow \nu & \downarrow \iota & & \downarrow \iota \\
 \Sigma^m(R/x) & \xrightarrow{x} & R/x & \xrightarrow[\phi]{\iota \wedge \text{id}} & (R/x) \wedge_R (R/x) & \xrightarrow[\sigma]{\pi} & \Sigma^{m+1}(R/x) & \xrightarrow{x} & \Sigma(R/x) \\
 & & & & \downarrow \pi' & & \\
 & & & & \Sigma^{2m+2}R & &
 \end{array}$$

The map x is that specified by (9.4). The bottom row is the cofibre sequence that results from the equivalence

$$(R/x) \wedge_R (R/x) \simeq (R/x)/x$$

of Lemma 9.9, and the column is also a cofibre sequence. The composite $x \circ \iota$ is null homotopic since $\iota \circ x$ is null homotopic and the square commutes. Therefore there is a map ν such that $\pi \circ \nu = \iota$, and ν is unique since $\pi_{m+1}(R/x) = 0$. Since $\pi \circ \nu \circ x = \iota \circ x = 0$, $\nu \circ x$ factors through a map $\Sigma^{2m+1}R \rightarrow R/x$. Since $\pi_{2m+1}(R/x) = 0$, such maps are null homotopic. Thus $\nu \circ x$ is null homotopic. Therefore there is a map σ such that $\sigma \circ \iota = \nu$. Now $\pi \circ \sigma \circ \iota = \pi \circ \nu = \iota$, hence $(\pi \circ \sigma - \text{id})\iota = 0$. Therefore $\pi \circ \sigma - \text{id}$ factors through a map $\Sigma^{2m+2}R \rightarrow \Sigma^{m+1}(R/x)$. Again, such maps are null homotopic. Therefore $\pi \circ \sigma = \text{id}$. Thus the bottom cofibre sequence splits (proving in passing that $x : \Sigma^n(R/x) \rightarrow R/x$ is null homotopic, as it must be). A choice ϕ of a splitting gives a product on R/x . The unit condition $\phi \circ (\iota \wedge \text{id}) = \text{id}$ is automatic. To see that $\phi \circ (\text{id} \wedge \iota) = \text{id}$, we observe that, by the lemma,

$$(\phi \circ (\text{id} \wedge \iota) - \text{id}) \circ \iota = \phi \circ (\text{id} \wedge \iota - \iota \wedge \text{id}) \circ \iota = 0.$$

Therefore $\phi \circ (\text{id} \wedge \iota) - \text{id}$ factors through a map $\Sigma^{m+1}R \rightarrow R/x$. Again, such maps are null homotopic, hence $\phi \circ (\text{id} \wedge \iota) = \text{id}$. This completes the proof that R/x is an R -algebra with unit ι . The rest follows from Lemmas 9.9 and 9.2. \square

The product on R/x can be described a little more concretely. The wedge sum

$$(10.6) \quad (\iota \wedge \text{id}) \vee \sigma : (R/x) \vee \Sigma^{m+1}(R/x) \longrightarrow (R/x) \wedge_R (R/x)$$

is an equivalence. The product ϕ restricts to the identity on the first wedge summand and to the trivial map on the second wedge summand. Thus the product is determined by the choice of σ , and two choices of σ differ by a composite

$$(10.7) \quad \Sigma^{m+1}(R/x) \xrightarrow{\pi'} \Sigma^{2m+2}R \longrightarrow (R/x) \wedge_R (R/x).$$

By the splitting (10.6) and the assumption that $\pi_{m+1}(R/x) = 0$, we can view the second map as an element of $\pi_{2m+2}(R/x)$. If x is not a zero divisor, then $\pi'_* = 0$ on homotopy groups and any two products have the same effect on homotopy groups.

For an R -algebra A and an element x as in the theorem, we give $A/xA \simeq (R/x) \wedge_R A$ the product induced by one of our constructed products on R/x and the given product on A . We refer to any such product as a ‘‘canonical’’ product on A/xA . Observe that, by first using the product on A , the product on A/xA can be factored through

$$\phi \wedge_R \text{id} : (R/x) \wedge_R (R/x) \wedge_R A \longrightarrow (R/x) \wedge_R A.$$

This allows us to smash any diagram giving information about the product on R/x with A and so obtain information about the product on A/xA . Obviously any diagram so constructed is a diagram of right A -modules via the product action of A on itself. This smashing with A can kill obstructions. Clearly, a map of A -modules $\Sigma^q A \longrightarrow M$ is determined by its restriction $S^q \longrightarrow M$ along the unit of A regarded as a map of spectra (or S -modules), which is just an element of $\pi_q(M)$. This leads to the following result.

Theorem 10.8. Let $x \in R_m$ and assume that $\pi_{m+1}(R/x) = 0$ and $\pi_{2m+1}(R/x) = 0$. Let A be an R -algebra and assume that $\pi_{2m+2}(A/xA) = 0$. Then there is a unique canonical product on A/xA . If A is commutative, then A/xA is commutative. If A is associative and $\pi_{3m+3}(A/xA) = 0$, then A/xA is associative.

Proof. The second arrow of (10.7) becomes zero after smashing with A since it is then given by an element of $\pi_{2m+2}(A/xA) = 0$. This proves the uniqueness statement. The commutativity statement follows since if ϕ is a canonical product on A/xA , then so is $\phi\tau$. The associativity statement requires consideration of the restriction of the iterated product to the wedge summands of $A/xA \wedge_R A/xA \wedge_R A/xA$. The details are similar to, but simpler than, those in the proof of Theorem 10.4. \square

Iterating and observing that passage to telescopes can kill obstructions, we arrive at the following fundamental conclusion.

Theorem 10.9. Assume that $R_i = 0$ if i is odd. Let X be a sequence of non zero divisors in R_* such that $\pi_*(R/X)$ is concentrated in degrees congruent to zero mod 4. Then R/X has a unique canonical structure of R -algebra, and it is commutative and associative.

11. THE SPECIALIZATION TO MU -MODULES AND ALGEBRAS

The classical Thom spectra arise in nature as E_∞ ring spectra. In fact, it was inspection of their prespectrum level definition in terms of Grassmannians that first led to the theory of E_∞ ring spectra [19]. Of course, the homotopy groups of MU are concentrated in even degrees, and every non-zero element is a non zero divisor. Thus the results above have the following immediate corollary.

Theorem 11.1. Let X be a regular sequence in MU_* , let I be the ideal generated by X , and let Y be any sequence in MU_* . Then there is an MU -algebra $(MU/X)[Y^{-1}]$ and a natural map of MU -algebras (the unit map)

$$\eta : MU \longrightarrow (MU/X)[Y^{-1}]$$

such that

$$\eta_* : MU_* \longrightarrow \pi_*((MU/X)[Y^{-1}])$$

realizes the natural homomorphism of MU_* -algebras

$$MU_* \longrightarrow (MU_*/I)[Y^{-1}].$$

If MU_*/I is concentrated in degrees congruent to zero mod 4, then there is a unique canonical product on $(MU/X)[Y^{-1}]$, and this product is commutative and associative.

In comparison with earlier constructions of this sort based on the Baas-Sullivan theory of manifolds with singularities or on Landweber's exact functor theorem (where it applies), we have obtained a simpler proof of a substantially stronger result. We emphasize that an MU -algebra is a much richer structure than just a ring spectrum and that commutativity and associativity in the MU -algebra sense are much more stringent conditions than mere commutativity and associativity of the underlying ring spectrum.

We illustrate by explaining how BP appears in this context. Fix a prime p and write $(?)_p$ for localization at p . Let BP be the Brown-Peterson spectrum at p . We are thinking of Quillen's idempotent construction [24], and we have the splitting maps $i : BP \longrightarrow MU_p$ and $e : MU_p \longrightarrow BP$. These are maps of commutative and associative ring spectra such that $e \circ i = \text{id}$. Let I be the kernel of the composite

$$MU_* \longrightarrow MU_{p*} \longrightarrow BP_*.$$

Then I is generated by a regular sequence X , and our MU/X is a canonical integral version of BP . For the moment, let $BP' = (MU/X)_p$. Let $\xi : BP \rightarrow BP'$ be the composite

$$BP \xrightarrow{i} MU_p \xrightarrow{\zeta_p} BP'.$$

It is immediate that ξ is an equivalence. In effect, since we have arranged that ζ_p has the same effect on homotopy groups as e , ξ induces the identity map of $(MU_*/I)_p$ on homotopy groups. By the splitting of MU_p and the fact that self-maps of MU_p are determined by their effect on homotopy groups [2, II.9.3], maps $MU_p \rightarrow BP$ are determined by their effect on homotopy groups. This implies that $\xi \circ e = \zeta_p : MU_p \rightarrow BP'$. The product on BP is the composite

$$BP \wedge BP \xrightarrow{i \wedge i} MU_p \wedge MU_p \xrightarrow{\phi} MU_p \xrightarrow{e} BP.$$

Since ζ_p is a map of MU -algebras and thus of ring spectra, a trivial diagram chase now shows that the equivalence $\xi : BP \rightarrow BP'$ is a map of ring spectra.

We conclude that our BP' is a model for BP that is an MU -algebra, commutative and associative if $p > 2$. The situation for $p = 2$ is interesting. We conclude from the equivalence that BP' is commutative and associative as a ring spectrum, although we do not know that it is commutative or associative as an MU -algebra.

Recall that $\pi_*(BP) = \mathbb{Z}_{(p)}[v_i | \deg(v_i) = 2(p^i - 1)]$, where the generators v_i come from $\pi_*(MU)$ (provided that we use the Hazewinkel generators). We list a few of the spectra derived from BP , with their coefficient rings. Let \mathbb{F}_p denote the field with p elements.

$$\begin{array}{ll} BP\langle n \rangle & \mathbb{Z}_{(p)}[v_1, \dots, v_n] \\ P(n) & \mathbb{F}_p[v_n, v_{n+1}, \dots] \\ k(n) & \mathbb{F}_p[v_n] \end{array} \quad \begin{array}{ll} E(n) & \mathbb{Z}_{(p)}[v_1, \dots, v_n, v_n^{-1}] \\ B(n) & \mathbb{F}_p[v_n^{-1}, v_n, v_{n+1}, \dots] \\ K(n) & \mathbb{F}_p[v_n, v_n^{-1}] \end{array}$$

By the method just illustrated, we can construct canonical integral versions of the $BP\langle n \rangle$ and $E(n)$. All of these spectra fit into the context of Theorem 10.1. If $p > 2$, they all have unique canonical commutative and associative MU -algebra structures. Further study is needed when $p = 2$. In any case, this theory makes it unnecessary to appeal to Baas-Sullivan theory or to Landweber's exact functor theorem for the construction and analysis of spectra such as these.

With more sophisticated techniques, the second author [14] has proven that BP can be constructed as an E_∞ ring spectrum, and in fact admits uncountably many distinct E_∞ ring structures. There is much other ongoing work on the construction and application of new A_∞ and E_∞ ring spectra, by Hopkins, Miller, McClure, and others. The enriched multiplicative structures on rings and modules that we have

discussed are rapidly becoming a standard tool in the study of periodicity phenomena in stable homotopy theory.

REFERENCES

1. J.F. Adams. Lectures on generalised cohomology. Springer Lecture Notes in Mathematics Vol. 99, 1969, 1-138.
2. J.F. Adams. Stable homotopy and generalized homology. University of Chicago. 1974, 1994.
3. J.F. Adams. A variant of E.H.Brown's representability theorem. *Topology*, **10** (1971), 185-198.
4. J.M. Boardman. Stable homotopy theory. Thesis, Warwick 1964; mimeographed notes from Warwick and Johns Hopkins Universities, 1965-1970.
5. J.M. Boardman. Conditionally convergent spectral sequences. Preprint. 1981.
6. E.H. Brown, Jr. Abstract homotopy theory. *Trans. Amer. Math. Soc.* **119** (1965), 79-85.
7. R.R. Bruner, J.P. May, J.E. McClure, and M. Steinberger. H_∞ ring spectra and their applications. Springer Lecture Notes in Mathematics Vol. 1176. 1986.
8. W.G. Dwyer and J. Spalinski. Homotopy theories and model categories. This volume.
9. S. Eilenberg and J.C. Moore. Homology and fibrations, I. *Comm. Math. Helv.* **40** (1966), 199-236.
10. A.D. Elmendorf, J.P.C. Greenlees, I. Kriz, and J.P. May. Commutative algebra in stable homotopy theory and a completion theorem. *Mathematical Research Letters* **1** (1994), 225-239.
11. A.D. Elmendorf, I. Kriz, and J.P. May. Rings, modules, and algebras in stable homotopy theory. In preparation.
12. J.P.C. Greenlees and J.P. May. Generalized Tate cohomology. *Memoirs Amer. Math. Soc.* To appear.
13. V.K.A.M. Gugenheim and J.P. May. On the theory and applications of differential torsion products. *Memoirs Amer. Math. Soc.* **142**, 1974.
14. I. Kriz. Towers of E_∞ ring spectra with an application to BP . Preprint. 1993.
15. I. Kriz, and J.P. May. Operads, algebras, modules, and motives. *Astérisque*. To appear.
16. L.G. Lewis, Jr., J.P. May, and M. Steinberger (with contributions by J.E. McClure). Equivariant stable homotopy theory. Springer Lecture Notes in Mathematics Vol. 1213. 1986.
17. S. Mac Lane. *Categories for the Working Mathematician*. Springer-Verlag. 1971.
18. J.P. May. The Geometry of Iterated Loop Spaces. Springer Lecture Notes in Mathematics Vol. 271. 1972.
19. J.P. May (with contributions by N. Ray, F. Quinn and J. Tornehave). E_∞ ring spaces and E_∞ ring spectra. Springer Lecture Notes in Mathematics Vol. 577. 1977.
20. J.P. May. Multiplicative infinite loop space theory. *J. Pure and Applied Algebra*, **26** (1982), 1-69.
21. J.P. May. Derived categories in algebra and topology. In the Proceedings of the Eleventh International Conference on Topology, *Rendiconti dell Istituto Matematico dell Università di Trieste*. To appear.
22. J.P. May. Equivariant homotopy and cohomology theory. *CBMS Lectures*. In preparation.
23. D.G. Quillen. Homotopical algebra. Springer Lecture Notes in Mathematics Vol. 43. 1967.
24. D.G. Quillen. On the formal group laws of unoriented and complex cobordism theory. *Bull. Amer. Math. Soc.* **75** (1969), 1293-1298.
25. A. Robinson. Derived tensor products in stable homotopy theory. *Topology* **22** (1983), 1-18.
26. A. Robinson. Spectra of derived module homomorphisms. *Math. Proc. Camb. Phil. Soc.* **101** (1987), 249-257.
27. A. Robinson. The extraordinary derived category. *Math. Z.* **196**(1987), 231-238.

28. J.L. Verdier. *Catégories dérivées*. Springer Lecture Notes in Mathematics Vol. 569, 1977, 262-311.
29. G.W. Whitehead. *Elements of homotopy theory*. Springer-Verlag. 1978.

PURDUE UNIVERSITY CALUMET, HAMMOND, IN 46323 USA
E-mail address: aelmendo@math.purdue.edu

THE UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109-1003 USA
E-mail address: ikriz@math.lsa.umich.edu

THE UNIVERSITY OF CHICAGO, CHICAGO, IL 60637 USA
E-mail address: may@math.uchicago.edu