

TATE COHOMOLOGY AND PERIODIC LOCALIZATION OF POLYNOMIAL FUNCTORS

NICHOLAS J. KUHN

ABSTRACT. In this paper, we show that Goodwillie calculus, as applied to functors from stable homotopy to itself, interacts in striking ways with chromatic aspects of the stable category.

Localized at a fixed prime p , let $T(n)$ be the telescope of a v_n self map of a finite S -module of type n . The Periodicity Theorem of Hopkins and Smith implies that the Bousfield localization functor associated to $T(n)_*$ is independent of choices.

Goodwillie's general theory says that to any homotopy functor F from S -modules to S -modules, there is an associated tower under F , $\{P_d F\}$, such that $F \rightarrow P_d F$ is the universal arrow to a d -excisive functor.

Our first main theorem says that $P_d F \rightarrow P_{d-1} F$ always admits a homotopy section after localization with respect to $T(n)_*$ (and so also after localization with respect to Morava K -theory $K(n)_*$). Thus, after periodic localization, polynomial functors split as the product of their homogeneous factors.

This theorem follows from our second main theorem which is equivalent to the following: for any finite group G , the Tate spectrum $t_G(T(n))$ is weakly contractible. This strengthens and extends previous theorems of Greenlees–Sadofsky, Hovey–Sadofsky, and Mahowald–Shick. The Periodicity Theorem is used in an essential way in our proof.

The connection between the two theorems is via a reformulation of a result of McCarthy on dual calculus.

1. INTRODUCTION AND MAIN RESULTS

Over the past twenty years, beginning with the Nilpotence and Periodicity Theorems of E. Devinatz, M. Hopkins, and J. Smith [DHS, HopSm, R2], there has been a steady deepening of our understanding of stable homotopy as organized by the chromatic, or periodic, point of view. During this same period, there have been many new results in homotopical algebra, many following the conceptual model offered by T. Goodwillie's calculus of functors [G1, G2, G3].

Date: July, 2003.

2000 Mathematics Subject Classification. Primary 55P65; Secondary 55N22, 55P60, 55P91.

This research was partially supported by a grant from the National Science Foundation.

Here, and in a previous paper [Ku2], I prove theorems illustrating a beautiful interaction between these two strands of homotopy theory. These results say that certain homotopy functors, stratified via Goodwillie calculus, decompose into their homogeneous strata, after periodic localization. The first paper concerned a highly structured splitting of the important functor $\Sigma^\infty\Omega^\infty$. Ignoring the extra structure, one is left with an illustration of the main result here: after Bousfield localization with respect to a periodic homology theory, *all* polynomial endofunctors of stable homotopy split into a product of their homogeneous components.

We now explain our main results in more detail.

The periodic homology theories we consider are $K(n)_*$, the n^{th} Morava K -theory at a fixed prime p and with $n > 0$, and the ‘telescopic’ variants $T(n)_*$, where $T(n)$ denotes the telescope of a v_n -self map of a finite complex of type n . A consequence of the Periodicity Theorem is that the associated Bousfield class $\langle T(n) \rangle$ is independent of the choice of both the complex and self map. Also, we recall that $T(n)_*$ -acyclics are $K(n)_*$ -acyclic¹; thus the associated localization functors are related by $L_{K(n)} \simeq L_{K(n)}L_{T(n)}$.

Our use of concepts from Goodwillie calculus and localization theory require that we work within a good model category with homotopy category equivalent to the standard stable homotopy category. Thus we work within the category \mathcal{S} , the category of S -modules of [EKMM].

Goodwillie’s general theory then says that a homotopy functor $F : \mathcal{S} \rightarrow \mathcal{S}$ admits a universal tower of fibrations under F ,

$$\begin{array}{ccc}
 & & \vdots \\
 & & \downarrow \\
 & & P_2F(X) \\
 & \nearrow e_2 & \downarrow p_2 \\
 & & P_1F(X) \\
 & \nearrow e_1 & \downarrow p_1 \\
 F(X) & \xrightarrow{e_0} & P_0F(X),
 \end{array}$$

such that

- (1) P_dF is d -excisive, and
- (2) $e_d : F \rightarrow P_dF$ is the universal natural transformation to a d -excisive functor.

¹The Telescope Conjecture asserts the converse, and, these days, is considered unlikely to hold for $n \geq 2$.

Our splitting theorem then is as follows.

Theorem 1.1. *Let $F : \mathcal{S} \rightarrow \mathcal{S}$ be any homotopy functor. For all primes p , $n \geq 1$, and $d \geq 1$, the natural map*

$$p_d(X) : P_d F(X) \rightarrow P_{d-1} F(X)$$

admits a natural homotopy section after applying $L_{T(n)}$.

The theorem can be reformulated as follows. Let $D_d F(X)$ be the fiber of $p_d(X) : P_d F(X) \rightarrow P_{d-1} F(X)$. Then $D_d F$ is both d -excisive and homogeneous: $P_{d-1} D_d F \simeq *$. The theorem is equivalent to the statement that there is a natural weak equivalence of filtered spectra

$$L_{T(n)} P_d F(X) \simeq \prod_{c=0}^d L_{T(n)} D_c F(X).$$

Example 1.2. Here is the simplest example illustrating our theorem. Let $p = 2$. For $k \in \mathbb{Z}$, let $\mathbb{R}P_k^\infty$ be the Thom spectrum of k copies of the canonical line bundle over $\mathbb{R}P^\infty$. [Ku1, Ex.5.7] implies that the cofibration sequence

$$(1.1) \quad \mathbb{R}P_{-1}^\infty \rightarrow \mathbb{R}P_0^\infty \rightarrow S^0$$

splits after $K(n)$ -localization, for all n , even though the connecting map $\delta : S^0 \rightarrow \Sigma \mathbb{R}P_{-1}^\infty$ is nonzero in mod 2 homology.

As was, in essence, observed in a 1983 paper by J.Jones and S.Wegmann [JW], (1.1) is the suspension of the special case $X = S^{-1}$ of a natural cofibration sequence of functors

$$(1.2) \quad (X \wedge X)_{h\mathbb{Z}/2} \rightarrow P_2(X) \rightarrow X.$$

One can also construct this sequence using Goodwillie calculus: see §3.

Theorem 1.1 says that (1.2) splits after applying $L_{T(n)}$ for all n and X , even though the connecting map

$$\delta : X \rightarrow \Sigma(X \wedge X)_{h\mathbb{Z}/2}$$

is often nontrivial before localization.

Remark 1.3. There are various sorts of polynomial functors studied in the literature differing slightly from Goodwillie's d -excisive functors: R.McCarthy has studied d -additive functors [McC], and his student A.Mauer-Oats [MO] has studied an infinite family interpolating between additive and excisive. As will be explained more fully in §6, the analogue of Theorem 1.1 holds in all these generalized settings.

Remark 1.4. Theorem 1.1 and Corollary 1.7 below also has consequences for using the tower $\{P_d F(X)\}$ to understand $E_n^*(F(X))$, where E_n is the usual p -complete integral height n complex oriented commutative S -algebra.

Since it is known [H] that $K(n)_*(X) = 0$ if and only if $E_n^*(X) = 0$, our theorem says that the spectral sequence associated to the tower will collapse at E_1 .

Theorem 1.1 is deduced from a rather different result in equivariant stable homotopy theory that we now describe.

If G is a finite group, let $G\text{-}\mathcal{S}$ denote the category of S -modules with G -action: the category of so-called ‘naive G -spectra’. Note that any S -module can be considered as an object in $G\text{-}\mathcal{S}$ by giving it trivial G -action.

For $Y \in G\text{-}\mathcal{S}$, we let Y_{hG} and Y^{hG} respectively denote associated homotopy orbit and homotopy fixed point S -modules. There are various constructions in the literature, more [GM] or less [ACD, AK, Kl1, WW1] sophisticated, of a natural ‘Norm’ map

$$N(Y) : Y_{hG} \rightarrow Y^{hG}$$

satisfying the key property that $N(Y)$ is an equivalence if Y is a finite free G -CW spectrum. Let the Tate spectrum $t_G(Y)$ be defined as the cofiber of $N(Y)$. As recently observed by J.Klein [Kl2], up to weak equivalence, these constructions are unique: see §2.

We prove the following vanishing theorem.

Theorem 1.5. *For all finite groups G , primes p , and $n \geq 1$,*

$$L_{T(n)}t_G(L_{T(n)}S) \simeq *.$$

This theorem will turn out to be equivalent to the following corollary.

Corollary 1.6. *If $T(n)$ is the telescope of any v_n -self map of a type n complex, then $t_G(T(n)) \simeq *$.*

Besides implying Theorem 1.1, Theorem 1.5 also leads to the following splitting result.

Corollary 1.7. *For any $Y \in G\text{-}\mathcal{S}$, the fundamental cofibration sequence*

$$Y_{hG} \xrightarrow{N(Y)} Y^{hG} \rightarrow t_G(Y)$$

splits after applying $L_{T(n)}$ for any n .

One also immediately deduces results similar to [HSt, Cor.8.7].

Corollary 1.8. *For all finite groups G , the norm map induces an isomorphism*

$$T(n)_*(BG) \xrightarrow{\sim} T(n)^{-*}(BG).$$

Similarly, $L_{T(n)}(\Sigma^\infty BG_+)$ is self dual in the category of $T(n)$ -local spectra.

Our two theorems are supported by three propositions.

The first of these is a slight variant of results of R. McCarthy in [McC], and establishes the connection between our two theorems.

We need to recall Goodwillie's classification of homogeneous polynomial functors [G3]. Let Σ_d denote the d^{th} symmetric group. If our original functor F is *finitary* (terminology from [G3]), i.e. commutes with directed homotopy colimits, then $D_d F(X)$ is weakly equivalent to a homotopy orbit spectrum of the form

$$(C_F(d) \wedge X^{\wedge d})_{h\Sigma_d},$$

where $C_F(d) \in \Sigma_d\text{-}\mathcal{S}$ is determined naturally by F . Important to us is that, even without the finitary hypothesis, there is a natural weak equivalence of the form

$$D_d F(X) \simeq (\Delta_d F(X))_{h\Sigma_d},$$

where $\Delta_d F$ is a functor determined naturally by F , taking values in the category $\Sigma_d\text{-}\mathcal{S}$.

Proposition 1.9. *Let $F : \mathcal{S} \rightarrow \mathcal{S}$ be any homotopy functor. For all $d \geq 1$, there is a homotopy pullback diagram*

$$\begin{array}{ccc} P_d F(X) & \longrightarrow & (\Delta_d F(X))^{h\Sigma_d} \\ \downarrow p_d & & \downarrow \\ P_{d-1} F(X) & \longrightarrow & t_{\Sigma_d}(\Delta_d F(X)). \end{array}$$

This diagram is natural in both X and F .

Our other two propositions together imply Theorem 1.5. The first is a new very general observation about Tate spectra.

Proposition 1.10. *Let R be a ring spectrum and E_* a homology theory. If $t_{\mathbb{Z}/p}(R)$ is E_* -acyclic for all primes p , then so is $t_G(M)$ for all R -modules M and for all finite groups G .*

We remark that, by standard arguments, $t_{\mathbb{Z}/p}(R) \simeq *$, and thus is certainly E_* -acyclic, for all primes p such that R_* is uniquely p -divisible. In particular, to apply the proposition to the pair $(R, E_*) = (L_{T(n)}\mathcal{S}, T(n)_*)$, one need to only look at the single prime involved in the periodic theory.

It is in proving our last proposition that deep results in periodic stable homotopy will be used.

Proposition 1.11. *For all primes p and $n \geq 1$, $L_{T(n)}t_{\mathbb{Z}/p}(L_{T(n)}\mathcal{S}) \simeq *$.*

At this point we need to comment on results like Theorem 1.5 in the literature.

The main theorem of the 1988 article by M.Mahowald and P.Shick [MS] can be restated as

$$(1.3) \quad t_{\mathbb{Z}/2}(T(n)) \simeq *.$$

A proof along their lines can presumably be done at odd primes as well. We will see that the generalization of their theorem to all primes is equivalent to Proposition 1.11, yielding one possible proof of that result. We will offer a rather different proof, using the telescopic functors of Bousfield and the author [B1, Ku1, B2].

The main theorem of the 1996 article by J.Greenlees and H.Sadofsky [GS] reads

$$(1.4) \quad t_G(K(n)) \simeq *.$$

Their proof is elementary (in the sense that consequences of the Nilpotence Theorem are not needed), but heavily uses two special facts about $K(n)$: it is complex oriented, and $K(n)_*(B\mathbb{Z}/p)$ is a finitely generated $K(n)_*$ -module. Note that neither of these two facts is available when considering $T(n)_*$. For readers interested in the simplest proof of (1.4), it is hard to imagine improving upon the clever argument given in [GS, Lemma 2.1] showing that $t_{\mathbb{Z}/p}(K(n)) \simeq *$, but our Proposition 1.10 offers an alternative way to proceed starting from this.

The most substantial part of the main theorem of [HSa] says that

$$(1.5) \quad L_{K(n)}t_G(L_{K(n)}S) \simeq *.$$

Note that, were the Telescope Conjecture true, then (1.5) and Theorem 1.5 would be equivalent; at any rate, the latter implies the former. The authors prove their theorem by starting from (1.4), and then using the Periodicity Theorem, together with the technical heart of Hopkins and D. Ravenel's proof [R2] that $L_{E(n)}$ is a smashing localization. Our proof of Theorem 1.5 bypasses the need for the Hopkins–Ravenel argument.

The rest of the paper is organized as follows. In §2, we review properties of the norm map and t_G , leading to a proof of Proposition 1.10. In §3, supported by the appendix, we first discuss models for $L_E t_{\mathbb{Z}/p}(L_E S)$ for a general spectrum E , and then use telescopic functors to show that the model is contractible when $E = T(n)$. The results of the previous two sections are combined in §4 yielding a proof of Theorem 1.5. Also in this section is a discussion of the equivalence of Theorem 1.5 and Corollary 1.6, with arguments similar in spirit to ones in [MS, HSa]. In §5, we review what we need to about d -excisive functors, and prove Proposition 1.9 with arguments similar to those in [McC]. In §6, we prove our splitting results, Theorem 1.1 and Corollary 1.7.

As is already evident, if E is an S -module, we let L_E denote Bousfield localization with respect to the associated homology theory E_* . Throughout we also use the following conventions regarding functors taking values

in \mathcal{S} . We write $F \xrightarrow[\sim]{f} G$ if $f(X) : F(X) \rightarrow G(X)$ is a weak equivalence for all X . By a weak natural transformation $f : F \rightarrow G$ we mean a pair of natural transformations of the form $F \xleftarrow[\sim]{g} H \xrightarrow{h} G$ or $F \xrightarrow{h} H \xleftarrow[\sim]{g} G$. Finally, we say that a diagram of weak natural transformations commutes if, after evaluation on any object X , the associated diagram commutes in the stable homotopy category.

Acknowledgements I would like to thank various people who have helped me with this project. Randy McCarthy and Greg Arone have helped me learn about Goodwillie towers. Obviously Randy's paper [McC] has been important to my thinking, and Greg suggested the compelling reformulation of Randy's results given in Proposition 1.9. Neil Strickland alerted me to the fact a conjecture of mine, that (1.5) was true, was already a theorem in the literature, and Hal Sadofsky similarly told me about Mahowald and Shick's theorem (1.3). Our main results have been reported on in various seminars and conferences, e.g. at the A.M.S. meetings in January, 2003, and in Oberwolfach in March, 2003.

2. TATE SPECTRA AND PROPOSITION 1.10

2.1. Homotopy orbit and fixed point spectra. For G a fixed finite group, and $Y \in G\text{-}\mathcal{S}$, the S -modules Y_{hG} and Y^{hG} are defined in the usual way:

$$Y_{hG} = (EG_+ \wedge Y)/G, \quad \text{and} \quad Y^{hG} = (\text{Map}_S(EG_+, Y))^G.$$

Both of these functors take weak equivalences and cofibration sequences in $G\text{-}\mathcal{S}$ to weak equivalences and cofibration sequences in \mathcal{S} . (See [GM, Part I] for these sorts of facts.)

Y_{hG} has an important additional property not shared with Y^{hG} : it commutes with filtered homotopy colimits.

We record the following well known facts, which are fundamental when one considers the behavior of Y_{hG} and Y^{hG} under Bousfield localization.

Lemma 2.1. *If $f : Y \rightarrow Z$ is a map in $G\text{-}\mathcal{S}$ that is an E_* -isomorphism, then $f_{hG} : Y_{hG} \rightarrow Z_{hG}$ is also an E_* -isomorphism.*

Lemma 2.2. *If $Y \in G\text{-}\mathcal{S}$ is E -local, so is Y^{hG} .*

2.2. A characterization of the norm map. A recent paper by Klein [Kl2] exploring axioms for generalized Farrell–Tate cohomology leads to a nice characterization of norm maps, and thus Tate spectra.

Proposition 2.3. *Let $N_G(Y), N'_G(Y) : Y_{hG} \rightarrow Y^{hG}$ be natural transformations such that both $N_G(\Sigma^\infty G_+)$ and $N'_G(\Sigma^\infty G_+)$ are weak equivalences.*

Then there is a unique weak natural equivalence $f(Y) : Y_{hG} \xrightarrow{\sim} Y_{hG}$ such that the diagram

$$\begin{array}{ccc} Y_{hG} & \xrightarrow{N_G(Y)} & Y^{hG} \\ & \searrow f(Y) & \uparrow N'_G(Y) \\ & & Y_{hG} \end{array}$$

commutes. It follows that the cofibers of $N_G(Y)$ and $N'_G(Y)$ are naturally weakly equivalent.

We sketch the proof, using the sorts of arguments in [Kl2].

Call a homotopy functor $H : G\text{-}\mathcal{S} \rightarrow \mathcal{S}$ *homological* if it preserves homotopy pushout squares and filtered homotopy colimits. Then Klein, in the spirit of [WW2], observes that any homotopy functor $F : G\text{-}\mathcal{S} \rightarrow \mathcal{S}$ admits a universal left approximation by a homological functor, i.e. there exists homological functor F^{hom} , and a natural transformation $F^{hom}(Y) \rightarrow F(Y)$ satisfying the expected universal property.

Applying this to the case $F(Y) = Y^{hG}$, and observing that Y_{hG} is homological, shows that there is a unique weak natural transformation $g : Y_{hG} \rightarrow Y^{hG, hom}$ yielding a commutative diagram of weak natural transformations

$$\begin{array}{ccc} Y_{hG} & \xrightarrow{N_G(Y)} & Y^{hG} \\ & \searrow g(Y) & \uparrow \\ & & Y^{hG, hom} \end{array}$$

The right upward map is certainly an equivalence for $Y = \Sigma^\infty G_+$, and, by assumption, so is the top map. Thus g is a weak natural transformation between homological functors that is an equivalence when $Y = \Sigma^\infty G_+$. It follows that g is weak equivalence.

Applying this same argument to N'_G yields the proposition.

2.3. Tate spectra. We refer to any natural transformation N_G as in the last proposition as a norm map. The cofiber of $N_G(Y)$ is the associated Tate spectrum, denoted $t_G(Y)$. Both N_G and t_G are unique in the sense of Proposition 2.3; their existence is shown in the various references cited in the introduction.

It is immediate that t_G preserves weak equivalences and cofibration sequences.

From [GM, Prop. I.3.5], we deduce

Lemma 2.4. *If R is a (homotopy) ring spectrum with trivial G action, and M is an R -module, then $t_G(R)$ is a ring spectrum, and $t_G(M)$ is a $t_G(R)$ -module. Furthermore, $R^{hG} \rightarrow t_G(R)$ is a map of R -algebra spectra.*

Fix $Y \in G\text{-}\mathcal{S}$. For each subgroup H of G , Y can be regarded as being in $H\text{-}\mathcal{S}$ by restriction. From [GM, pp.28–29], one deduces

Lemma 2.5. *The assignment $G/H \mapsto t_H(Y)$ defines a Mackey functor to the stable homotopy category. Furthermore, $Y^{hH} \rightarrow t_H(Y)$ is a map of Mackey functors.*

In §5, we will use the following familiar property of the norm map. In the literature, this explicitly appears, with a short axiomatic proof, as [AK, Prop.2.10].

Lemma 2.6. *If K is a finite free G -CW complex, then for all $Y \in G\text{-}\mathcal{S}$, $t_G(\text{Map}_{\mathcal{S}}(K, Y)) \simeq *$.*

2.4. Proof of Proposition 1.10. Recall that R is a ring spectrum, and we are assuming that $t_{\mathbb{Z}/p}(R)$ is E_* -acyclic. We wish to show that $t_G(M)$ is also E_* -acyclic, for all R -modules M , and for all G .

We first note that we can assume $M = R$. For $t_G(M)$ is a $t_G(R)$ -module, and thus the former will be E_* -acyclic if the latter is.

Next we show that we can reduce to the case when G is a p -group. For each prime p dividing the order of G , let $G_p < G$ be a p -Sylow subgroup. Then we have

Lemma 2.7. *Given $Y \in G\text{-}\mathcal{S}$ and E_* a generalized homology theory, $t_G(Y)$ will be E_* -acyclic if $t_{G_p}(Y)$ is E_* -acyclic for all p dividing the order of G .*

Proof. We recall that the completion of the Burnside ring $A(H)$ is denoted $\widehat{A}(H)$. The assignment $G/H \mapsto Y^{hH}$ is then an $\widehat{A}(H)$ -module Mackey functor in the sense of [MM]. Thus so is $G/H \mapsto t_H(Y)$, and then also $G/H \mapsto E_*(t_H(Y))$. Now [MM, Cor.4] implies the lemma. \square

Having reduced Proposition 1.10 to the case when G is a p -group, and is thus solvable, the next lemma implies the proposition.

Lemma 2.8. *Let K be a normal subgroup of G , $Q = G/K$, R a ring spectrum, and E_* a homology theory. If $t_K(R)$ and $t_Q(R)$ are both E_* -acyclic, so is $t_G(R)$.*

Proof. For $Y \in G\text{-}\mathcal{S}$, consider the composite

$$Y_{hG} \simeq (Y_{hK})_{hQ} \xrightarrow{N_K(Y)_{hQ}} (Y^{hK})_{hQ} \xrightarrow{N_Q(Y^{hK})} (Y^{hK})^{hQ} \simeq Y^{hG}.$$

We will know that this composite can be considered a norm map if we check that each of these maps is an equivalence when $Y = \Sigma^\infty G_+$.

As there is an equivalence of S -modules with K -action

$$\Sigma^\infty G_+ \simeq \bigvee_{gK \in Q} \Sigma^\infty K_+,$$

it follows that $N_K(\Sigma^\infty G_+)$, and thus $N_K(\Sigma^\infty G_+)_{hQ}$, is an equivalence.

As there are equivalences of S -modules with Q -action

$$(\Sigma^\infty G_+)_{hK} \xleftarrow[\sim]{N_K(\Sigma^\infty G_+)} (\Sigma^\infty G_+)_{hK} \xrightarrow[\sim]{} \Sigma^\infty Q_+,$$

it follows that $N_Q((\Sigma^\infty G_+)_{hK})$ is an equivalence.

We conclude from this discussion that if both $N_K(R)_{hQ}$ and $N_Q(R^{hK})$ are E_* -isomorphisms, then $N_G(R)$ will also be an E_* -isomorphism, and thus $t_G(R)$ will be E_* -acyclic.

By assumption, $t_K(R)$ is E_* -acyclic. Thus $N_K(R)$ is an E_* -isomorphism. By Lemma 2.1, $N_K(R)_{hQ}$ is also.

By assumption, $t_Q(R)$ is E_* -acyclic. As $t_Q(R^{hK})$ is a $t_Q(R)$ -module, we conclude that $t_Q(R^{hK})$ is also E_* -acyclic, so that $N_Q(R^{hK})$ is an E_* -isomorphism. \square

3. TELESCOPIC FUNCTORS AND PROPOSITION 1.11

The goal of this section is to prove that $L_{T(n)}t_{\mathbb{Z}/p}L_{T(n)}S \simeq *$. We will prove this by establishing that the localized unit map

$$L_{T(n)}S \rightarrow L_{T(n)}t_{\mathbb{Z}/p}L_{T(n)}S$$

is null.

In outline our argument showing this is as follows. It is well known that $t_{\mathbb{Z}/p}S$ can be written as certain inverse limit of Thom spectra. Starting from this, we will show that the unit map $S \rightarrow t_{\mathbb{Z}/p}S$ factors through an inverse limit of ‘connecting maps’ associated to the Goodwillie tower of the functor $\Sigma^\infty \Omega^\infty$ applied to spheres in negative dimensions. We warn the reader of technical complications: odd primes are less pleasant than $p = 2$, we use the theorems of W.H.Lin and J.Gunawardena establishing the Segal conjecture for \mathbb{Z}/p , and a key homological calculation is deferred to an appendix.

It will follow that the localized unit will factor through the inverse limit of the localized connecting maps. That this inverse limit is null will then be an easy consequence of constructions of Bousfield and the author [B1, Ku1, B2] showing that $L_{T(n)}$ factors through Ω^∞ . These ‘telescopic’ constructions heavily use the Periodicity Theorem of Hopkins and Smith [HopSm], and thus are also heavily dependent on the Nilpotence Theorem of [DHS].

3.1. Models for $L_E t_{\mathbb{Z}/p} L_E Y$ and $L_E t_{\Sigma_p} L_E Y$. If α is an orthogonal real representation of a finite group G , we let $S(\alpha)$ and S^α respectively denote the associated unit sphere and one point compactified sphere. Thus $S(\alpha)$

has an unbased G -action while the G -action on S^α is based, and there is a cofibration sequence of based G -spaces

$$S(\alpha)_+ \rightarrow S^0 \rightarrow S^\alpha.$$

Fix a prime p , and let ρ denote Σ_p acting on $\mathbb{R}^p/\Delta(\mathbb{R})$ in the usual way. The action of $\mathbb{Z}/p < \Sigma_p$ on $S(\rho)$ is free, and one concludes that the infinite join $S(\infty\rho)$ is a model for $E\mathbb{Z}/p$. This quickly leads to the following well known description of $t_{\mathbb{Z}/p}$.

Lemma 3.1. *(Compare with [GM, Thm.16.1].) For $Y \in G\text{-}\mathcal{S}$, there is a natural weak equivalence*

$$t_{\mathbb{Z}/p}Y \simeq \operatorname{holim}_k \Sigma \operatorname{Map}_S(S^{k\rho}, Y)_{h\mathbb{Z}/p}.$$

We need a generalization of this.

Lemma 3.2. *For $Y \in G\text{-}\mathcal{S}$, there is a natural weak equivalence*

$$L_E t_{\mathbb{Z}/p} L_E Y \simeq \operatorname{holim}_k \Sigma L_E(\operatorname{Map}_S(S^{k\rho}, Y)_{h\mathbb{Z}/p}).$$

If $(p-1)!$ acts invertibly on E_ , e.g. if E is p -local, there is a natural weak equivalence*

$$L_E t_{\Sigma_p} L_E Y \simeq \operatorname{holim}_k \Sigma L_E(\operatorname{Map}_S(S^{k\rho}, Y)_{h\Sigma_p}).$$

These equivalences are also natural with respect to the partially ordered set of Bousfield classes $\langle E \rangle$, and there are commutative diagrams

$$\begin{array}{ccc} L_E t_{\Sigma_p} L_E Y & \xrightarrow{\sim} & \operatorname{holim}_k \Sigma L_E(\operatorname{Map}_S(S^{k\rho}, Y)_{h\Sigma_p}) \\ \downarrow & & \downarrow \\ L_E t_{\mathbb{Z}/p} L_E Y & \xrightarrow{\sim} & \operatorname{holim}_k \Sigma L_E(\operatorname{Map}_S(S^{k\rho}, Y)_{h\mathbb{Z}/p}). \end{array}$$

Proof. By definition, $L_E t_{\mathbb{Z}/p} L_E Y$ is the cofiber of

$$L_E N_{\mathbb{Z}/p}(L_E Y) : L_E(L_E Y)_{h\mathbb{Z}/p} \rightarrow L_E(L_E Y)^{h\mathbb{Z}/p}.$$

The domain of this map can be simplified:

$$L_E Y_{h\mathbb{Z}/p} \rightarrow L_E(L_E Y)_{h\mathbb{Z}/p}$$

is an equivalence. Meanwhile, the range of this map rewritten via the following chain of natural weak equivalences:

$$\begin{aligned}
L_E(L_E Y)^{h\mathbb{Z}/p} &\xleftarrow{\sim} (L_E Y)^{h\mathbb{Z}/p} \\
&\xrightarrow{\sim} \text{Map}_S(S(\infty\rho)_+, L_E Y)^{h\mathbb{Z}/p} \\
&\xrightarrow{\sim} \text{holim}_k \text{Map}_S(S(k\rho)_+, L_E Y)^{h\mathbb{Z}/p} \\
&\xrightarrow{\sim} \text{holim}_k L_E \text{Map}_S(S(k\rho)_+, L_E Y)^{h\mathbb{Z}/p} \\
&\xleftarrow{\sim} \text{holim}_k L_E \text{Map}_S(S(k\rho)_+, L_E Y)_{h\mathbb{Z}/p} \\
&\xleftarrow{\sim} \text{holim}_k L_E \text{Map}_S(S(k\rho)_+, Y)_{h\mathbb{Z}/p}.
\end{aligned}$$

The crucial second to last equivalence here is induced by norm maps which are equivalences since \mathbb{Z}/p acts freely on $S(k\rho)$.

Thus $L_E t_{\mathbb{Z}/p} L_E Y$ has been identified:

$$\begin{aligned}
L_E t_{\mathbb{Z}/p} L_E Y &\xrightarrow{\sim} \text{holim}_k \text{cofiber} \{L_E Y_{h\mathbb{Z}/p} \rightarrow L_E \text{Map}_S(S(k\rho)_+, Y)_{h\mathbb{Z}/p}\} \\
&\xrightarrow{\sim} \text{holim}_k \Sigma L_E \text{Map}_S(S^{k\rho}, Y)_{h\mathbb{Z}/p}.
\end{aligned}$$

The proof of the statements for t_{Σ_p} are similar, noting that, under the hypothesis that $(p-1)!$ acts invertibly on E_* , the norm maps

$$\text{Map}_S(S(k\rho)_+, L_E Y)_{h\Sigma_p} \rightarrow \text{Map}_S(S(k\rho)_+, L_E Y)^{h\Sigma_p}$$

will still be equivalences. \square

For $r \geq 0$, and X an S -module, we let $D_r X = (X^{\wedge r})_{h\Sigma_r}$, and we recall that there are natural transformations $\Sigma D_r X \rightarrow D_r \Sigma X$. Specializing to $r = p$, a quick check of definitions verifies the next lemma.

Lemma 3.3. *There is a natural weak equivalence*

$$\Sigma^k D_p \Sigma^{-k} X \simeq \text{Map}_S(S^{k\rho}, X^{\wedge p})_{h\Sigma_p},$$

and thus there is a p -local equivalence

$$t_{\Sigma_p} S \simeq \text{holim}_k \Sigma^{k+1} D_p S^{-k}.$$

Define $d_k : S \rightarrow \Sigma^{k+1} D_p S^{-k}$ to be the composite

$$S \xrightarrow{\text{unit}} t_{\Sigma_p} S \longrightarrow \Sigma^{k+1} D_p S^{-k}.$$

As the restriction map $t_{\Sigma_p} S \rightarrow t_{\mathbb{Z}/p} S$ is unital, our various observations combine to yield the following proposition.

Proposition 3.4. *If E is p -local, $L_E t_{\mathbb{Z}/p} L_E S \simeq *$ if and only if*

$$\text{holim}_k L_E d_k : L_E S \rightarrow \text{holim}_k L_E \Sigma^{k+1} D_p S^{-k}$$

is null.

3.2. The Goodwillie tower of $\Sigma^\infty\Omega^\infty$. Recall that $\Sigma^\infty Z$ denotes the suspension spectrum of a space Z , and that Σ^∞ has right adjoint Ω^∞ , where $\Omega^\infty X$ is the zeroth space of a spectrum X .

Let $P_r(X)$ denote the r^{th} functor in the Goodwillie tower of the functor

$$\Sigma^\infty\Omega^\infty : \mathcal{S} \rightarrow \mathcal{S}.$$

Thus this Goodwillie tower has the form

$$\begin{array}{c}
 \vdots \\
 \downarrow \\
 P_3(X) \\
 \downarrow p_3(X) \\
 P_2(X) \\
 \downarrow p_2(X) \\
 P_1(X).
 \end{array}
 \begin{array}{l}
 \\
 \nearrow e_3(X) \\
 \nearrow e_2(X) \\
 \xrightarrow{e_1(X)}
 \end{array}
 \Sigma^\infty\Omega^\infty X$$

This tower has the following fundamental properties.

- (1) If X is 0-connected, then $\Sigma^\infty\Omega^\infty X \rightarrow \text{holim}_r P_r(X)$ is an equivalence.
- (2) The fiber of $p_r(X) : P_r(X) \rightarrow P_{r-1}(X)$ is naturally weakly equivalent to $D_r(X)$.
- (3) There are equivalences $D_1 X \simeq P_1 X \simeq X$, and via the second of these, $e_1(X) : \Sigma^\infty\Omega^\infty X \rightarrow P_1 X$ can be identified with evaluation map $\epsilon(X) : \Sigma^\infty\Omega^\infty X \rightarrow X$.

All of these properties can be deduced from Goodwillie's general theory. For an explicit discussion of these (and more) see [AK] or [Ku2].

3.3. Telescopic functors. Bousfield and the author have deduced the the following consequence of the Periodicity Theorem.

Theorem 3.5. *There exists a functor $\Phi_n : \text{Spaces} \rightarrow S\text{-modules}$ and a natural weak equivalence $\Phi_n\Omega^\infty X \simeq L_{T(n)}X$.*

With the result stated at the level of homotopy categories, and with $K(n)$ replacing $T(n)$, this is the main theorem of [Ku1]. However the sorts of constructions given there, and in [B1] (for $n = 1$), yield the theorem as stated: see [B2].

This has the following immediate corollary [Ku1, Ku2].

Corollary 3.6. *There is a natural factorization by weak S -module maps*

$$\begin{array}{ccc} & L_{T(n)}\Sigma^\infty\Omega^\infty X & \\ \eta_n(X) \nearrow & & \searrow L_{T(n)}\epsilon(X) \\ L_{T(n)}X & \xlongequal{\quad\quad\quad} & L_{T(n)}X. \end{array}$$

To use this, we recall an observation about *reduced* homotopy functors, functors $F : \mathcal{S} \rightarrow \mathcal{S}$ such that $F(X)$ is contractible whenever X is. Goodwillie observes that then there is an induced weak natural transformation

$$\Sigma F(X) \longrightarrow F(\Sigma X).$$

The naturality is with respect to both X and F . For example, if $F = D_r$, this natural transformation agrees with the one discussed previously.

In particular, we can apply this construction to both the domain and range of the natural transformation

$$L_{T(n)}P_p \rightarrow L_{T(n)}P_1,$$

evaluated on $\Sigma^{-k}X$ for all $k \geq 0$. Recalling that $L_{T(n)}$ commutes with suspension and $P_1(X) \simeq X$, we obtain maps

$$\operatorname{holim}_k \Sigma^k L_{T(n)}P_p(\Sigma^{-k}X) \rightarrow L_{T(n)}X.$$

Theorem 3.7. $\operatorname{holim}_k \Sigma^k L_{T(n)}P_p(\Sigma^{-k}X) \rightarrow L_{T(n)}X$ admits a homotopy section.

Proof. A section is given by $\operatorname{holim}_k \Sigma^k (L_{T(n)}e_p(\Sigma^{-k}X) \circ \eta_n(\Sigma^{-k}X))$. \square

3.4. Specialization to odd spheres. Standard homology calculations as in [CLM, BMMS] imply the next lemma.

Lemma 3.8. *Localized at an odd prime p , $D_r S^k \simeq *$ for odd $k \in \mathbb{Z}$, and for $2 \leq r \leq p-1$. Thus (for all primes p) the natural map*

$$\operatorname{holim}_k \Sigma^k P_{p-1}(S^{-k}) \rightarrow S$$

is a p -local equivalence.

Continuing the cofibration sequence $D_p X \rightarrow P_p(X) \rightarrow P_{p-1}(X)$ one step to the right defines a natural transformation

$$\delta(X) : P_{p-1}(X) \rightarrow \Sigma D_p X.$$

Localized at p , define $\delta_k : S \rightarrow \Sigma^{k+1} D_p S^{-k}$ to be the composite

$$S \xleftarrow{\sim} \Sigma^k P_{p-1}(S^{-k}) \xrightarrow{\Sigma^k \delta(S^{-k})} \Sigma^{k+1} D_p S^{-k}.$$

Proposition 3.9. $\operatorname{holim}_k L_{T(n)}\delta_k : L_{T(n)}S \rightarrow \operatorname{holim}_k L_{T(n)}\Sigma^{k+1} D_p S^{-k}$ is null.

Proof. Localized at p , there is a cofibration sequence

$$\operatorname{holim}_k L_{T(n)} \Sigma^k P_p S^{-k} \rightarrow L_{T(n)} S \rightarrow \operatorname{holim}_k L_{T(n)} \Sigma^{k+1} D_p S^{-k}.$$

Theorem 3.7 says that the first map has a section. Thus the second map is null. \square

3.5. Proof of Proposition 1.11. A comparison of Proposition 3.4 with Proposition 3.9 shows that we will have proved Proposition 1.11 once we check the following lemma.

Lemma 3.10. $\operatorname{holim}_k d_k : S \rightarrow \operatorname{holim}_k \Sigma^{k+1} D_p S^{-k}$ *factors through*

$$\operatorname{holim}_k \delta_k : S \rightarrow \operatorname{holim}_k \Sigma^{k+1} D_p S^{-k}.$$

Proof. W.H.Lin's theorem [L], when $p = 2$, and J.Gunawardena's theorem [Gun, AGM], when p is odd, can be stated in the following way:

$$\operatorname{holim}_k d_k : S \rightarrow \operatorname{holim}_k \Sigma^{k+1} D_p S^{-k}$$

is p -adic completion. It follows that we need to check that $\operatorname{holim}_k \delta_k \in \pi_0(\operatorname{holim}_k \Sigma^{k+1} D_p S^{-k}) \simeq \mathbb{Z}_p$ is a topological generator. As topological generators of \mathbb{Z}_p are detected mod p , the next lemma, whose proof is deferred to the appendix, completes our argument. \square

Lemma 3.11. $\delta(S^{-1}) : P_{p-1}(S^{-1}) \rightarrow \Sigma D_p S^{-1}$ *is nonzero in mod p homology.*

4. THE PROOFS OF THEOREM 1.5 AND COROLLARY 1.6

We begin this section by noting how Proposition 1.11 and Proposition 1.10 together imply Theorem 1.5. Proposition 1.11 can be restated as saying that $t_{\mathbb{Z}/p} L_{T(n)} S$ is $T(n)_*$ -acyclic. Recalling that the localization of a ring spectrum (e.g. S) is again a ring spectrum, Proposition 1.10 can then be applied to the pair $(R, E_*) = (L_{T(n)} S, T(n)_*)$, to conclude that $t_G L_{T(n)} S$ is $T(n)_*$ -acyclic for all G . This is a restatement of Theorem 1.5.

Now we turn to showing how Corollary 1.6 can be deduced from Theorem 1.5, and vice versa.

We need to review some of the fine points of the Periodicity Theorem. (A good reference for this is [R2].) We fix a prime p , and work with p -local spectra. A finite spectrum F is of type n if $K(n)_*(F) \neq 0$, but $K(i)_*(F) = 0$ for $i < n$. Let $\mathcal{C}_n = \{\text{finite } F \mid F \text{ has type at least } n\}$. Then every $F \in \mathcal{C}_n$ admits a v_n self map: a map $f : \Sigma^d F \rightarrow F$ such that $K(n)_*(f)$ is an isomorphism, but $K(i)_*(f) = 0$ for all $i \neq n$. If $n > 0$, then d will necessarily be positive. In all cases, f is unique and natural up to iteration. Thus there is a well defined functor from \mathcal{C}_n to spectra sending F to $v_n^{-1} F$, the telescope of any v_n self map of F . We note that v_n preserves both cofibration sequences and retracts.

The Thick Subcategory Theorem says that any thick subcategory of the category of p -local spectra, i.e. any collection of p -local finite spectra closed under cofibration sequences and retracts, is \mathcal{C}_n for some $n \geq 0$.

We recall that $L_{T(n)}$ denotes $L_{v_n^{-1}F}$ for any F of type n . From the facts stated above, it is easily verified that this is independent of choice of F , and that for all $F \in \mathcal{C}_n$, $L_{T(n)}(F) = v_n^{-1}F$. Finally we note that if F has type n and F' has type $i \neq n$, then $v_n^{-1}F \wedge v_i^{-1}F' \simeq *$.

Lemma 4.1. *Fix a finite group G . The following conditions are equivalent.*

- (1) $t_G(L_{T(n)}S)$ is $T(n)_*$ -acyclic.
- (2) For all $F \in \mathcal{C}_n$, $t_G(v_n^{-1}F) \simeq *$.
- (3) For all type n complexes F , $t_G(v_n^{-1}F) \simeq *$.
- (4) There exists a type n complex F such that $t_G(v_n^{-1}F) \simeq *$.

Note that statement (1) is the conclusion of Theorem 1.5 and (3) is the conclusion of Corollary 1.6.

Clearly (2) implies (3), which in turn implies (4). To see that (4) implies (2), note that the collection of $F \in \mathcal{C}_n$ such that $t_G(v_n^{-1}F) \simeq *$ forms a thick subcategory contained in \mathcal{C}_n . Such a thick subcategory will equal all of \mathcal{C}_n if it contains *any* type n finite. (This type of reasoning appears in [MS].)

Now suppose (1) holds. Since $v_n^{-1}F \simeq L_{T(n)}F$, it is an $L_{T(n)}S$ -module, and we see that $t_G(v_n^{-1}F)$ is $(v_n^{-1}F)_*$ -acyclic for all finite F of type n . It is easy to find a type n finite F that is a ring spectrum; thus so is $R = v_n^{-1}F$. But then $t_G(R)$ will be an R_* -acyclic R -module, and thus contractible, i.e. statement (4) holds.

It remains to show that (2) implies (1). We reason as in [HSa].

Define finite spectra $F(0), \dots, F(n)$ by first setting $F(0) = S$, and then recursively defining $F(i+1)$ to be the cofiber of a v_i self map of $F(i)$.

Ravenel [R1] observes that if $f : \Sigma^d X \rightarrow X$ is a self map with cofiber C and telescope T , then $\langle X \rangle = \langle C \vee T \rangle$. Applying this n times leads to an equality of Bousfield classes

$$\langle S \rangle = \langle F(n) \vee \bigvee_{i=0}^{n-1} v_i^{-1}F(i) \rangle.$$

Smashing this with $t_G(L_{T(n)}S)$, and noting that

$$t_G(L_{T(n)}S) \wedge F(n) \simeq t_G(L_{T(n)}F(n)) \simeq t_G(v_n^{-1}F(n)),$$

leads to

$$\langle t_G(L_{T(n)}\mathcal{S}) \rangle = \langle t_G(v_n^{-1}F(n)) \vee \bigvee_{i=0}^{n-1} t_G(L_{T(n)}\mathcal{S}) \wedge v_i^{-1}F(i) \rangle.$$

Smashing this with $T(n)$, and noting that $T(n) \wedge v_i^{-1}F(i) \simeq *$ if $i < n$, leads to

$$\langle T(n) \wedge t_G(L_{T(n)}\mathcal{S}) \rangle = \langle T(n) \wedge t_G(v_n^{-1}F(n)) \rangle.$$

If (2) holds, then the right side of this last equation is the Bousfield class of a contactible spectrum. Thus so is the left, i.e. (1) holds.

5. POLYNOMIAL FUNCTORS AND TATE COHOMOLOGY

In this section we sketch a proof of Proposition 1.9. As I hope will be clear, this proposition is just a variant of [McC, Prop.4], and our proof uses precisely the same ideas that McCarthy does.

5.1. Review of Goodwillie calculus. In the series of papers [G1, G2, G3], Tom Goodwillie has developed his theory of polynomial resolutions of homotopy functors. We need to summarize some aspects of Goodwillie's work as they apply to functors from S -modules to S -modules. Throughout we cite the version of [G3] of June, 2002.

In [G2], Goodwillie begins by defining and studying the *total homotopy fiber* of a cubical diagram. For example the total homotopy fiber of a square

$$\begin{array}{ccc} X_0 & \longrightarrow & X_1 \\ \downarrow & & \downarrow \\ X_2 & \longrightarrow & X_{12} \end{array}$$

is the homotopy fiber of the evident map from X_0 to the homotopy pullback of the square with X_0 omitted. A cubical diagram is then *homotopy cartesian* if its total fiber is weakly contractible. Dual constructions similarly define *total homotopy cofibers* and *homotopy cocartesian cubes*. We note that in a stable model category like \mathcal{S} , a cubical diagram is homotopy cartesian exactly when it is homotopy cocartesian.

A cubical diagram is *strongly homotopy cocartesian* if each of its 2 dimensional faces is homotopy cocartesian. A functor is then said to be *d-excise* if it takes strongly homotopy cocartesian $(d+1)$ -cubical diagrams to homotopy cartesian cubical diagrams.

In [G3], given a functor F , Goodwillie proves the existence of a tower $\{P_d F\}$ under F so that $F \rightarrow P_d F$ is the universal arrow to a d -excise functor, up to weak equivalence.

For functors with range in a stable model category, Goodwillie [G3] gives a description of how $D_d F(X)$, the fiber of $P_d F(X) \rightarrow P_{d-1} F(X)$, can be computed by means of cross effects. We describe how this goes in our setting.

Let $F : \mathcal{S} \rightarrow \mathcal{S}$ be a functor. Let $\mathbf{d} = \{1, 2, \dots, d\}$. In [G3, §3], $cr_d F$, the d^{th} cross effect of F , is defined to be the functor of d variables given as the total homotopy fiber

$$(cr_d F)(X_1, \dots, X_d) = \text{TotFib}_{T \subset \mathbf{d}} F\left(\bigvee_{i \in \mathbf{d}-T} X_i\right).$$

A d -variable homotopy functor $H : \mathcal{S}^d \rightarrow \mathcal{S}$ is *reduced* if $H(X_1, \dots, X_d)$ is contractible whenever any of the X_i are. Given such a functor, its *multilinearization* $\mathcal{L}(H) : \mathcal{S}^d \rightarrow \mathcal{S}$ is defined by the formula

$$(5.1) \quad \mathcal{L}(H)(X_1, \dots, X_d) = \text{hocolim}_{n_i \rightarrow \infty} \Omega^{n_1 + \dots + n_d} H(\Sigma^{n_1} X_1, \dots, \Sigma^{n_d} X_d).$$

This will be 1-excisive in each variable.

Now define $\Delta_d F : \mathcal{S} \rightarrow \Sigma_{d-}\mathcal{S}$ by the formula

$$\Delta_d F(X) = \mathcal{L}(cr_d F)(X, \dots, X).$$

Then [G3, Theorems 3.5, 6.1] says that there is a natural weak equivalence

$$(5.2) \quad D_d F(X) \simeq (\Delta_d F)(X)_{h\Sigma_d}.$$

We need to explain some of the ideas behind this formula.

Firstly, $\Delta_d(F) \rightarrow \Delta_d(P_d F)$ is always an equivalence, and it follows that one can assume the original functor F is d -excisive.

If F is d -excisive then $cr_d F$ is already 1-excisive in each variable [G3, Prop.3.3], and so $\Delta_d F(X)$ can be identified with $(cr_d F)(X, \dots, X)$. In this case, the natural map

$$D_d F(X) \rightarrow P_d F(X)$$

identifies with the natural transformation

$$\alpha_d(X) : (\Delta_d F)(X)_{h\Sigma_d} \rightarrow F(X)$$

defined to be the composite

$$(\Delta_d F)(X)_{h\Sigma_d} \rightarrow F\left(\bigvee_{i=1}^d X\right)_{h\Sigma_d} \rightarrow F(X).$$

Here the second map is induced by the fold map $\bigvee_{i=1}^d X \rightarrow X$.

Goodwillie proves (5.2) by verifying that $cr_d(\alpha_d)$ is an equivalence, so that $D_d(\alpha_d)$ is an equivalence. Enroute to this, he shows that there is a natural equivariant weak equivalence

$$cr_d(\Delta_d F) \simeq \Sigma_{d+} \wedge cr_d F.$$

5.2. Dual constructions. In [McC], McCarthy investigates ‘dual calculus’. In this spirit, replacing wedges by products, fibers by cofibers, etc., leads to constructions dual to the above. In particular, given $F : \mathcal{S} \rightarrow \mathcal{S}$, we define $cr^d F : \mathcal{S}^d \rightarrow \mathcal{S}$ by the formula

$$(cr^d F)(X_1, \dots, X_d) = \text{TotCofib}_{T \subset \mathbf{d}} F\left(\prod_{i \in T} X_i\right),$$

and then we define $\Delta^d F : \mathcal{S} \rightarrow \Sigma_d \mathcal{S}$ by

$$\Delta^d F(X) = \mathcal{L}(cr^d F)(X, \dots, X).$$

Because both the domain and range of F is a stable model category, one sees that each of the natural transformations

$$cr_d F \rightarrow cr^d F$$

and

$$\Delta_d F \rightarrow \Delta^d F$$

are weak equivalences.

If F is d -excisive then $\Delta^d F(X)$ can be identified with $(cr^d F)(X, \dots, X)$. In this case, we define the weak natural transformation

$$\alpha^d(X) : F(X) \rightarrow (\Delta_d F)(X)^{h\Sigma_d}$$

to be the zig-zag composite

$$F(X) \rightarrow F(X^d)^{h\Sigma_d} \rightarrow (\Delta^d F)(X)^{h\Sigma_d} \xleftarrow{\sim} (\Delta_d F)(X)^{h\Sigma_d}.$$

Here the first map is induced by the diagonal $X \rightarrow X^d$.

Arguments dual to Goodwillie's show that the next lemma holds.

Lemma 5.1. *(Compare with [McC, Lemmas 3.7,3.8].) Let $F : \mathcal{S} \rightarrow \mathcal{S}$ be d -excisive.*

(1) $cr^d(\alpha^d)$, and thus $D_d(\alpha^d)$, is an equivalence.

(2) There is a natural equivariant weak equivalence

$$cr^d(\Delta^d F) \simeq \text{Map}_{\mathcal{S}}(\Sigma_+, cr^d F).$$

5.3. Proof of Proposition 1.9. Proposition 1.9 is a formal consequence of Lemma 5.1. First of all, we observe the following.

Lemma 5.2. *(Compare with [McC, proof of Prop.4].) Let F be d -excisive. Then $t_{\Sigma_d}(\Delta_d F)$ is $(d-1)$ -excisive. Thus the cofibration sequence*

$$D_d((\Delta_d F)^{h\Sigma_d}) \rightarrow P_d(\Delta_d F)^{h\Sigma_d} \rightarrow P_{d-1}((\Delta_d F)^{h\Sigma_d})$$

identifies with the norm sequence

$$(\Delta_d F)_{h\Sigma_d} \rightarrow (\Delta_d F)^{h\Sigma_d} \rightarrow t_{\Sigma_d}(\Delta_d F).$$

Proof. For the first statement, we check that $cr^d(t_{\Sigma_d}(\Delta_d F)) \simeq *$:

$$cr^d(t_{\Sigma_d}(\Delta_d F)) \simeq t_{\Sigma_d}(cr^d(\Delta_d F)) \simeq t_{\Sigma_d}(\text{Map}_{\mathcal{S}}(\Sigma_+, cr^d F)) \simeq *.$$

Here we have used Lemma 5.1(2) and Lemma 2.6.

As $(\Delta_d F)_{h\Sigma_d}$ is d -excisive and homogeneous, the second statement follows. \square

Now we turn to the proof of Proposition 1.9. We can assume that F is d -excisive. Assuming this, the last lemma implies that the weak natural transformation $\alpha^d(X) : F(X) \rightarrow (\Delta_d F(X))^{h\Sigma_d}$ induces a commutative diagram of weak natural transformations

$$\begin{array}{ccc} D_d F(X) & \longrightarrow & (\Delta_d F(X))_{h\Sigma_d} \\ \downarrow & & \downarrow \\ P_d F(X) & \longrightarrow & (\Delta_d F(X))^{h\Sigma_d} \\ \downarrow & & \downarrow \\ P_{d-1} F(X) & \longrightarrow & t_{\Sigma_d}(\Delta_d F(X)). \end{array}$$

In this diagram each of the vertical columns is a homotopy fibration sequence of S -modules. The top map is a weak equivalence thanks to Lemma 5.1(1). Thus the bottom square is a homotopy pullback diagram.

5.4. Polynomial functor variants. McCarthy and his student Mauer-Oats [MO] have explored various different notions of what it might mean to say a functor $F : \mathcal{A} \rightarrow \mathcal{B}$ is polynomial of degree at most d , with d -excisive and d -additive as two special cases. In these variants \mathcal{B} should surely be a reasonable model category, but \mathcal{A} can often be a category with much less structure. As a hint of why this might be true, note that the definition of cross effects only uses the existence of finite coproducts in \mathcal{A} .

If \mathcal{B} is any stable model category admitting norm maps, and \mathcal{A} is also appropriately stable, then the evident analogue of Proposition 1.9 still holds. The discussion above goes through with one little change: the formula (5.1) for the (multi)linearization process \mathcal{L} needs to be adjusted to reflect the notion of degree 1 functor at hand. Note that our proof of Proposition 1.9 didn't use this formula (nor did McCarthy's arguments in [McC]).

Of relevance to the next section, we note that these variants of \mathcal{L} are still homotopy colimits, and thus preserve E_* -isomorphisms.

6. LOCALIZATION AND THE PROOFS OF THEOREM 1.1 AND COROLLARY 1.7

In this section, we show how our vanishing Tate cohomology result, Theorem 1.5, leads to the splitting results Theorem 1.1 and Corollary 1.7. To simplify notation, we let $L = L_{T(n)}$.

Proof of Corollary 1.7. Let Y be an S -module with G action. We wish to show that the norm sequence

$$Y_{hG} \xrightarrow{N(Y)} Y^{hG} \rightarrow t_G(Y)$$

splits after applying L . Thus we need to construct a left homotopy inverse to $L(N(Y))$.

The localization map $Y \rightarrow LY$ induces a commutative diagram

$$\begin{array}{ccc} Y_{hG} & \xrightarrow{N(Y)} & Y^{hG} \\ \downarrow & & \downarrow \\ (LY)_{hG} & \xrightarrow{N(LY)} & (LY)^{hG}. \end{array}$$

Applying L to this yields the diagram

$$\begin{array}{ccc} L(Y_{hG}) & \xrightarrow{L(N(Y))} & L(Y^{hG}) \\ \downarrow \wr & & \downarrow L(\eta^{hG}) \\ L((LY)_{hG}) & \xrightarrow[\sim]{L(N(LY))} & L((LY)^{hG}). \end{array}$$

Here the left vertical map is an equivalence, as homology isomorphisms are preserved by taking homotopy orbits (Lemma 2.1). The lower map, $L(N(LY))$, is an equivalence by Theorem 1.5: its cofiber, $L(t_G(LY))$, is a module over $L(t_G(LS))$, and is thus contractible.

Our desired left homotopy inverse is now obtained by composing the right vertical map of the diagram with the inverses of the two indicated equivalences. \square

Proof of Theorem 1.1. We are given a functor $F : \mathcal{S} \rightarrow \mathcal{S}$ and wish to prove that

$$D_d F(X) \rightarrow P_d F(X) \rightarrow P_{d-1} F(X)$$

splits after applying L . Thus we need to construct a left homotopy inverse to $LD_d F(X) \rightarrow LP_d F(X)$.

We need a lemma that plays the role that Lemma 2.1 played in the previous proof. Call a natural transformation $F \rightarrow G$ an E_* -isomorphism, if $F(X) \rightarrow G(X)$ is an E_* -isomorphism for all X .

Formula (5.2) says that D_d is the composition of constructions each of which preserve E_* -isomorphisms, and thus we have

Lemma 6.1. *If $F \rightarrow G$ is an E_* -isomorphism, then so is $D_d F \rightarrow D_d G$.*

Remark 6.2. This lemma holds for the variants on the notion of d -excisive, as discussed above in §5.4.

Armed with this lemma, Theorem 1.1 is proved as follows.

The localization natural transformation $F \rightarrow LF$, together with Proposition 1.9, induce a commutative diagram

$$\begin{array}{ccccc} D_d F & \longrightarrow & D_d(LF) & \xrightarrow{\sim} & \Delta_d(LF)_{h\Sigma_d} \\ \downarrow & & \downarrow & & \downarrow \\ P_d F & \longrightarrow & P_d(LF) & \longrightarrow & \Delta_d(LF)^{h\Sigma_d}. \end{array}$$

Applying L to this, gives the diagram

$$\begin{array}{ccccc} LD_d F & \xrightarrow{\sim} & LD_d(LF) & \xrightarrow{\sim} & L(\Delta_d(LF))^{h\Sigma_d} \\ \downarrow & & \downarrow & & \downarrow \wr \\ LP_d F & \longrightarrow & LP_d(LF) & \longrightarrow & L(\Delta_d(LF))^{h\Sigma_d}. \end{array}$$

Here the top left natural transformation is an equivalence by the lemma just stated. The right vertical natural transformation is an equivalence by Theorem 1.5, as its cofiber, $L(t_{\Sigma_d}(\Delta_d(LF)))$, is an $L(t_G(LS))$ -module, when evaluated on any X . (Though not necessarily local, due to the hocolimit construction \mathcal{L} , $\Delta_d(LF)(X)$ is nevertheless an LS -module.)

Our desired left homotopy inverse is now obtained by composing the natural transformation along the bottom of this diagram with the inverses of the three indicated equivalences. \square

APPENDIX A. PROOF OF LEMMA 3.11

We begin with some needed notation.

Recall that P_r denotes the r^{th} Goodwillie approximation to the functor $\Sigma^\infty \Omega^\infty$. We let

$$\delta(X) : P_{r-1}(X) \rightarrow \Sigma D_r X$$

denote the connecting map for the cofibration sequence

$$D_r X \rightarrow P_r(X) \rightarrow P_{r-1}(X).$$

Given any reduced homotopy functor $F : \mathcal{S} \rightarrow \mathcal{S}$, we let

$$\Delta(X) : \Sigma F(X) \rightarrow F(\Sigma X)$$

denote the canonical natural map.

Fixing a prime p , all homology will be with \mathbb{Z}/p coefficients. The Steenrod operations act on $H_*(X)$ as operations lowering dimensions. To unify the ‘even’ prime and odd prime cases, we let $\mathcal{P}^1 = Sq^2$, when $p = 2$. Thus, for all primes p , \mathcal{P}^1 lowers degree by $2p - 2$.

The goal of this appendix is to prove Lemma 3.11, which we restate more precisely.

Lemma A.1. $\delta_* : H_{-1}(P_{p-1}(S^{-1})) \rightarrow H_{-1}(\Sigma D_p(S^{-1}))$ is an isomorphism of one dimensional \mathbb{Z}/p -modules.

Recall that $H_*(D_r X)$ is a known functor of $H_*(X)$, both additively, and as a module over the Steenrod algebra. Furthermore, the behavior of $\Delta_* : H_*(\Sigma D_r X) \rightarrow H_*(D_r \Sigma X)$ is known. See [CLM, BMMS].

Naturality implies that there is a commutative diagram:

$$\begin{array}{ccc}
H_{-1}(P_{p-1}(S^{-1})) & \xrightarrow{\delta_*} & H_{-1}(\Sigma D_p(S^{-1})) \\
\downarrow \wr & & \downarrow \wr \\
H_{-1}(P_{p-1}(\Sigma^{-1}H\mathbb{Z})) & \xrightarrow{\delta_*} & H_{-1}(\Sigma D_p(\Sigma^{-1}H\mathbb{Z})) \\
\uparrow \mathcal{P}_*^1 & & \uparrow \wr \mathcal{P}_*^1 \\
H_{2p-3}(P_{p-1}(\Sigma^{-1}H\mathbb{Z})) & \xrightarrow{\delta_*} & H_{2p-3}(\Sigma D_p(\Sigma^{-1}H\mathbb{Z})) \\
\downarrow \Delta_* & & \downarrow \wr \Delta_* \\
H_{2p-3}(\Sigma^{-2}P_{p-1}(\Sigma H\mathbb{Z})) & \xrightarrow{\Sigma^{-2}\delta_*} & H_{2p-3}(\Sigma^{-1}D_p(\Sigma H\mathbb{Z})),
\end{array}$$

where the top vertical maps are induced by the inclusion $S^{-1} \rightarrow \Sigma^{-1}H\mathbb{Z}$. The top square is a square of homology groups of lowest degree. That the indicated maps are isomorphisms, all between one dimensional vector spaces, is an easy consequence of facts from [CLM, BMMS]. For example, the middle right map is an isomorphism due to the Nishida relation

$$\mathcal{P}_*^1 \beta Q^1 x = \beta Q^0 x \in H_{-2}(D_p(\Sigma^{-1}H\mathbb{Z})),$$

for $x \in H_{-1}(\Sigma^{-1}H\mathbb{Z})$.

Using this diagram, to show that the top map is nonzero, and thus an isomorphism, it suffices to show that the lower left map and the bottom map are each isomorphisms. We state each of these as a separate lemma (one in dual form).

Lemma A.2. $\Delta_* : H_{2p-3}(P_{p-1}(\Sigma^{-1}H\mathbb{Z})) \rightarrow H_{2p-3}(\Sigma^{-2}P_{p-1}(\Sigma H\mathbb{Z}))$ is an isomorphism of one dimensional \mathbb{Z}/p -modules.

Proof. When $p = 2$, Δ is an equivalence, and so Δ_* is an isomorphism.

When p is odd, the situation is more complicated, and we proceed as follows. We have a commutative diagram

$$\begin{array}{ccc}
H_{2p-3}(P_{p-1}(\Sigma^{-1}H\mathbb{Z})) & \xrightarrow{\Delta_*} & H_{2p-3}(\Sigma^{-2}P_{p-1}(\Sigma H\mathbb{Z})) \\
\downarrow \wr & & \downarrow \wr \\
H_{2p-3}(P_2(\Sigma^{-1}H\mathbb{Z})) & & \\
\downarrow & & \downarrow \wr \\
H_{2p-3}(\Sigma^{-1}H\mathbb{Z}) & \xrightarrow[\sim]{\Delta_*} & H_{2p-3}(\Sigma^{-1}H\mathbb{Z})
\end{array}$$

with indicated isomorphisms. Thus, to show the top map is an isomorphism, we need to check that the lower left map is an isomorphism. Equivalently, we need to check that

$$\delta_* : H_{2p-3}(\Sigma^{-1}H\mathbb{Z}) \rightarrow H_{2p-3}(\Sigma D_2 \Sigma^{-1}H\mathbb{Z})$$

is zero. The map $\delta : \Sigma^{-1}HZ \rightarrow \Sigma D_2 \Sigma^{-1}HZ$ factors through

$$\Delta : \Sigma^2 D_2 \Sigma^{-2}HZ \rightarrow \Sigma D_2 \Sigma^{-1}HZ,$$

and this map is zero on H_{2p-3} : the range is one dimensional, spanned by the suspension of a $*$ -decomposable of the form $x * y$, with $x \in H_{-1}(\Sigma^{-1}HZ)$ and $y \in H_{2p-3}(\Sigma^{-1}HZ)$. But nonzero $*$ -decomposables are never in the image of $\Delta_* : H_*(\Sigma D_2(X)) \rightarrow H_*(D_2(\Sigma X))$. \square

With our final lemma, we have reached the heart of the matter.

Lemma A.3. $\delta^* : H^{2p-1}(\Sigma D_p(\Sigma H\mathbb{Z})) \rightarrow H^{2p-1}(P_{p-1}(\Sigma H\mathbb{Z}))$ is an isomorphism of one dimensional \mathbb{Z}/p -modules.

Proof. Since $\Sigma H\mathbb{Z}$ is 0-connected, the Goodwillie tower $P_r(\Sigma H\mathbb{Z})$ converges strongly to $\Sigma^\infty \Omega^\infty(\Sigma H\mathbb{Z}) = \Sigma^\infty S^1$. Thus the associated 2nd quadrant spectral sequence converges strongly to $H^*(S^1)$. For this to happen, $\mathcal{P}^1(x)$ must be in the image of δ^* , where $x \in H^1(P_{p-1}(\Sigma H\mathbb{Z}))$ is a nonzero element, for otherwise $\mathcal{P}^1(x) \neq 0 \in H^{2p-1}(S^1)$.

Thus δ^* is nonzero, and is thus an isomorphism. \square

Remark A.4. In work in progress, the author is studying the spectral sequence converging to $H^*(\Omega^\infty X)$ with $E_1^{-r,*+r} = H^*(D_r X)$. The sort of argument just given generalizes to show that the first interesting differential is $d_{p-1} : H^{*-1}(D_p X) \rightarrow H^*(X)$. This differential is determined by $H^*(X)$ as a module over the Steenrod algebra, and has image imposing the unstable condition on $H^*(X)$.

REFERENCES

- [AGM] J. F. Adams, J. H. Gunawardena, and H. Miller, *The Segal conjecture for elementary abelian p -groups*, *Topology* **24** (1985), 435–460.
- [ACD] A. Adem, R. L. Cohen, and W. G. Dwyer, *Generalized Tate homology, homotopy fixed points and the transfer*, *Algebraic topology* (Evanston, 1988), *A.M.S.Cont.Math.* **96** (1989), 1–13.
- [AK] S. T. Ahearn and N. J. Kuhn, *Product and other fine structure in polynomial resolutions of mapping spaces*, *Alg. Geom. Topol.* **2** (2002), 591–647.
- [B1] A. K. Bousfield, *Uniqueness of infinite deloopings for K -theoretic spaces*, *Pacific J. Math.* **129** (1987), 1–31.
- [B2] A. K. Bousfield, *On the telescopic homotopy theory of spaces*, *Trans. A. M. S.* **353** (2001), 2391–2426.
- [BMMS] R. R. Bruner, J. P. May, J. McClure, and M. Steinberger *H_∞ Ring Spectra and their Applications*, *Springer L. N. Math.* **1176**, 1986.
- [CLM] F. R. Cohen, T. Lada, and J. P. May, *The Homology of Iterated Loop Spaces*, *Springer L. N. Math.* **533**, 1976.
- [DHS] E. S. Devinatz, M. J. Hopkins, and J. H. Smith, *Nilpotence and stable homotopy theory. I*, *Ann. Math.* **128** (1988), 207–241.
- [EKMM] A.D. Elmendorf, I. Kriz, M.A. Mandell, J.P. May, *Rings, modules, and algebras in stable homotopy theory*, *A.M.S. Math. Surveys and Monographs* **47**, 1997.
- [G1] T. G. Goodwillie, *Calculus I: the first derivative of pseudoisotopy*, *K-theory* **4** (1990), 1–27.
- [G2] T. G. Goodwillie, *Calculus II: analytic functors*, *K-theory* **5** (1992), 295–332.

- [G3] T. G. Goodwillie, *Calculus III: the Taylor series of a homotopy functor*, preprint, Brown University, June, 2002.
- [GM] J. P. C. Greenlees and J. P. May, *Generalized Tate cohomology*, *Memoirs A.M.S.* **113** (1995), no. 543.
- [GS] J. P. C. Greenlees and H. Sadofsky, *The Tate spectrum of v_n -periodic complex oriented theories*, *Math. Zeit.* **222** (1996), 391–405.
- [Gun] J. H. C. Gunawardena, *Segal's conjecture for cyclic groups of (odd) prime order*, J. T. Knight Prize Essay, Cambridge, 1980.
- [HopSm] M. J. Hopkins and J. H. Smith, *Nilpotence and stable homotopy theory. II*, *Ann. Math.* **148** (1998), 1–49.
- [H] M. Hovey, *Cohomological Bousfield classes*, *J. Pure Appl. Alg.* **103** (1995), 45–59.
- [HSa] M. Hovey and H. Sadofsky, *Tate cohomology lowers chromatic Bousfield classes*, *Proc. A. M. S.* **124** (1996), 3579–3585.
- [HSt] M. Hovey and N. P. Strickland, *Morava K -theories and localisation*, *Memoirs A.M.S.* **139** (1999), no. 666.
- [JW] J. D. S. Jones and S. A. Wegmann, *Limits of stable homotopy and cohomotopy groups*, *Math. Proc. Camb. Phil. Soc.* **94** (1983), 473–482.
- [K11] J. R. Klein, *The dualizing spectrum of a topological group*, *Math. Ann.* **319** (2001), 421–456.
- [K12] J. R. Klein, *Axioms for generalized Farrell-Tate cohomology*, *J. Pure Appl. Algebra* **172** (2002), 225–238.
- [Ku1] N. J. Kuhn, *Morava K -theories and infinite loop spaces*, Algebraic topology (Arcata, CA, 1986), Springer Lecture Notes in Math. **1370** (1989), 243–257.
- [Ku2] N. J. Kuhn, *Localization of André-Quillen-Goodwillie towers, and the periodic homology of infinite loopspaces*, preprint, 2003.
- [L] W. H. Lin, *On conjectures of Mahowald, Segal and Sullivan*, *Math. Proc. Camb. Phil. Soc.* **87** (1980), 449–458.
- [MS] M. Mahowald and P. Shick, *Root invariants and periodicity in stable homotopy theory*, *Bull. London Math. Soc.* **20** (1988), 262–266.
- [MO] A. J. Mauer-Oats, *Goodwillie Calculi*, thesis, University of Illinois, 2002.
- [MM] J. P. May and J. E. McClure, *A reduction of the Segal conjecture*, Current trends in Algebraic Topology (London, Ontario, 1981), Canadian Math. Soc. Conference Proc. **2**, Part 2 (1982), 209–222.
- [McC] R. McCarthy, *Dual calculus for functors to spectra*, Homotopy methods in algebraic topology (Boulder, 1999), A.M.S. Contemp. Math. Series **271** (2001), 183–215.
- [R1] D. C. Ravenel, *Localization with respect to certain periodic homology theories*, *Amer. J. Math.* **106** (1984), 351–414.
- [R2] D. C. Ravenel, *Nilpotence and Periodicity in Stable Homotopy Theory*, *Annals of Mathematics Study* **128**, 1992.
- [WW1] M. Weiss and B. Williams, *Automorphisms of manifolds and algebraic K -theory II*, *J. Pure Appl. Algebra* **62** (1989), 47–107.
- [WW2] M. Weiss and B. Williams, *Assembly, Novikov conjectures, index theorems and rigidity*, Vol. 2 (Oberwolfach, 1993), London Math. Soc. Lecture Note Ser. **227** (1995), 332–352.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF VIRGINIA, CHARLOTTESVILLE, VA 22903

E-mail address: njk4x@virginia.edu