

THE GHOST DIMENSION OF A RING

MARK HOVEY AND KEIR LOCKRIDGE

ABSTRACT. We introduce the concept of the ghost dimension $\text{gh. dim. } R$ of a ring R . This is the longest nontrivial chain of maps in the derived category emanating from a perfect complex such that each map is zero on homology. We show that $\text{gh. dim. } R \leq \text{w. dim. } R$, with equality if R is coherent or $\text{w. dim. } R = 1$.

INTRODUCTION

Given a ring R , one can study R by looking at properties of elements of R , properties of the category $R\text{-mod}$ of right R -modules, or properties of its unbounded derived category $\mathcal{D}(R)$. If we allow ourselves to remember the object R (thus ruling out Morita and derived equivalences), these approaches all contain the same information. However, there are generalized rings where not all of these approaches are available. In algebraic topology, for example, there are structured ring spectra E (often called S -algebras or symmetric ring spectra). For such an E , there are no elements, and there is no abelian category of E -modules. But there is a triangulated category $\mathcal{D}(E)$ analogous to $\mathcal{D}(R)$. If one wants to define various homological dimensions of E , then, one needs to look at how to define homological dimensions of R solely in terms of the derived category $\mathcal{D}(R)$ (with its distinguished object R). In this paper, we describe how to recover the right global dimension of R from $\mathcal{D}(R)$ and its distinguished object R , and we introduce another invariant, the ghost dimension of R , that we originally thought would be the weak dimension or the finitistic right global dimension of R . However, we now believe this to be a new invariant of R .

In more detail, given a ring R , let $\mathcal{D}(R)$ denote the unbounded derived category of right R -modules. A map $f: X \rightarrow Y$ in $\mathcal{D}(R)$ is called a **ghost** if $H_*f = 0$. (Note that to define a ghost, we need to remember the object R , since $H_*X = \mathcal{D}(R)(s^*R, X)$, so ghosts are not derived invariant). A complex X is said to have **ghost dimension** n , written $\text{gh. dim. } X = n$, if every composite

$$X \xrightarrow{f_1} Y_1 \xrightarrow{f_2} \dots \xrightarrow{f_{n+1}} Y_{n+1}$$

of $n + 1$ ghosts is 0 in $\mathcal{D}(R)$, and there exists a composite of n ghosts from X that is not 0 in $\mathcal{D}(R)$. Since the complex R itself has ghost dimension 0, one might think that the ghost dimension of a complex is analogous to the projective dimension of a module. Hence, the maximal ghost dimension should be the right global dimension of R . This is correct, and we prove it in Theorem 1.5, following the second author's thesis [Loc06, Theorem 4.4.4]. Now a finitely presented module is analogous to a perfect complex in $\mathcal{D}(R)$, so the maximal ghost dimension of a perfect complex should be related to the maximal projective dimension of a finitely

presented module, or perhaps to the weak dimension of R (since these two invariants are equal when R is coherent). However, we have not been able to prove this, and so we define the **ghost dimension of R** , written $\text{gh. dim. } R$, to be the maximum ghost dimension of any perfect complex in $\mathcal{D}(R)$ (or ∞ if there is no such maximum).

The authors and G. Puninski proved that ghost dimension 0 is equivalent to weak dimension 0 (i.e., von Neumann regular) in [HLP07], although the authors referred to R having ghost dimension 0 as R satisfying the strong generating hypothesis, based on analogy with the stable homotopy category in algebraic topology. In the present paper, we prove that $\text{w. dim. } R \leq \text{gh. dim. } R$, with equality holding if R is right coherent or $\text{w. dim. } R \leq 1$. We have not been able to resolve the general case, though we suspect there is a counterexample.

Ghost dimension in the stable module category of a finite group has been studied in [CCM07].

1. GLOBAL DIMENSION

The object of this section is to prove that $\text{r. gl. dim. } R$ is the maximum ghost dimension of a complex in $\mathcal{D}(R)$. Along the way, we prove some basic facts about ghost dimension that we will need later.

We first recall that the ghost maps \mathcal{I} are part of a projective class $(\mathcal{P}, \mathcal{I})$ in $\mathcal{D}(R)$ [Chr98, Section 8]. Here \mathcal{P} is the collection of all complexes isomorphic in $\mathcal{D}(R)$ to a complex of projectives with zero differential. To say that $(\mathcal{P}, \mathcal{I})$ is a projective class means three things:

- (1) $f: X \rightarrow Y \in \mathcal{I}$ if and only if the composite fg is 0 for all $g: P \rightarrow X$ with $P \in \mathcal{P}$;
- (2) $P \in \mathcal{P}$ if and only if $fg = 0$ for all $f: X \rightarrow Y \in \mathcal{I}$ and all $g: P \rightarrow X$; and
- (3) For all $X \in \mathcal{D}(R)$ there is a cofiber sequence

$$P \rightarrow X \xrightarrow{f} Y \rightarrow \Sigma P$$

with $P \in \mathcal{P}$ and $f \in \mathcal{I}$.

These conditions are all easy to check for the ghost projective class. For example, given X , choose a free module P_n mapping onto $H_n X$ for all n , and let $P = \bigoplus_n S^n P_n$, where $S^n P_n$ is the complex consisting of P_n concentrated in degree n . There is an obvious map $S^n P_n \rightarrow X$ that is onto in homology, from which it follows that the cofiber $X \rightarrow Y$ is a ghost.

As Christensen describes in [Chr98, Theorem 3.5], there is an induced projective class $(\mathcal{P}_{n+1}, \mathcal{I}^{n+1})$, where \mathcal{I}^{n+1} consists of the $n+1$ -fold composites of ghost maps, and so, by definition, \mathcal{P}_{n+1} is the collection of complexes of ghost dimension $\leq n$.

In particular, from Christensen's work we deduce the following proposition.

Proposition 1.1. *Let R be a ring.*

- (1) *A complex $X \in \mathcal{D}(R)$ has $\text{gh. dim. } X \leq n$ if and only if there is an exact triangle*

$$Z \rightarrow X \rightarrow P \rightarrow sZ$$

in $\mathcal{D}(R)$, where $\text{gh. dim. } Z \leq n - 1$ and $\text{gh. dim. } P = 0$, and s is the shift functor.

- (2) *X has finite ghost dimension if and only if it is in the thick subcategory generated by the complexes of projective modules with zero differential.*

Recall that a full subcategory \mathcal{C} is thick if it is closed under shifts and summands, and if two out of three objects in an exact triangle are in \mathcal{C} , so is the third (where we view an object and its shift as the same). Note also that this means that all perfect complexes have finite ghost dimension, since they are in the thick subcategory generated by the complex R .

Proof. The first statement follows from [Chr98, Theorem 3.5] and the discussion immediately preceding it. For the second statement, note first that complexes of projective modules with zero differential have ghost dimension 0, so [Chr98, Note 3.6] implies that everything in the thick subcategory generated by them has finite ghost dimension. For the converse, we proceed by induction on the ghost dimension of X . The induction step follows from part (a). The base case is clear, since every complex of ghost dimension 0 is a complex of projectives with zero differential. \square

Corollary 1.2. *Suppose X is a complex of projectives such that $X_i = 0$ for all but $n + 1$ values of i . Then $\text{gh. dim. } X \leq n$.*

Proof. Use induction and part (1) of Proposition 1.1. \square

Christensen's work also gives us the following proposition, which is a restatement of [Chr98, Theorem 8.3].

Proposition 1.3. *Suppose $\text{r. gl. dim. } R \leq n$. Then every complex X in $\mathcal{D}(R)$ has $\text{gh. dim. } X \leq n$.*

One way to think about this proposition is in terms of the spectral sequence

$$E_2^{s,t} = \text{Ext}_R^s(H_*X, H_*Y)_{t-s} \Rightarrow [X, Y]_s.$$

This is sometimes known as the hypercohomology spectral sequence in algebra, and the universal coefficient spectral sequence in topology. It is conditionally convergent, and the associated filtration on $[X, Y]$ is the ghost filtration. This spectral sequence is described in [Chr98, Section 4] for a general projective class, but is well-known in this case. In any case, if the right global dimension of R is $\leq n$, then this spectral sequence has only $n + 1$ nonzero rows at E_2 , so of course also at E_∞ , so the ghost dimension is at most n .

To prove the converse of this proposition, we note one simple source of ghosts. For any complex X , we can let $X[m, \infty]$ be the complex with $X[m, \infty]_i = X_i$ if $i \geq m$ and 0 if $i < m$, with the nonzero differentials coinciding with those of X . Then there is an obvious map $X \rightarrow X[m, \infty]$ of complexes, and this map will be a ghost if X has no homology in degrees $\geq m$. In particular, we get the following lemma.

Lemma 1.4. *If X is the projective resolution of a module M , then $\text{gh. dim. } X = \text{proj. dim. } M$. In particular, $\text{gh. dim. } R$ is at least as big as the maximal finite projective dimension of an FP_∞ -module.*

Recall that an FP_∞ -module is a module which has a projective resolution by finitely generated projectives. If the FP_∞ module M has finite projective dimension, then there is a projective resolution of M that is a perfect complex, so the second statement of the lemma follows from the first.

Proof. Write X as the sequence

$$\cdots \xrightarrow{d_{k+1}} X_k \xrightarrow{d_k} X_{k-1} \cdots \xrightarrow{d_1} X_0.$$

Suppose $\text{proj. dim. } M = n$, so that $X_k = 0$ for $k > n$. Then Corollary 1.2 implies $\text{gh. dim. } X \leq n$. Now, the sequence of ghosts

$$X \rightarrow X[1, \infty] \rightarrow X[2, \infty] \rightarrow \cdots \rightarrow X[n, \infty]$$

cannot be 0 in $\mathcal{D}(R)$, since if it were it would be chain homotopic to 0, and so null homotopic, so that $d_n: X_n \rightarrow X_{n-1}$ would be a split monomorphism. This would mean that the kernel of d_{n-2} is projective, so that $\text{proj. dim. } M \leq n - 1$. Hence $\text{gh. dim. } X \geq n$ (even if $n = \infty$). \square

This lemma together with Proposition 1.3 gives us the following theorem, due to the second author [Loc06, Theorem 4.4.4].

Theorem 1.5. *The maximum ghost dimension of an object in $\mathcal{D}(R)$ is $\text{r. gl. dim. } R$. Hence $\text{gh. dim. } R \leq \text{r. gl. dim. } R$.*

2. WEAK DIMENSION

The object of this section is to show that $\text{gh. dim. } R \geq \text{w. dim. } R$, with equality holding when R is right coherent or when $\text{w. dim. } R \leq 1$.

We first explain why the coherent case is easier.

Proposition 2.1. *Suppose R is right coherent. Then $\text{gh. dim. } R \leq \text{w. dim. } R$.*

Proof. We will show that in the spectral sequence

$$E_2^{s,t} = \text{Ext}_R^s(H_*X, H_*Y)_{t-s} \Rightarrow [X, Y]_s.$$

discussed after Proposition 1.3, E_2 vanishes for $s > \text{w. dim. } R$ if X is perfect. Hence the same will be true for E_∞ . Since $E_\infty^{s,*}$ represents the maps that are s -fold composites of ghost maps but not $(s-1)$ -fold such composites, we conclude that every nonzero map is a composite of at most $\text{w. dim. } R$ ghosts, giving us the desired result.

First note that because R is right coherent, finitely presented right modules are closed under kernels and cokernels. Thus if X is a perfect complex, then H_iX is a finitely presented R -module for all i . On the other hand, again because R is right coherent, $\text{proj. dim. } M$ is equal to the flat dimension of M for all finitely presented M . Indeed, because R is right coherent, there is a projective resolution

$$\cdots \rightarrow P_k \rightarrow P_{k-1} \rightarrow \cdots \rightarrow P_0 \rightarrow M$$

of M in which each P_k is finitely generated projective. If the flat dimension of M is n , then the kernel of $P_{n-1} \rightarrow P_{n-2}$ is flat. But it is also finitely presented, again since R is right coherent. So it is projective, and so $\text{proj. dim. } M \leq n$. Hence we have $\text{proj. dim. } M \leq \text{w. dim. } R$ for all finitely presented M , completing the proof. \square

The general case is addressed in the following theorem.

Theorem 2.2. *Suppose R is a ring. Then $\text{w. dim. } R \leq \text{gh. dim. } R$. In particular, if R is right coherent, then $\text{gh. dim. } R = \text{w. dim. } R$.*

Proof. Let $n = \text{gh. dim. } R$, which we can of course assume is finite. Let M be an R -module, and let

$$\cdots \xrightarrow{d_{k+1}} P_k \xrightarrow{d_k} P_{k-1} \xrightarrow{d_{k-1}} \cdots \xrightarrow{d_1} P_0 \xrightarrow{d_0} M$$

be a free resolution of M . Let M_{k+1} be the kernel of d_k , so we have short exact sequences

$$0 \rightarrow M_{k+1} \xrightarrow{i_{k+1}} P_k \xrightarrow{q_k} M_k \rightarrow 0,$$

with $M_0 = M$ and $i_k q_k = d_k$. We need to show that M_n is flat, which we will do by applying [Lam99, Theorem 4.23] to the short exact sequence

$$0 \rightarrow M_{n+1} \rightarrow P_n \rightarrow M_n \rightarrow 0.$$

So take a $c \in M_{n+1}$. To use [Lam99, Theorem 4.23], we need to construct a map $P_n \rightarrow M_{n+1}$ that takes c to itself. We first build a perfect subcomplex of $P = P_*$ that contains c (or really $i_{n+1}c$) by a simple induction. To start, c lies in a finitely generated summand Q_n of P_n . But then $d(Q_n)$ lies in a finitely generated summand Q_{n-1} of P_{n-1} . Continuing in this fashion, we get a perfect subcomplex Q of P with $c \in Q_n$.

Now consider the sequence of ghosts

$$Q \rightarrow P[1, \infty] \rightarrow P[2, \infty] \rightarrow \cdots \rightarrow P[n, \infty],$$

where the first map is the composite $Q \xrightarrow{j} P \rightarrow P[1, \infty]$. Since $\text{gh. dim. } Q \leq n$, this composite is nullhomotopic. Thus there are maps $r: Q_n \rightarrow P_{n+1}$ and $s: Q_{n-1} \rightarrow P_n$ such that $d_{n+1}r + sd_n = j_n$, the inclusion of Q_n into P_n . Consider the map

$$P_n \xrightarrow{\tau} Q_n \xrightarrow{r} P_{n+1} \xrightarrow{q_{n+1}} M_{n+1},$$

where τ is a splitting of the inclusion j_n . Then

$$i_{n+1}q_{n+1}r\tau(c) = d_{n+1}rc = i_{n+1}c - sd_n c = i_{n+1}c,$$

since c is a cycle. This means $q_{n+1}r\tau(c) = c$, as required to apply [Lam99, Theorem 4.23]. \square

We have not been able to prove that $\text{gh. dim. } R = \text{w. dim. } R$ for a general ring R . However, this is true if $\text{w. dim. } R = 1$. To prove this, we need the following slight generalization of the theorem of Villamayor [Lam99, Theorem 4.23] that we have already used in the proof of Theorem 2.2.

Lemma 2.3. *Suppose we have a short exact sequence of right modules over a ring R*

$$0 \rightarrow K \rightarrow P \rightarrow M \rightarrow 0$$

where P is projective. Then the following conditions are equivalent.

- (1) M is flat.
- (2) For every finitely generated submodule L of K , there is a map $\theta: P \rightarrow K$ such that $\theta(x) = x$ for all $x \in L$.
- (3) For every element c of K , there is a map $\theta: P \rightarrow K$ such that $\theta(c) = c$.

Proof. Find a Q such that $P \oplus Q$ is free. Then we have a short exact sequence

$$0 \rightarrow K \oplus Q \rightarrow P \oplus Q \rightarrow M \rightarrow 0.$$

We can now apply [Lam99, Theorem 4.23]. More precisely, suppose M is flat, and L is a finitely generated submodule of K . Then, by [Lam99, Theorem 4.23], there

is a map $f: P \oplus Q \rightarrow K \oplus Q$ such that $f(x, 0) = (x, 0)$ for all $x \in L$. But then we can define θ to be the composite

$$P \rightarrow P \oplus Q \xrightarrow{f} K \oplus Q \rightarrow K$$

to see that (1) \Rightarrow (2). It is obvious that (2) \Rightarrow (3). To see that (3) \Rightarrow (1), take an element (c, q) of $K \oplus Q$. By assumption, there is a map $\theta: P \rightarrow K$ such that $\theta(c) = c$. Define $f: P \oplus Q \rightarrow K \oplus Q$ by $f(x, y) = (\theta(x), y)$. Then $f(c, q) = (c, q)$. So [Lam99, Theorem 4.23] implies that M is flat. \square

We can use this improvement to characterize rings of weak dimension 1.

Proposition 2.4. *A ring R has $\text{w. dim. } R \leq 1$ if and only if, for every diagram*

$$P_2 \xrightarrow{d_2} P_1 \xrightarrow{d_1} P_0$$

of finitely generated projectives with $d_1 d_2 = 0$, there is a map $f: P_1 \rightarrow P_1$ with $d_1 f = 0$ and $f d_2 = d_2$.

Proof. Suppose R has weak dimension ≤ 1 , and consider the short exact sequence

$$0 \rightarrow \ker d_1 \rightarrow P_1 \rightarrow \text{im } d_1 \rightarrow 0.$$

Since R has weak dimension ≤ 1 , $\text{im } d_1 \subseteq P_0$ is flat. Apply Lemma 2.3 to the finitely generated submodule $\text{im } d_2$ of $\ker d_1$ to get a map $f: P_1 \rightarrow \ker d_1$ such that $f(x) = x$ for all $x \in \text{im } d_2$. This means that $d_1 f = 0$ and $f d_2 = d_2$, as required.

For the converse, it suffices to show that $\text{Tor}_2^R(N, -) = 0$ for all finitely presented N . Take a finite presentation

$$P_1 \xrightarrow{d_1} P_0 \rightarrow N \rightarrow 0$$

of N , where P_1 and P_0 are free and finitely generated. This gives us the short exact sequence

$$0 \rightarrow K \rightarrow P_1 \rightarrow \text{im } d_1 \rightarrow 0.$$

We need to show that $\text{im } d_1$ is flat. Using [Lam99, Theorem 4.23], it suffices to show that for any $c \in K$, there is a map $f: P_1 \rightarrow K$ such that $f(c) = c$. But c then gives us the complex

$$R \xrightarrow{c} P_1 \xrightarrow{d_1} P_0$$

of finitely generated projectives. By hypothesis, then, there is a map $f: P_1 \rightarrow K$ such that $f(c) = c$. \square

Theorem 2.5. *If $\text{w. dim. } R \leq 1$, then $\text{gh. dim. } R = \text{w. dim. } R$.*

Proof. If $\text{w. dim. } R = 0$, this is proved in [HLP07, Theorem 1.3]. Thus, in view of Theorem 2.2, we must show that $\text{w. dim. } R = 1$ implies $\text{gh. dim. } R \leq 1$. We do this by showing that every perfect complex is a retract of a direct sum of complexes with only two nonzero entries. In view of Corollary 1.2, this will complete the proof.

Suppose P is a perfect complex, with maps $d_n: P_n \rightarrow P_{n-1}$. Since $d_n d_{n+1} = 0$, Proposition 2.4 implies that there are maps $f_n: P_n \rightarrow \ker d_n$ such that $f_n d_{n+1} = d_{n+1}$. For each n , choose a surjection $q_n: Q_n \rightarrow \ker d_n$ where Q_n is projective, and choose a lift $g_n: P_n \rightarrow Q_n$ so that $q_n g_n = f_n$. Now define a two-term chain complex T^n with $T_n^n = P_n$ and $T_{n-1}^n = Q_{n-1}$, with differential $g_{n-1} d_n$.

We claim that there are chain maps

$$P \xrightarrow{\phi} \bigoplus_k T^k \xrightarrow{\psi} P$$

exhibiting P as a retract of $\bigoplus_k T^k$. Note that the n th term of the complex $\bigoplus_k T^k$ is $Q_n \oplus P_n$, with differential $D_n(x, y) = (g_{n-1}d_n y, 0)$. Define $\psi_n: Q_n \oplus P_n \rightarrow P_n$ by $\psi_n(x, y) = q_n(x) + y$. Then

$$d_n \psi_n(x, y) = d_n y$$

and

$$\psi_{n-1} D_n(x, y) = \psi_{n-1}(g_{n-1}d_n y, 0) = q_{n-1}g_{n-1}d_n y = f_{n-1}d_n y = d_n y,$$

so ψ is a chain map.

Now define $\phi_n: P_n \rightarrow Q_n \oplus P_n$ by $\phi_n(x) = (g_n(x), x - f_n x)$. Then

$$D_n \phi_n(x) = (g_{n-1}d_n(x - f_n x), 0) = (g_{n-1}d_n x, 0)$$

and

$$\phi_{n-1} d_n(x) = (g_{n-1}d_n x, d_n x - f_{n-1}d_n x) = (g_{n-1}d_n x, 0).$$

Thus ϕ is a chain map, and one can readily verify that

$$\psi_n \phi_n(x) = \psi_n(g_n x, x - f_n x) = q_n g_n x + x - f_n x = x,$$

showing that P is a retract of $\bigoplus_n T^n$. \square

REFERENCES

- [CCM07] Sunil K. Chebolu, J. Daniel Christensen, and Ján Mináč, *Ghosts in modular representation theory*, preprint, 2007.
- [Chr98] J. Daniel Christensen, *Ideals in triangulated categories: phantoms, ghosts and skeleta*, Adv. Math. **136** (1998), no. 2, 284–339. MR MR1626856 (99g:18007)
- [HLP07] Mark Hovey, Keir Lockridge, and Gena Puninski, *The generating hypothesis in the derived category of a ring*, Math. Z. **256** (2007), no. 4, 789–800. MR MR2308891
- [Lam99] T. Y. Lam, *Lectures on modules and rings*, Graduate Texts in Mathematics, vol. 189, Springer-Verlag, New York, 1999. MR MR1653294 (99i:16001)
- [Loc06] Keir H. Lockridge, *The generating hypothesis in general stable homotopy categories*, Ph.D. thesis, University of Washington, 2006.

DEPARTMENT OF MATHEMATICS, WESLEYAN UNIVERSITY, MIDDLETOWN, CT 06459

E-mail address: hovey@member.ams.org

DEPARTMENT OF MATHEMATICS, WAKE FOREST UNIVERSITY, WINSTON-SALEM, NC 27109

E-mail address: lockrikh@wfu.edu