

Basic constructions in rational homotopy theory of function spaces

Urtzi Buijs and Aniceto Murillo*

Departamento de Algebra, Geometría y Topología,
Universidad de Málaga,
Ap. 59, 29080 Málaga, Spain

June 17, 2005

Abstract

Via the Bousfield-Gugenheim realization functor, and starting from the Brown Szczarba model of a function space, we give a functorial framework to describe basic objects and maps concerning the rational homotopy type of function spaces and its path components.

1 Introduction

There are two choices to describe the equivalence between the homotopy categories of rational nilpotent CW-complexes of finite type and that of commutative differential graded algebras over the rationals. These choices depends on how to realize a given algebra. On one hand, one may proceed as in [16], considering a Sullivan model and then build its realization as a Postnikov tower whose topological description mimics the algebraic properties of the model. This is highly geometrical but quite unnatural. On the other hand, one can take a general algebra and realizing it as a simplicial set via the

*Partially supported by the Ministerio de Ciencia y Tecnología grant MTM2004-06262 and by the Junta de Andalucía grant FQM-213.

2000 M.S.C.: 55P62.

Keywords: Function space, mapping space, Sullivan model, rational homotopy theory.

Bousfield-Gugenheim realization functor [1]. This might be less transparent but totally functorial.

In particular both options are available to the study of the rational homotopy type of the function (or mapping) space $\mathcal{F}(X, Y)$. The first approach was first followed by Thom [18] and then by the fundamental work of Haefliger [8] where he supplied a model for the path component of sections, homotopic to a fixed one, of a nilpotent bundle. In particular, a model of a given component of a mapping space is produced. Later on, this approach was further developed by other authors [4, 14, 15, 19]. However, due to the non functorial description of the Haefliger model, it is not easy to use it to describe further geometrical behavior of function spaces.

Under the second, and more functorial approach, it is essential the paper of Brown and Szczarba [3] in which it is proved that the Bousfield-Gugenheim realization of a particular commutative differential \mathbb{Z} -graded algebra (CDGA henceforth), which already appears in [8], is homotopy equivalent to the (non connected!) function space $\mathcal{F}(X, Y)$. It is also shown how to obtain from this algebra a quotient CDGA whose realization is a given path component of the function space, and therefore, the Haefliger model of this component is retrieved.

In this paper we continue with the study under this natural approach, and present models for some of the basic objects and maps concerning function spaces. By a model of a non connected object (or a map between them) we mean a \mathbb{Z} -graded CDGA (or a morphism) whose simplicial realization has the homotopy type as the singular simplicial approximation of the chosen object. We also present an explicit way of restricting algebraically to components obtaining Sullivan models of nilpotent spaces and maps relating functions spaces, retrieving in particular the Haefliger model for its components.

We now state our main results: For any function space $\mathcal{F}(X, Y)$ we shall always assume X and Y CW-complexes with X finite, so that $\mathcal{F}(X, Y)$ is itself a CW-complex [13], and with both X and Y nilpotent of finite type over \mathbb{Q} , so that the components of $\mathcal{F}(X, Y)$ are also nilpotent and of finite \mathbb{Q} -type [10, Thm.2.5]. Given $f: X \rightarrow Y$, denote by $\mathcal{F}(X, Y; f)$ the component of $\mathcal{F}(X, Y)$ containing f . We first give an explicit model of the evaluation map which includes in particular the original result of Haefliger [8, Thm.3.2] and [11, Thm.1.1].

Theorem 1. *Let $(\Lambda(V \otimes B_*), \tilde{d})$ be the Brown-Szczarba model for $\mathcal{F}(X, Y)$*

(see §2). Then,

$$\omega: (\Lambda V, d) \rightarrow (\Lambda(V \otimes B_*), \tilde{d}) \otimes B, \quad \omega(v) = \sum_i (-1)^{\alpha|b_i|} (v \otimes \beta_i) \otimes b_i,$$

where $\{b_i\}$ is a basis for B , $\{\beta_i\}$ its dual and $\alpha(n)$ is the integer part of $(n+1)/2$, is a model of the evaluation map

$$\omega: \mathcal{F}(X, Y) \times Y \rightarrow Y, \quad \omega(f, x) = f(x).$$

In particular, the map

$$\omega_0: (\Lambda V, d) \rightarrow (\Lambda(V \otimes B_*), \tilde{d}), \quad \omega_0(v) = v \otimes 1^*,$$

is a model of the evaluation of the base point $\omega_0: \mathcal{F}(X, Y) \rightarrow Y$, $\omega_0(f) = f(x_0)$.

Moreover, given $\zeta: (\Lambda V, d) \longrightarrow (\Lambda W, d)$ a Sullivan model for $f: X \rightarrow Y$, there is an explicit morphism $\xi: (\Lambda(V \otimes C_*), \tilde{d}) \longrightarrow (\Lambda(W \otimes C_*), \tilde{d})$ (see §3) between the models of $\mathcal{F}(Z, Y)$ and $\mathcal{F}(Z, X)$ satisfying:

Theorem 2. *The commutative square*

$$\begin{array}{ccc} (\Lambda(W \otimes C_*), \tilde{d}) \otimes C & \xleftarrow{\xi \otimes 1_C} & (\Lambda(V \otimes C_*), \tilde{d}) \otimes C \\ \omega \uparrow & & \uparrow \omega \\ (\Lambda W, d) & \xleftarrow{\zeta} & (\Lambda V, d) \end{array}$$

is a model of

$$\begin{array}{ccc} \mathcal{F}(Z, X) \times Z & \xrightarrow{(f)_* \times 1_Z} & \mathcal{F}(Z, Y) \times Z \\ \omega \downarrow & & \downarrow \omega \\ X & \xrightarrow{f} & Y. \end{array}$$

Next, we show how to obtain from these results the corresponding Sullivan models when considering a fixed component of the function space. Indeed, given $(\Lambda(V \otimes B_*), \tilde{d})$ the model of $\mathcal{F}(X, Y)$ and a map $f: X \rightarrow Y$, there is a Sullivan model (in fact this is the Haefliger model) of the component $\mathcal{F}(X, Y; f)$ of the form $(\Lambda S_f, \bar{d})$ in which $S_f^{\geq 2} = (V \otimes B_*)^{\geq 2}$ and S_f^1 is a quotient of $(V \otimes B_*)^1$ (see §4). Then we prove:

Theorem 3. *The CDGA morphism*

$$\omega_0: (\Lambda V, d) \longrightarrow (\Lambda S, \bar{d}),$$

given by $\omega_0(v) = v \otimes 1^*$ if $v \in V^{\geq 2}$, or its projection over S_f^1 if $v \in V^1$, is a Sullivan model of the evaluation at the base point $\omega_0: \mathcal{F}(X, Y; f) \longrightarrow Y$. Moreover, with the notation of Theorem 2, ξ induces a morphism $\bar{\xi}$ for which the square

$$\begin{array}{ccc} (\Lambda S_g, \bar{d}) & \xleftarrow{\bar{\xi}} & (\Lambda S_{fg}, \bar{d}) \\ \omega_0 \uparrow & & \uparrow \omega_0 \\ (\Lambda W, d) & \xleftarrow{\zeta} & (\Lambda V, d) \end{array}$$

is a Sullivan model of

$$\begin{array}{ccc} \mathcal{F}(Z, X; g) & \xrightarrow{(f)_*} & \mathcal{F}(Z, Y; f \circ g) \\ \omega_0 \downarrow & & \downarrow \omega_0 \\ X & \xrightarrow{f} & Y. \end{array}$$

As in this result, ω_0 is a fibration with fibre $\mathcal{F}^*(X, Y; f)$, the space of pointed maps, we easily obtain the following consequence, a form of which already appeared in [4], [11] or [19].

Corollary 4. *The CDGA $(\Lambda(S_f/V), \bar{d})$ is a Sullivan model of $\mathcal{F}^*(X, Y; f)$.*

Finally, we translate the constructions above to rational homotopy groups by taking the homology of the indecomposables of the given sullivan models and maps.

2 Notation, basic facts and tools

All spaces considered henceforth shall be of the homotopy type of connected CW-complexes. We will heavily rely on known results from homotopy theory of simplicial sets and from rational homotopy theory. For that [7] or [12] and [6] are respectively standard and excellent references. Here, mainly to set the framework in which we work and to fix notation, we recall some basics:

Denote by

$$\mathbf{Top} \begin{array}{c} \xrightarrow{S_*} \\ \xleftarrow{|\cdot|} \end{array} \mathbf{SimpSet}$$

the singular and realization adjoint functors relating the categories of simplicial sets and topological spaces. These induce equivalences between the (pointed if we wish) homotopy category of CW-complexes and the homotopy category of Kan complexes [13]. For each space X and each simplicial set K , We shall denote by $\psi(K): K \xrightarrow{\simeq} S_*|K|$ and $\phi(X): |S_*X| \xrightarrow{\simeq} X$ the corresponding weak equivalences.

On the other hand, if **CDGA** is the category of cochain \mathbb{Z} -graded commutative differential graded algebras, there are adjoint functors [1]

$$\mathbf{SimpSet} \begin{array}{c} \xrightarrow{A_{PL}} \\ \xleftarrow{\langle \cdot \rangle} \end{array} \mathbf{CDGA},$$

where: $A_{PL}(X)$ is the CDGA of *piecewise-linear polynomial forms* on X with rational coefficients and $\langle A \rangle$ is the *simplicial realization* of the CDGA A : For any $q \geq 0$, $\langle A \rangle_q = \text{hom}(A, (A_{PL})_q)$, i.e., the set of morphisms from A to $\Lambda(x_1, \dots, x_q, dx_1, \dots, dx_q)$. The face and degeneracies operators are induced by their counterparts in the simplicial algebra $(A_{PL})_*$ [1, 6].

the composition of the above pairs of adjoint functors produce equivalences when restricted to the homotopy category of rational nilpotent space of finite type over \mathbb{Q} and the homotopy category of connected CDGA's over \mathbb{Q} of the same homotopy type of Sullivan algebras of finite type [1, 6]. Abusing of notation we shall write $A_{PL}(X) = A_{PL}(S_*X)$ and $|A| = |\langle A \rangle|$ for any space X and any CDGA A . Given a non necessarily connected space Z , a CDGA A is a *model* of Z if $S_*(Z)$ and $\langle A \rangle$ are simplicial sets of the same homotopy type. As we are mainly concerned in this paper on mapping spaces, let us describe these objects whenever we take simplicial sets or CDGA's. Given K, L simplicial sets, $\mathcal{F}(K, L)$ is the simplicial set defined

by $\mathcal{F}(K, L)_q = \text{hom}(K \times \Delta[q], L)$, where $\Delta[q] \in \mathbf{SimpSet}$ is the *standard n -simplex*: For each $n \geq 0$, $[n] = \{0, \dots, n\}$ and $(\Delta[n])_q$ is the set of non decreasing maps $[q] \rightarrow [n]$. Call Δ_q to $1_{[q]} \in \Delta[q]_q$. Faces and degeneracies operators on $\mathcal{F}(K, L)$ are defined by $\partial_i f = f \circ (1_K \times \delta_i)$ y $s_i f = f \circ (1_K \times \sigma_i)$. Whenever L is Kan, $\mathcal{F}(K, L)$ is also Kan and it has the same homotopy type of $S_*\mathcal{F}(|K|, |L|)$ via the equivalence

$$\alpha: \mathcal{F}(K, L) \xrightarrow{\cong} S_*\mathcal{F}(|K|, |L|),$$

in which $\alpha(f): \Delta^q \mathcal{F}(|K|, |L|)$ is given by the exponential law applied to the composition

$$|K| \times \Delta^q \xrightarrow{\cong} |K| \times |\Delta[q]| \xrightarrow{\cong} |K \times \Delta[q]| \xrightarrow{|f|} |L|.$$

The evaluation map $\omega: \mathcal{F}(K, L) \times K \rightarrow L$ is defined by $\omega(f, \sigma) = f(\sigma, \Delta_q)$, where $\Delta_q = (0, 1, \dots, q) \in \Delta[q]_q$ is the fundamental q -simplex. This map can be identified with $S_*\omega: S_*(\mathcal{F}(|K|, |L|) \times |K|) \cong S_*\mathcal{F}(|K|, |L|) \times S_*|K| \rightarrow S_*|L|$ via the following result obtained by direct calculation:

Proposition 5. *The following diagram commutes:*

$$\begin{array}{ccc} S_*\mathcal{F}(|K|, |L|) \times S_*|K| & \xleftarrow[\cong]{\alpha \times \psi(K)} & \mathcal{F}(K, L) \times K \\ S_*\omega \downarrow & & \downarrow \omega \\ S_*|L| & \xleftarrow[\cong]{\psi(L)} & L. \end{array}$$

□

On the other hand given CDGA's A and B define $\mathcal{F}(A, B) \in \mathbf{SimpSet}$ by $\mathcal{F}(A, B)_q = \text{hom}(A, (A_{PL})_q \otimes B)$ with faces and degeneracies given by $\partial_i f = (\partial_i \otimes 1_B) \circ f$, $s_j f = (s_j \otimes 1_B) \circ f$. If $\phi: B \rightarrow C$ is a CDGA morphism, we denote by

$$\phi_*: \mathcal{F}(A, B) \rightarrow \mathcal{F}(A, C)$$

the simplicial map given which assigns to each morphism $\psi: A \rightarrow (A_{PL})_q \otimes B$ the composition $\phi_*(\psi) = (1_{(A_{PL})_q} \otimes \phi) \circ \psi$. If A, B, C are connected CDGA's and ϕ is a quasi-isomorphism, then ϕ_* is a weak equivalence.

Next we recall the Brown-Szczarba model of the function space $\mathcal{F}(X, Y)$, with X as always a finite complex and both X and Y nilpotent of finite

type (over \mathbb{Q}). Consider $A = (\Lambda V, d) \xrightarrow{\cong} A_{PL}(Y)$ a Sullivan model (non necessarily minimal!) of Y and $B \xrightarrow{\cong} A_{PL}(X)$ a quasi-isomorphism with B a connected CDGA of finite type. Let $B_* = \text{hom}(B, \mathbb{Q})$ be the differential graded coalgebra dual of B , and consider the CDGA $\Lambda(A \otimes B_*)$ with the natural differential induced by the one on A and by the dual of the differential of B . Now, consider the differential ideal $I \subset \Lambda(A \otimes B_*)$ generated by $1 \otimes 1^* - 1$ and by the elements of the form

$$a_1 a_2 \otimes \beta - \sum_j (-1)^{|a_2||\beta_j'|} (a_1 \otimes \beta_j') (a_2 \otimes \beta_j''),$$

$a_1, a_2 \in A$, $\beta \in B$, and $\Delta\beta = \sum_j \beta_j' \otimes \beta_j''$. Then, the composition

$$\rho: \Lambda(V \otimes B_*) \subset \Lambda(A \otimes B_*) \longrightarrow \Lambda(A \otimes B_*)/I$$

is an isomorphism of graded algebras [3, Thm.1.2], and therefore, considering on $\Lambda(V \otimes B_*)$ the differential $\tilde{d} = \rho^{-1} d \rho$, ρ is also an isomorphism of CDGA's. Then, $(\Lambda(V \otimes B_*), \tilde{d})$ is a model of $\mathcal{F}(X, Y_{\mathbb{Q}})$ [3, Thm.1.3]. In other words, $S_* \mathcal{F}(X, Y)$ and $\langle (\Lambda(V \otimes B_*), \tilde{d}) \rangle$ are homotopy equivalent.

In order to explicitly determine \tilde{d} on $v \otimes \beta \in V \otimes B_*$, calculate $(dv) \otimes \beta + (-1)^{|v|} v \otimes d\beta$ and then use the relations which generate the ideal I to express $(dv) \otimes \beta$ as an element of $\Lambda(V \otimes B_*)$.

3 Compatible models for the evaluation map

In this section we properly state and prove Theorems 1 and 2. For that, fix as before $\varphi: (\Lambda V, d) \xrightarrow{\cong} A_{PL}(Y)$ a Sullivan model of Y (i.e., a KS-complex non necessarily minimal), and $\beta: B \xrightarrow{\cong} A_{PL}(X)$, a quasi-isomorphism in which B is finite dimensional and satisfies an additional condition: the inclusion of the base point $\{x_0\} \hookrightarrow X$ determines a canonical augmentation $\varepsilon: A_{PL}(X) \rightarrow A_{PL}(x_0) \cong \mathbb{Q}$. Thus we may, and will, choose B connected and β preserving augmentations, i.e., $\varepsilon \circ \beta(1) = 1$. Finally, fix $\{b_i\}$ (respec. $\{\beta_i\}$) a basis for B (respec. the dual basis of B_*) and denote by $\alpha(n)$ the integer part of $(n+1)/2$. Then we have:

Theorem 6. *the CDGA morphism*

$$\omega: (\Lambda V, d) \rightarrow (\Lambda(V \otimes B_*), \tilde{d}) \otimes B, \quad \omega(v) = \sum_i (-1)^{\alpha|b_i|} (v \otimes \beta_i) \otimes b_i,$$

is a model of the evaluation map

$$\omega: \mathcal{F}(X_{\mathbb{Q}}, Y_{\mathbb{Q}}) \times X_{\mathbb{Q}} \rightarrow Y_{\mathbb{Q}}, \quad \omega(f, x) = f(x).$$

That is to say, there is a commutative diagram like the following in which the vertical arrows are equivalences:

$$\begin{array}{ccc}
\mathcal{F}(X_{\mathbb{Q}}, Y_{\mathbb{Q}}) \times X_{\mathbb{Q}} & \xrightarrow{\omega} & Y_{\mathbb{Q}} & \text{Top} \\
& & \downarrow \text{\scriptsize } S_* & \\
S_*\mathcal{F}(X_{\mathbb{Q}}, Y_{\mathbb{Q}}) \times S_*X_{\mathbb{Q}} & \xrightarrow{S_*\omega} & S_*(Y_{\mathbb{Q}}) & \\
\downarrow \simeq & & \downarrow \simeq & \text{SimpSet} \\
\langle (\Lambda(V \otimes B_*), \tilde{d}) \otimes B \rangle & \xrightarrow{\langle \omega \rangle} & \langle (\Lambda, V, d) \rangle & \\
& & \uparrow \text{\scriptsize } \langle \cdot \rangle & \\
(\Lambda(V \otimes B_*), \tilde{d}) \otimes B & \xleftarrow{\omega} & (\Lambda V, d) & \text{CDGA}
\end{array}$$

As the inclusion $i: \mathcal{F}(X, Y) \hookrightarrow \mathcal{F}(X, Y) \times X$ on the base point x_0 is clearly modeled by $(\Lambda(V \otimes B_*), \tilde{d}) \otimes B \xrightarrow{1 \otimes \varepsilon} (\Lambda(V \otimes B_*), \tilde{d})$, and since $\omega_0 = \omega \circ i$, we immediately have:

Corollary 7. *The CDGA morphism*

$$\omega_0: (\Lambda V, d) \longrightarrow (\Lambda(V \otimes B_*), \tilde{d}), \quad \omega_0(v) = v \otimes 1^*,$$

is a model of the evaluation map at the base point $\omega_0: \mathcal{F}(X, Y_{\mathbb{Q}}) \longrightarrow Y_{\mathbb{Q}}$, $\omega_0(g) = g(x_0)$. \square

Note that in Theorem 6 we consider $\mathcal{F}(X_{\mathbb{Q}}, Y_{\mathbb{Q}})$ while in Corollary 7 the evaluation map is defined over $\mathcal{F}(X, Y_{\mathbb{Q}})$: although there is a weak equivalence [15, Thm.2.3] $\mathcal{F}(X_{\mathbb{Q}}, Y_{\mathbb{Q}}) \simeq \mathcal{F}(X, Y_{\mathbb{Q}})$, in the first result we evaluate at any point of $X_{\mathbb{Q}}$ while in the second we simply do it at the base point of X .

Proof. Start by choosing simplicial sets $K = S_*(X_{\mathbb{Q}})$, $L = S_*(Y_{\mathbb{Q}})$. As Y is nilpotent, the natural map induced by the adjunction $j: L \xrightarrow{\simeq} \langle A_{PL}(L) \rangle$, $j(\sigma)(\Phi) = \Phi_{\sigma}$, is an equivalence which may be composed with the map

$\langle \varphi \rangle: \langle A_{PL}(L) \rangle \xrightarrow{\cong} \langle A \rangle$ to produce an equivalence $h: L \xrightarrow{\cong} \langle A \rangle$. Apply just definitions to see that the following commutes:

$$\begin{array}{ccc}
\mathcal{F}(K, L) \times K & \xrightarrow[h_* \times 1_K]{\cong} & \mathcal{F}(K, \langle A \rangle) \times K \\
\omega \downarrow & & \downarrow \omega \\
L & \xrightarrow[h]{\cong} & \langle A \rangle.
\end{array} \tag{1}$$

Next, we use the homotopy equivalence $\gamma: \mathcal{F}(A, A_{PL}(K)) \xrightarrow{\cong} \mathcal{F}(K, \langle A \rangle)$ given in [2, Thm.20] which we now describe: Consider the morphism

$$A_{PL}(p_1) \otimes A_{PL}(p_2): A_{PL}(K) \otimes A_{PL}(\Delta[q]) \longrightarrow A_{PL}(K \times \Delta[q]),$$

induced by the projections $K \xleftarrow{p_1} K \times \Delta[q] \xrightarrow{p_2} \Delta[q]$, and the isomorphism of [6, Prop.10.4] $\theta: (A_{PL})_q \longrightarrow A_{PL}(\Delta[q])$ which assigns to each $\Omega \in (A_{PL})_q$ the only element $\theta(\Omega) \in A_{PL}(\Delta[q])$ satisfying $\theta(\Omega)(\Delta_q) = \Omega$. Then, given $\psi: A \longrightarrow (A_{PL})_q \otimes A_{PL}(K)$ an element of $\mathcal{F}(A, A_{PL}(K))_q$,

$$\gamma(\psi) = \langle A_{PL}(p_1) \otimes A_{PL}(p_2) \circ \theta \otimes 1 \circ \psi \rangle \circ j: K \times \Delta[q] \longrightarrow \langle A \rangle.$$

As before, $j: K \times \Delta[q] \xrightarrow{\cong} \langle A_{PL}(K \times \Delta[q]) \rangle$ is the natural equivalence induced by adjunction. On the other hand, define a simplicial map

$$\bar{\omega}: \mathcal{F}(A, A_{PL}(K)) \times \langle A_{PL}(K) \rangle \longrightarrow \langle A \rangle$$

which sends each $\psi: A \longrightarrow (A_{PL})_q \otimes A_{PL}(K)$ and $\phi: A_{PL}(K) \rightarrow (A_{PL})_q$ to the composition

$$\bar{\omega}(\psi, \phi): A \xrightarrow{\psi} (A_{PL})_q \otimes A_{PL}(K) \xrightarrow{1 \cdot \phi} (A_{PL})_q.$$

It is then straightforward to check that the following commutes:

$$\begin{array}{ccc}
\mathcal{F}(K, \langle A \rangle) \times K & \xleftarrow[\cong]{\gamma \times j^{-1}} & \mathcal{F}(A, A_{PL}(K)) \times \langle A_{PL}(K) \rangle \\
\omega \downarrow & & \downarrow \bar{\omega} \\
\langle A \rangle & \xlongequal{\quad\quad\quad} & \langle A \rangle.
\end{array} \tag{2}$$

For the simplicial map $\bar{\omega}: \mathcal{F}(A, B) \times \langle B \rangle \rightarrow \langle A \rangle$, defined as before by

$$\bar{\omega}(\psi, \phi) : A \xrightarrow{\psi} (A_{PL})_q \otimes B \xrightarrow{1 \cdot \phi} (A_{PL})_q,$$

it is also easy to see the commutativity of:

$$\begin{array}{ccc} \mathcal{F}(A, A_{PL}(K)) \times \langle A_{PL}(K) \rangle & \xleftarrow[\cong]{\beta_* \times \langle \beta \rangle} & \mathcal{F}(A, B) \times \langle B \rangle \\ \downarrow \bar{\omega} & & \downarrow \bar{\omega} \\ \langle A \rangle & \xlongequal{\hspace{10em}} & \langle A \rangle. \end{array} \quad (3)$$

Finally, consider the isomorphism of CDGA's

$$\rho: (\Lambda(V \otimes B_*), \tilde{d}) \xrightarrow{\cong} (\Lambda(A \otimes B_*)/I, d),$$

recalled in §2, and the simplicial isomorphism [3, Thm.3.1]

$$\Psi: \langle \Lambda(A \otimes B_*)/I \rangle \xrightarrow{\cong} \mathcal{F}(A, B),$$

defined as follows: for each $\eta \in \langle \Lambda(A \otimes B_*)/I \rangle_q$ and each $v \in V$,

$$\Psi(\eta)(v) = \sum_i (-1)^{\alpha|b_i|} \eta[v \otimes \beta_i] \otimes b_i.$$

On the other hand observe that

$$\langle \Lambda(A \otimes B_*)/I \otimes B \rangle \cong \langle \Lambda(A \otimes B_*)/I \rangle \times \langle B \rangle.$$

Then, under this identification, we may show that the following commutes:

$$\begin{array}{ccc} \langle \Lambda(A \otimes B_*)/I \rangle \times \langle B \rangle & \xrightarrow{\Psi \times 1_{\langle B \rangle}} & \mathcal{F}(A, B) \times \langle B \rangle \\ \downarrow \langle (\rho \otimes 1_B) \circ \omega \rangle & & \downarrow \bar{\omega} \\ \langle A \rangle & \xlongequal{\hspace{10em}} & \langle A \rangle. \end{array} \quad (4)$$

Indeed, given each $\eta \in \langle \Lambda(A \otimes B_*)/I \rangle_q$, $\phi \in \langle B \rangle$ and $v \in V$:

$$\begin{aligned} (\bar{\omega} \circ \Psi \times 1_{\langle B \rangle})(\eta, \phi)(v) &= \bar{\omega}(\Psi(\eta), \phi)(v) = \sum_i (-1)^{\alpha|b_i|} \eta[v \otimes \beta_i] \cdot \phi(b_i) \\ &= \langle (\rho \otimes 1_B) \circ \omega \rangle(\eta, \phi)(v). \end{aligned}$$

To finish, use diagrams (1), (2), (3) and (4), together with proposition 5, to obtain the following diagram which completes the proof:

$$\begin{array}{ccc}
\mathcal{F}(X_{\mathbb{Q}}, Y_{\mathbb{Q}}) \times X_{\mathbb{Q}} & \xrightarrow{\omega} & Y_{\mathbb{Q}} & \text{Top} \\
& & \vdots \downarrow S_* & \\
S_* \mathcal{F}(X_{\mathbb{Q}}, Y_{\mathbb{Q}}) \times S_* X_{\mathbb{Q}} & \xrightarrow{S_*(\omega)} & S_*(Y_{\mathbb{Q}}) & \\
\uparrow \simeq & & \uparrow \simeq & \\
\mathcal{F}(K, L) \times K & \xrightarrow{\omega} & L & \\
h_* \times 1_K \downarrow \simeq & & \downarrow \simeq h & \\
\mathcal{F}(K, \langle A \rangle) \times K & \xrightarrow{\omega} & \langle A \rangle & \\
\uparrow \simeq & & \parallel & \\
\mathcal{F}(A, A_{PL}(K)) \times \langle A_{PL}(K) \rangle & \xrightarrow{\bar{\omega}} & \langle A \rangle & \text{SimpSet} \\
\uparrow \simeq & & \parallel & \\
\mathcal{F}(A, B) \times \langle B \rangle & \xrightarrow{\bar{\omega}} & \langle A \rangle & \\
\uparrow \cong & & \parallel & \\
\langle \Lambda(A \otimes B_*)/I \rangle \times \langle B \rangle & \xrightarrow{\langle (\rho \otimes 1_B) \circ \omega \rangle} & \langle A \rangle & \\
\downarrow \cong & & \parallel & \\
\langle \Lambda(V \otimes B_*) \rangle \times \langle B \rangle & \xrightarrow{\langle \omega \rangle} & \langle \Lambda V \rangle & \\
& & \uparrow \langle \cdot \rangle & \\
\Lambda(V \otimes B_*) \otimes B & \xleftarrow{\omega} & \Lambda V & \text{AGDC}
\end{array}$$

□

Next, fix a map $f: X \rightarrow Y$ between nilpotent complexes of finite type over \mathbb{Q} and let Z be a finite nilpotent complex. Let $A = (\Lambda W, d) \xrightarrow[\simeq]{\varphi} A_{PL}(X)$ and $B = (\Lambda V, d) \xrightarrow[\simeq]{\psi} A_{PL}(Y)$ be Sullivan models (again non necessar-

ily minimal!) of X and Y respectively, let $C = (C, \delta) \xrightarrow[\cong]{\nu} A_{PL}(Z)$ be a quasi-isomorphism with C finite dimensional connected of finite type, and let $\zeta: (\Lambda V, d) \longrightarrow (\Lambda W, d)$ be a Sullivan model for f . Define

$$\xi: (\Lambda(V \otimes C_*), \tilde{d}) \longrightarrow (\Lambda(W \otimes C_*), \tilde{d}), \quad \xi(v \otimes c) = \rho^{-1}[\zeta(v) \otimes c],$$

being $(\Lambda(V \otimes C_*), \tilde{d})$ and $(\Lambda(W \otimes C_*), \tilde{d})$ the models of $\mathcal{F}(Z, Y_{\mathbb{Q}})$ and $\mathcal{F}(Z, X_{\mathbb{Q}})$ respectively, and $\rho: (\Lambda(W \otimes C_*), \tilde{d}) \xrightarrow{\cong} (\Lambda(A \otimes C_*), d)/I$ the CDGA isomorphism described in §2. In other words, to compute effectively $\xi(v \otimes c)$ use the relations which define I to express $\zeta(v) \otimes c$ as an element of $\Lambda(V \otimes C_*)$. For instance, if $\zeta(v) = w_1 w_2$ and $\Delta c = \sum_i c'_i \otimes c''_i$, $\xi(v \otimes c) = \sum_i (-1)^{|w_2||c'_i|} (w_1 \otimes c'_i)(w_2 \otimes c''_i)$.

Then, we prove this more explicit version of Theorem 2:

Theorem 8. *The commutative square*

$$\begin{array}{ccc} (\Lambda(W \otimes C_*) \otimes C, \tilde{d}) & \xleftarrow{\xi \otimes 1_C} & (\Lambda(V \otimes C_*) \otimes C, \tilde{d}) \\ \omega \uparrow & & \uparrow \omega \\ (\Lambda W, d) & \xleftarrow{\zeta} & (\Lambda V, d) \end{array}$$

is a model of

$$\begin{array}{ccc} \mathcal{F}(Z_{\mathbb{Q}}, X_{\mathbb{Q}}) \times Z_{\mathbb{Q}} & \xrightarrow{(f_{\mathbb{Q}})_* \times 1} & \mathcal{F}(Z_{\mathbb{Q}}, Y_{\mathbb{Q}}) \times Z_{\mathbb{Q}} \\ \omega \downarrow & & \downarrow \omega \\ X_{\mathbb{Q}} & \xrightarrow{f_{\mathbb{Q}}} & Y_{\mathbb{Q}}, \end{array}$$

i.e., there exist a commutative cube of simplicial sets as the following in which

the vertical arrows are homotopy equivalences:

$$\begin{array}{ccccc}
S_*(\mathcal{F}(Z_{\mathbb{Q}}, X_{\mathbb{Q}}) \times Z_{\mathbb{Q}}) & \xrightarrow{S_*((f_{\mathbb{Q}})_* \times 1)} & S_*(\mathcal{F}(Z_{\mathbb{Q}}, Y_{\mathbb{Q}}) \times Z_{\mathbb{Q}}) & & \\
\downarrow \simeq & \searrow S_*(\omega) & \downarrow \simeq & \searrow S_*(\omega) & \\
& & S_*(X_{\mathbb{Q}}) & \xrightarrow{S_*(f_{\mathbb{Q}})} & S_*(Y_{\mathbb{Q}}) \\
& & \downarrow \simeq & & \downarrow \simeq \\
\langle \Lambda(W \otimes C_*) \otimes C \rangle & \xrightarrow{\langle \xi \otimes 1_C \rangle} & \langle \Lambda(V \otimes C_*) \otimes C \rangle & & \\
\searrow \langle \omega \rangle & & \searrow \langle \omega \rangle & & \\
& & \langle \Lambda W, d \rangle & \xrightarrow{\langle \zeta \rangle} & \langle \Lambda V, d \rangle.
\end{array}$$

As in Corollary 7, evaluating at the base point we immediately obtain:

Corollary 9. *The commutative square*

$$\begin{array}{ccc}
(\Lambda(W \otimes C_*), \tilde{d}) & \xleftarrow{\xi} & (\Lambda(V \otimes C_*), \tilde{d}) \\
\omega_0 \uparrow & & \uparrow \omega_0 \\
(\Lambda W, d) & \xleftarrow{\zeta} & (\Lambda V, d)
\end{array}$$

is a model of

$$\begin{array}{ccc}
\mathcal{F}(Z, X_{\mathbb{Q}}) & \xrightarrow{(f_{\mathbb{Q}})_*} & \mathcal{F}(Z, Y_{\mathbb{Q}}) \\
\omega_0 \downarrow & & \downarrow \omega_0 \\
X_{\mathbb{Q}} & \xrightarrow{f_{\mathbb{Q}}} & Y_{\mathbb{Q}}.
\end{array}$$

□

Proof. Fix $N = S_*(Z_{\mathbb{Q}})$, $K = S_*(X_{\mathbb{Q}})$, $L = S_*(Y_{\mathbb{Q}})$ and $\lambda = S_*(f_{\mathbb{Q}}): K \longrightarrow L$. Then we have:

Lemma 10. *The following commutes:*

$$\begin{array}{ccc}
S_*\mathcal{F}(|N|, |K|) \times S_*|N| & \xrightarrow{S_*|\lambda|_* \times 1} & S_*\mathcal{F}(|N|, |L|) \times S_*|N| \\
\downarrow S_*(\omega) & \swarrow \alpha \times \psi(N) \cong & \downarrow S_*(\omega) \\
S_*|K| & \xrightarrow{\omega} \mathcal{F}(N, K) \times N \xrightarrow{\lambda_* \times 1} \mathcal{F}(N, L) \times N & \xrightarrow{\alpha \times \psi(N) \cong} \\
\downarrow \cong & \downarrow \omega & \downarrow \omega \\
S_*|K| & \xrightarrow{S_*(|\lambda|)} S_*|L| & \xrightarrow{\lambda} L \\
\downarrow \cong & \downarrow \cong & \downarrow \cong \\
K & \xrightarrow{\lambda} & L
\end{array}$$

Proof. the lateral faces commute by Proposition 5, while the back and bottom faces are trivially commutative. For the front face observe that given $g: N \times \Delta[q] \longrightarrow K$ in $\mathcal{F}(N, K)_q$, $(\lambda \circ \omega)(g, \sigma) = \lambda(g(\sigma, \Delta_q)) = \omega \circ (\lambda_* \times 1)(g, \sigma)$. Finally, to check the commutativity of the top face, choose as before $g \in \mathcal{F}(N, K)_q$ and note that $\alpha(g) \in S_q\mathcal{F}(|N|, |K|)$ is the map $\Delta^q \longrightarrow \mathcal{F}(|N|, |K|)$ associated by the exponential law to

$$\tau: |N| \times \Delta^q \xrightarrow{\cong} |N| \times |\Delta[q]| \xrightarrow{\cong} |N \times \Delta[q]| \xrightarrow{|g|} |K|.$$

Taking into account that $|\lambda_*(g)| = |\lambda \circ g| = |\lambda| \circ |g|$, we observe that $\alpha(\lambda_*(g)) \in S_q\mathcal{F}(|N|, |L|)$ is again associated by the exponential law to

$$\tau \circ |\lambda|: |N| \times \Delta^q \xrightarrow{1 \times \iota} |N| \times |\Delta[q]| \xrightarrow{\bar{\eta}} |N \times \Delta[q]| \xrightarrow{|\lambda| \circ |g|} |L|.$$

Then, $(S_*(|\lambda|_*) \circ \alpha)(g) = |\lambda|_* \circ \alpha(g) = \alpha(\lambda_*(g)) = (\alpha \circ \lambda_*)(g)$. \square

Lemma 11. *The following commutes:*

$$\begin{array}{ccc}
\mathcal{F}(N, K) \times N & \xrightarrow{\lambda_* \times 1} & \mathcal{F}(N, L) \times N \\
\downarrow \omega & \searrow (h_K)_* \times 1 \cong & \downarrow \omega \\
& \mathcal{F}(N, \langle A \rangle) \times N & \xrightarrow{\langle \zeta \rangle_* \times 1} & \mathcal{F}(N, \langle B \rangle) \times N \\
& \downarrow \omega & & \downarrow \omega \\
K & \xrightarrow{\lambda} & L & \\
\downarrow h_K \cong & & \downarrow h_L \cong & \\
\langle A \rangle & \xrightarrow{\langle \zeta \rangle} & \langle B \rangle.
\end{array}$$

Proof. Side faces are square of type (1); commutativity of the back face has been checked in Lemma above while that of the front and bottom face are trivial. For the top face, given $g \in \mathcal{F}(N, K)_q$, $(h_L)_*(\lambda_*(g)) = h_L \circ \lambda \circ g$, while $\langle \zeta \rangle_*((h_K)_*(g)) = \langle \zeta \rangle \circ h_K \circ g$. But these are equal in view of the following commutative diagram in which the composition of the vertical arrows are precisely h_K and h_L :

$$\begin{array}{ccc}
K & \xrightarrow{\lambda} & L \\
\downarrow \cong & & \downarrow \cong \\
\langle A_{PL}(S_*(X)) \rangle & \xrightarrow{\langle A_{PL}(S_*(f)) \rangle} & \langle A_{PL}(S_*(Y)) \rangle \\
\downarrow \langle \varphi \rangle & & \downarrow \langle \psi \rangle \\
\langle A \rangle & \xrightarrow{\langle \zeta \rangle} & \langle B \rangle.
\end{array}$$

□

Lemma 12. *The following commutes:*

$$\begin{array}{ccc}
\mathcal{F}(N, \langle A \rangle) \times N & \xrightarrow{\langle \zeta \rangle_* \times 1} & \mathcal{F}(N, \langle B \rangle) \times N \\
\downarrow \omega_A & \swarrow \gamma_A \times j^{-1} & \downarrow \omega_B \\
& \mathcal{F}(A, A_{PL}(N)) \times \langle A_{PL}(N) \rangle & \xrightarrow{\zeta^* \times 1} & \mathcal{F}(B, A_{PL}(N)) \times \langle A_{PL}(N) \rangle \\
& \downarrow \bar{\omega}_A & & \downarrow \bar{\omega}_B \\
\langle A \rangle & \xrightarrow{\langle \zeta \rangle} & \langle B \rangle \\
& \searrow & \swarrow \\
& \langle A \rangle & \xrightarrow{\langle \zeta \rangle} & \langle B \rangle.
\end{array}$$

Proof. The lateral squares are as in (2). For the front face, given $g \in \mathcal{F}(A, A_{PL}(N))_q$, both $(\bar{\omega}_B \circ \zeta^* \times 1)(g, \phi)$ and $(\langle \zeta \rangle \circ \bar{\omega}_A)(g, \phi)$ are the same composition

$$B \xrightarrow{\zeta} A \xrightarrow{g} (A_{PL})_q \otimes A_{PL}(N) \xrightarrow{1 \cdot \phi} (A_{PL})_q.$$

For the top face recall that $\gamma_A: \mathcal{F}(A, A_{PL}(N)) \rightarrow \mathcal{F}(N, \langle A \rangle)$ assigns to each $g: A \rightarrow A_{PL}(N) \otimes (A_{PL})_q$ the simplicial map $\gamma(g) = \langle A_{PL}(p_1) \otimes A_{PL}(p_2) \otimes 1 \otimes \theta \circ g \rangle \circ j$ in which $j: N \times \Delta[q] \xrightarrow{\zeta} A_{PL}(N \times \Delta[q])$ is induced by adjunction. Taking into account that $\langle \zeta^*(g) \rangle = \langle g \circ \zeta \rangle = \langle \zeta \rangle \circ \langle g \rangle = \langle \zeta \rangle_*(g)$, it is immediate to see that both $(\langle \zeta \rangle_* \circ \gamma_A)(g)$ and $(\gamma_B \circ \zeta^*)(g)$ correspond to the same map $N \times \Delta[q] \xrightarrow{\gamma(g)} \langle A \rangle \xrightarrow{\langle \zeta \rangle} \langle B \rangle$. \square

Lemma 13. *The following commutes:*

$$\begin{array}{ccccc}
\mathcal{F}(A, A_{PL}(N)) \times \langle A_{PL}(N) \rangle & \xrightarrow{\zeta^* \times 1} & \mathcal{F}(B, A_{PL}(N)) \times \langle A_{PL}(N) \rangle & & \\
\downarrow \bar{\omega}_A & \swarrow \nu_* \times \langle \nu \rangle & \downarrow \bar{\omega}_B & \swarrow \nu_* \times \langle \nu \rangle & \\
\mathcal{F}(A, C) \times \langle C \rangle & \xrightarrow{\zeta^* \times 1} & \mathcal{F}(B, C) \times \langle C \rangle & & \\
\downarrow \bar{\omega}_A & \searrow \langle \zeta \rangle & \downarrow \bar{\omega}_B & \searrow \langle \zeta \rangle & \\
\langle A \rangle & \xrightarrow{\langle \zeta \rangle} & \langle B \rangle & & \\
\downarrow & \searrow & \downarrow & \searrow & \\
\langle A \rangle & \xrightarrow{\langle \zeta \rangle} & \langle B \rangle & &
\end{array}$$

Proof. Lateral faces are squares like (3). The front face has been checked in the past Lemma and the rest of the faces are trivially commutative. \square

Lemma 14. *The following commutes:*

$$\begin{array}{ccccc}
\mathcal{F}(A, C) \times \langle C \rangle & \xrightarrow{\zeta^* \times 1} & \mathcal{F}(B, C) \times \langle C \rangle & & \\
\downarrow \bar{\omega} & \swarrow \Psi_A \times 1 & \downarrow \bar{\omega} & \swarrow \Psi_B \times 1 & \\
\langle \Lambda A \otimes C_* / I \rangle \times \langle C \rangle & \xrightarrow{\langle [\Lambda(\zeta \otimes 1)] \rangle \times 1} & \langle \Lambda B \otimes C_* / I \rangle \times \langle C \rangle & & \\
\downarrow \langle \rho \otimes 1_C \circ \omega \rangle & \searrow \langle \zeta \rangle & \downarrow \langle \rho \otimes 1_C \circ \omega \rangle & \searrow \langle \zeta \rangle & \\
\langle A \rangle & \xrightarrow{\langle \zeta \rangle} & \langle B \rangle & & \\
\downarrow & \searrow & \downarrow & \searrow & \\
\langle A \rangle & \xrightarrow{\langle \zeta \rangle} & \langle B \rangle & &
\end{array}$$

Proof. First, observe that the morphism $\Lambda(\zeta \otimes 1): \Lambda(B \otimes C_*) \rightarrow \Lambda(A \otimes C_*)$ sends I_B into I_A and therefore it induces $[\Lambda(\zeta \otimes 1)]: \Lambda(B \otimes C_*)/I_B \rightarrow \Lambda(A \otimes C_*)/I_A$. Now, the back face has been checked in the past Lemma while lateral faces are squares like (4). For the top face observe that given $g \in \langle \Lambda(A \otimes C_*)/I_A \rangle_q$ and $b \in B$, $\zeta^*(\Psi_A(g))(b) = (\Psi_A(g) \circ \zeta)(b) = \sum_i (-1)^{\alpha(c_i)} g([\zeta b \otimes c_i^*]) \otimes c_i$.

On the other hand,

$$\begin{aligned}
\Psi_B(\langle [\Lambda(\zeta \otimes 1)] \rangle(g))(b) &= \sum_i (-1)^{\alpha(|c_i|)} \langle [\Lambda(\zeta \otimes 1)] \rangle(g)([b \otimes c_i^*]) \otimes c_i \\
&= \sum_i (-1)^{\alpha(|c_i|)} (g \circ [\Lambda(\zeta \otimes 1)])([b \otimes c_i^*]) \otimes c_i \\
&= \sum_i (-1)^{\alpha(|c_i|)} g([\zeta b \otimes c_i^*]) \otimes c_i.
\end{aligned}$$

Finally, the square

$$\begin{array}{ccc}
\Lambda(A \otimes C_*)/I_A \otimes C & \xleftarrow{[\Lambda(\zeta \otimes 1)] \otimes 1_C} & \Lambda(B \otimes C_*)/I_B \otimes C \\
\rho \otimes 1_C \circ \omega \uparrow & & \uparrow \rho \otimes 1_C \circ \omega \\
A & \xleftarrow{\zeta} & B,
\end{array}$$

is trivially commutative, so is the front face. \square

We shall also need an additional cube whose commutativity is trivial:

Lemma 15. *The following commutes:*

$$\begin{array}{ccccc}
\langle \Lambda A \otimes C_*/I \otimes C \rangle & \xrightarrow{\langle [\Lambda(\zeta \otimes 1)] \otimes 1_C \rangle} & \langle \Lambda B \otimes C_*/I \otimes C \rangle & & \\
\downarrow \langle \rho \otimes 1_C \rangle & \searrow \langle \rho \otimes 1_C \rangle & \downarrow \langle \rho \otimes 1_C \otimes \omega \rangle & \searrow \langle \rho \otimes 1_C \rangle & \\
\langle \Lambda(W \otimes C_*) \otimes C \rangle & \xrightarrow{\langle \xi \otimes 1_C \rangle} & \langle \Lambda(V \otimes C_*) \otimes C \rangle & & \\
\downarrow \langle \omega \rangle & & \downarrow \langle \omega \rangle & & \\
\langle A \rangle & \xrightarrow{\langle \zeta \rangle} & \langle B \rangle & & \\
\downarrow & & \downarrow & & \\
\langle \Lambda W \rangle & \xrightarrow{\langle \zeta \rangle} & \langle \Lambda V \rangle & &
\end{array}$$

\square

To finish the proof of Theorem 8 join the cubes of past lemmas by the back and front face to obtain the required commutative diagram. \square

4 The restriction to components

Here we restrict the results of past sections to the components of function spaces to get, in particular, Theorem 3 of the introduction. For this we need some algebraic tools: let $(\Lambda W, d)$ be a CDGA in which W is \mathbb{Z} -graded, and let $u: \Lambda W \longrightarrow \mathbb{Q}$ be an augmentation. In $(\Lambda W, d)$ we consider as in [3] the differential ideal K_u generated by $A_1 \cup A_2 \cup A_3$, where $A_1 = (\Lambda W)^{<0}$, $A_2 = d(\Lambda W)^0$ and $A_3 = \{\alpha - u(\alpha) : \alpha \in (\Lambda W)^0\}$.

Lemma 16. *The ideal K_u coincides with K_u' generated by $A_1' \cup A_2' \cup A_3'$, where $A_1' = W^{<0}$, $A_2' = dW^0$ and $A_3' = \{w - u(w) : w \in W^0\}$.*

Proof. The inclusion $K_u' \subset K_u$ is trivial, as it is $A_1 \subset K_u'$.

Let $\Phi = \alpha - u(\alpha) \in A_3$ and write $\alpha = a + b$, $a \in \Lambda^+(W^{<0}) \cdot \Lambda W$, $b \in \Lambda W^0$. Then, $\alpha - u(\alpha) = a + b - u(a) - u(b) = a + b - u(b)$. As $a \in A_1 \subset K_u'$, it remains to see that $b - u(b) \in K_u'$. Assume $b \in \Lambda^n W^0$ and argue by induction on n . For $n = 1$, trivially $b - u(b) \in A_3' \subset K_u'$. Let $b = b_1 \cdots b_n$, $b_i \in W^0$. Then, $b - u(b) = b_1((b_2 \cdots b_n) - u(b_2 \cdots b_n)) + u(b_2 \cdots b_n)(b_1 - u(b_1)) \in K_u'$ by induction hypothesis.

Finally, let $\Phi = d\alpha$, $\alpha \in (\Lambda W)^0$, be a generator of A_2 and write $\alpha = a + b$ where $a \in (\Lambda^+ W^{<-1}) \cdot \Lambda W$, $b \in \Lambda(W^0)$ and $c \in \Lambda(W^0) \cdot W^{-1} \cdot W^1$. Obviously $da, db \in K_u'$. Write $c = \sum \Phi c_1 c_2$, $\Phi \in \Lambda(W^0)$, $|c_1| = -1$ y $|c_2| = 1$. Thus, $dc = \sum d\Phi c_1 c_2 + \Phi dc_1 c_2 - \Phi c_1 dc_2$. While the first and third set of summands are trivially in K_u' , $u(dc_1) = du(c_1) = 0$ and therefore $\phi dc_1 c_2 = \phi(dc_1 - u(dc_1))c_2$ is also in K_u' . \square

Next, we see that $(\Lambda W, d)/K_u$ is itself a free commutative graded algebra. Let \widetilde{K}_u be the ideal of $(\Lambda W, d)$ generated by $A_1 \cup A_3$ and observe that the projection $\rho: W^1 \xrightarrow{\cong} (\Lambda W/\widetilde{K}_u)^1$ is a vector space isomorphism. Consider the linear map

$$\partial: W^0 \xrightarrow{d} (\Lambda W)^1 \longrightarrow (\Lambda W/\widetilde{K}_u)^1 \xrightarrow{\rho^{-1}} W^1$$

and call \overline{W}^1 a complement of the image of this map, $W^1 = \partial W^0 \oplus \overline{W}^1$.

In which follows, given $\Phi = \alpha \cdot \Psi$, $\alpha \in (\Lambda^+ W^0)$ and $\Psi \in \Lambda W$, we denote by Φ/u the element $u(\alpha)\Psi$. Hence, if for $w \in W^0$, $dw = \Phi_0 + \Phi_1 + \Phi_2$, with $\Phi_0 \in (\Lambda^+ W^{<0}) \cdot (\Lambda W)$, $\Phi_1 \in (\Lambda^+ W^0) \cdot W^1$, $\Phi_2 \in W^1$, then $\partial(w) = \Phi_1/u + \Phi_2$.

Proposition 17. *For a certain \bar{d} , $(\Lambda W/K_u, d) \cong \Lambda(\overline{W}^1 \oplus W^{\geq 2}, \bar{d})$.*

Proof. It is easy to see that the surjective morphism of graded algebras $\varphi: \Lambda W \rightarrow \Lambda \overline{W}^1 \oplus W^{\geq 2}$, given by

$$\varphi(w) = \begin{cases} 0 & \text{if } |w| < 0 \text{ or } w \in \partial W^0, \\ u(w) & \text{if } |w| = 0, \\ w & \text{otherwise,} \end{cases}$$

has K_u as kernel so it induces the required isomorphism. To finish, define $\overline{d} = \overline{\varphi} \circ d \circ \overline{\varphi}^{-1}$. \square

Remark 18. (1) To effectively compute \overline{d} , choose $w \in \Lambda(\overline{W}^1 \oplus W^{\geq 2})$ and write $dw = \Phi_0 + \Phi_1 + \Phi_2 + \Phi_3$, in which $\Phi_0 \in \Lambda^+ W^{<0} \cdot \Lambda W$, $\Phi_1 \in \Lambda^+(\partial W^0) \cdot \Lambda W^{\geq 0}$, $\Phi_2 \in (\Lambda^+ W^0) \cdot (\Lambda \overline{W}^1 \oplus W^{\geq 2})$ and $\Phi_3 \in \Lambda \overline{W}^1 \oplus W^{\geq 2}$. Then, $\overline{d}w = \overline{\varphi} d \overline{\varphi}^{-1} w = \varphi(\Phi_2 + \Phi_3) = \Phi_2/u + \Phi_3$.

(2) Note that if we have in W a basis w_i for which $dw_i \in \Lambda W_{<i}$, then the image of this basis in $\Lambda(\overline{W}^1 \oplus W^{\geq 2}, \overline{d})$ makes it a Sullivan model. However, even when d is decomposable in ΛW , \overline{d} might not be, i.e., $\Lambda(\overline{W}^1 \oplus W^{\geq 2}, \overline{d})$ is not necessarily minimal. This depends on u . In fact, as we just remarked, for each $w \in W$, Φ_2/u could be the linear part of $\overline{d}w$.

Next, consider again the function space $\mathcal{F}(X, Y)$ and recall that, if $h: Y \rightarrow Y_{\mathbb{Q}}$ denotes rationalization, then $\mathcal{F}(X, Y; f) \rightarrow \mathcal{F}(X, Y_{\mathbb{Q}}; h \circ f)$ is also the rationalization map [10, Thm.3.11], i.e., $\mathcal{F}(X, Y; f)_{\mathbb{Q}} = \mathcal{F}(X, Y_{\mathbb{Q}}; h \circ f)$.

On the other hand, recall that for a simplicial set L and a 0-simplex $x_0 \in L_0$, the component of L containing x_0 is the subsimplicial set L_{x_0} in which

$$(L_{x_0})_q = \{x \in L_q \mid \partial^{i_1} \dots \partial^{i_q} x = x_0, \text{ for some } i_1, \dots, i_q\}.$$

Then, given a non necessarily connected CW-complex X , if we denote by $\mathcal{C}(X; x_0)$ the path component of X containing x_0 , there exists a homotopy equivalence $(S_* X)_{x_0} \xrightarrow{\cong} S_* \mathcal{C}(X; x_0)$.

Let $(\Lambda(V \otimes B_*), \tilde{d})$ be the model of $\mathcal{F}(X, Y)$ and let $h \circ f \in \mathcal{F}(X, Y_{\mathbb{Q}})$ considered also as a 0-simplex of $S_0 \mathcal{F}(X, Y_{\mathbb{Q}})$. As before, denote by $S_* \mathcal{F}(X, Y_{\mathbb{Q}})_{hf}$ the component of $S_* \mathcal{F}(X, Y_{\mathbb{Q}})$ containing hf , and by u the 0-simplex of $\langle \Lambda(V \otimes B_*) \rangle$, i.e., the CDGA morphism $u: (\Lambda(V \otimes B_*), \tilde{d}) \rightarrow \mathbb{Q}$, which corresponds to hf through the equivalence $S_* \mathcal{F}(X, Y_{\mathbb{Q}}) \xrightarrow{\cong} \langle \Lambda(V \otimes B_*) \rangle$.

We first prove a generalization of [3, Thm.6.1]:

Proposition 19. *The projection $(\Lambda(V \otimes B_*), \tilde{d}) \rightarrow (\Lambda(V \otimes B_*), \tilde{d})/K_u$ induces a homotopy equivalence*

$$\langle \Lambda(V \otimes B_*)/K_u \rangle \xrightarrow{\cong} \langle \Lambda(V \otimes B_*) \rangle_u$$

which makes the following commutative:

$$\begin{array}{ccc} \langle \Lambda(V \otimes B_*) \rangle_u & \hookrightarrow & \langle \Lambda(V \otimes B_*) \rangle \\ \uparrow \cong & & \parallel \\ \langle \Lambda(V \otimes B_*)/K_u \rangle & \rightarrow & \langle \Lambda(V \otimes B_*) \rangle. \end{array}$$

Proof. We shall need the following version of [9, Thm. 2.2] for $(\Lambda(V \otimes B_*), \tilde{d})$:

Write $B_* = A \oplus \delta A \oplus C$ in which $C \cong H(B_*)$ and choose a basis $\{a_j, b_j, c_k\}$ with $\delta a_j = b_j$ and $\delta c_k = 0$. Also, as ΛV is a Sullivan model choose $\{v_i\}$ a basis for V satisfying $dv_i \in \Lambda(V_{<i})$. Then [3, Prop.5.1], there is a CDGA isomorphism $\eta: (\Lambda(V \otimes B_*), \tilde{d}) \xrightarrow{\cong} \Lambda(U \oplus dU) \otimes (\Lambda W, d)$ in which:

1. U is generated by $u_{ij} = v_i \otimes a_j$.
2. W is generated by $w_{ik} = v_i \otimes c_k + x_{ik}$, for some $x_{ik} \in \Lambda(V_{<i} \otimes B_*)$.
3. If the differential in ΛV is decomposable, the one in ΛW is also decomposable.

Consider now $u: \Lambda(V \otimes B_*) \longrightarrow \mathbb{Q}$ and $u' = u \circ \eta^{-1}$, and let K_u and $K_{u'}$ be the associated differential ideals of $(\Lambda(V \otimes B_*), \tilde{d})$ and $(\Lambda W, d) \otimes \Lambda(U \oplus dU)$ respectively. With the aid of Lemma 16 is easy to see that $\eta(K_u) = K_{u'}$.

On the other hand, consider in $(\Lambda W, d)$ and $\Lambda(U \oplus dU)$ the ideals K_p and K_q associated to the augmentations $p: (\Lambda W, d) \rightarrow (\Lambda W, d) \otimes \Lambda(U \oplus dU) \xrightarrow{u'} \mathbb{Q}$ and $q: \Lambda(U \oplus dU) \rightarrow (\Lambda W, d) \otimes \Lambda(U \oplus dU) \xrightarrow{u'} \mathbb{Q}$ respectively. Again, it is immediate to check that $K_{u'} = K_p \otimes K_q$, and thus,

$$\begin{aligned} (\Lambda(V \otimes B_*), \tilde{d})/K_u &\cong (\Lambda W, d) \otimes \Lambda(U \oplus dU)/K_{u'} \\ &\cong (\Lambda W, d)/K_p \otimes \Lambda(U \oplus dU)/K_q. \end{aligned}$$

Next, observe that $\Lambda(U \oplus dU)/K_q \cong \Lambda(U^{\geq 1} \oplus d(U^{\geq 1}))$, and therefore, it is acyclic. Indeed, the surjective morphism $\varphi: \Lambda(U \oplus dU) \rightarrow \Lambda(U^{\geq 1} \oplus d(U^{\geq 1}))$,

$$\varphi(z) = \begin{cases} 0 & \text{if } z \in U^{(0)} \text{ or } z \in d(U^{\leq 0}), \\ u(z) & \text{if } z \in U^0, \\ z & \text{if } z \in U^{\geq 1}, \end{cases}$$

has K_q as kernel. Then:

$$\begin{aligned} \langle \langle \Lambda(V \otimes B_*), \tilde{d} \rangle / K_u \rangle &\cong \langle \langle \Lambda W, d \rangle / K_p \otimes \Lambda(U \oplus dU) / K_q \rangle \\ &\cong \langle \langle \Lambda W, d \rangle / K_p \rangle \times \langle \Lambda(U \oplus dU) / K_q \rangle \\ &\simeq \langle \langle \Lambda W, d \rangle / K_p \rangle. \end{aligned}$$

Finally, in [3, Thm.6.1] is proved that the projection $\Lambda W \rightarrow \Lambda W / K_p$ induces an equivalence $\langle \Lambda W / K_p \rangle \xrightarrow{\simeq} \langle \Lambda W \rangle_p$. Therefore, as $\langle \Lambda(V \otimes B_*) \rangle_u \simeq \langle \Lambda W \rangle_p$ we get an equivalence $\langle \Lambda(V \otimes B_*) / K_u \rangle \xrightarrow{\simeq} \langle \Lambda(V \otimes B_*) \rangle_u$ which satisfies the required property. \square

Then, we have:

Theorem 20. *The projection $(\Lambda(V \otimes B_*), \tilde{d}) \rightarrow (\Lambda(V \otimes B_*), \tilde{d}) / K_u$ is a model of $i : \mathcal{F}(X, Y; f) \hookrightarrow \mathcal{F}(X, Y)$.*

Proof. Indeed, we have the following diagram

$$\begin{array}{ccc} \mathcal{F}(X, Y_{\mathbb{Q}}; h \circ f) & \xrightarrow{i} & \mathcal{F}(X, Y_{\mathbb{Q}}) & \text{Top} \\ & & \downarrow S_* & \\ S_* \mathcal{F}(X, Y_{\mathbb{Q}}; h \circ f) & \xrightarrow{S_* i} & S_* \mathcal{F}(X, Y_{\mathbb{Q}}) & \\ \simeq \uparrow & & \parallel & \\ S_* \mathcal{F}(X, Y_{\mathbb{Q}})_{hf} & \hookrightarrow & S_* \mathcal{F}(X, Y_{\mathbb{Q}}) & \\ \simeq \downarrow & & \downarrow \simeq & \text{SimpSet} \\ \langle \Lambda(V \otimes B_*) \rangle_u & \hookrightarrow & \langle \Lambda(V \otimes B_*) \rangle & \\ \uparrow \simeq & & \parallel & \\ \langle \Lambda(V \otimes B_*) / K_u \rangle & \longrightarrow & \langle \Lambda(V \otimes B_*) \rangle & \\ & & \uparrow \langle \cdot \rangle & \\ & & \Lambda(V \otimes B_*) & \text{AGDC} \\ & \longleftarrow & \Lambda(V \otimes B_*) / K_u & \end{array}$$

in which the top and middle square are trivially commutative and the bottom is given in Proposition 19. \square

The next results prove Theorem 3 and Corollary 4.

Theorem 21. *The CDGA morphism*

$$\omega_0: (\Lambda V, d) \longrightarrow (\Lambda(V \otimes B_*), \tilde{d})/K_u, \quad \omega(v) = [v \otimes 1^*],$$

is a model of $\omega_0: \mathcal{F}(X, Y; f) \longrightarrow Y$.

Proof. Write ω_0 as the composition

$$\mathcal{F}(X, Y; f) \xrightarrow{i} \mathcal{F}(X, Y) \xrightarrow{\omega_0} Y$$

and apply Theorems 6 y 20. □

In particular, via Proposition 17, we have:

Corollary 22. *The CDGA morphism*

$$\omega_0: (\Lambda V, d) \longrightarrow (\Lambda S_u, \bar{d}) = (\Lambda \overline{V \otimes B_*^1} \oplus (V \otimes B_*)^{\geq 2}, \bar{d}),$$

$\omega_0(v) = v \otimes 1^*$ if $v \in V^{\geq 2}$, or its projection over $\overline{V \otimes B_*^1}$ if $v \in V^1$, is a Sullivan model of the evaluation at the base point $\omega_0: \mathcal{F}(X, Y; f) \rightarrow Y$. □

Observe that, while $\omega_0(v)$ could vanish if $|v| = 1$, when $(\Lambda V, d)$ is 1-connected,

$$\omega_0: (\Lambda V, d) \longrightarrow (\Lambda S_u, \bar{d}) = (\Lambda \overline{V \otimes B_*^1} \oplus (V \otimes B_*)^{\geq 2}, \bar{d}),$$

is a KS-extension or a relative Sullivan algebra. The fibre is of the form

$$(\Lambda(\overline{V \otimes B_*^1} \oplus (V \otimes B_*)^{\geq 2})/(V \otimes 1^*), \bar{d}) = (\Lambda(S_u/V), \bar{d}).$$

Hence, as the fibre of $\omega_0: \mathcal{F}(X, Y; f) \rightarrow Y$ is $\mathcal{F}^*(X, Y; f)$, the path component of f of the space of pointed maps, we immediately obtain:

Corollary 23. *For a 1-connected space Y , $(\Lambda(S_u/V), \bar{d})$ is a Sullivan model of $\mathcal{F}^*(X, Y; f)$.*

Finally, let X, Y, Z and $f: X \rightarrow Y$ be as in Theorem 8. Let $u_g: \Lambda(W \otimes C_*) \rightarrow \mathbb{Q}$ and $u_{f \circ g}: \Lambda(V \otimes C_*) \rightarrow \mathbb{Q}$ be the 0-simplices of $\langle \Lambda(W \otimes C_*) \rangle$ and $\langle \Lambda(V \otimes C_*) \rangle$ corresponding to $h \circ g: Z \rightarrow X_{\mathbb{Q}}$ and $h \circ f \circ g = f_{\mathbb{Q}} \circ h \circ g: Z \rightarrow Y_{\mathbb{Q}}$ respectively.

Theorem 24. *The square*

$$\begin{array}{ccc}
(\Lambda S_{u_g}, \bar{d}) & \xleftarrow{\bar{\xi}} & (\Lambda S_{u_{f \circ g}}, \bar{d}) \\
\omega_0 \uparrow & & \uparrow \omega_0 \\
(\Lambda W, d) & \xleftarrow{\zeta} & (\Lambda V, d)
\end{array}$$

is a Sullivan model of

$$\begin{array}{ccc}
\mathcal{F}(Z, X; g) & \xrightarrow{(f)_*} & \mathcal{F}(Z, Y; f \circ g) \\
\omega_0 \downarrow & & \downarrow \omega_0 \\
X & \xrightarrow{f} & Y,
\end{array}$$

Proof. First observe that the $\Lambda(W \otimes C_*)/K_{u_g} \xleftarrow{\bar{\xi}} \Lambda(V \otimes C_*)/K_{u_{f \circ g}}$ is well defined, i.e., $\xi(K_{u_{f \circ g}}) \subset K_{u_g}$. For that note that $u_{f \circ g}$ is the composition $u_g \circ \xi$. If A_1', A_2' y A_3' is the set of generators of $K_{u_{f \circ g}}$ given in Lemma 16, then $\xi(A_1') \subset K_{u_g}$ trivially. Let $\tilde{d}(w \otimes c)$, $w \otimes c \in (W \otimes C_*)^0$, an element of A_2' . Then, $\xi(\tilde{d}(w \otimes c)) = \tilde{d}\xi(w \otimes c) = \tilde{d}\rho^{-1}[\zeta w \otimes c]$. Write $\rho^{-1}[\zeta w \otimes c] = a + b$, $a \in \Lambda^+(V \otimes C_*)^{(0)} \cdot \Lambda(V \otimes C_*)$, $b \in \Lambda(V \otimes C_*)^0$. Hence, $\tilde{d}(a), \tilde{d}(b) \in K_{u_g}$. Finally consider $w \otimes c - u_{f \circ g}(w \otimes c)$, $w \otimes c \in (W \otimes C_*)^0$, a generator of A_3' . Then,

$$\xi(w \otimes c - u_{f \circ g}(w \otimes c)) = \xi(w \otimes c) - u_{f \circ g}(w \otimes c) = \xi(w \otimes c) - u_g(\xi(w \otimes c)),$$

which again, by Lemma 16, lives in K_{u_g} . To finish the proof we have to show the existence of a commutative cube in **SimpSet** as the following, in which the vertical arrows are homotopy equivalences:

$$\begin{array}{ccccc}
S_*\mathcal{F}(Z, X_{\mathbb{Q}}; h \circ g) & \xrightarrow{S_*(f_{\mathbb{Q}})_*} & S_*\mathcal{F}(Z, Y_{\mathbb{Q}}; f_{\mathbb{Q}} \circ h \circ g) & & \\
\downarrow \simeq & \searrow S_*\omega & \downarrow \simeq & \searrow S_*\omega & \\
\langle \Lambda(W \otimes C_*)/K_{u_g} \rangle & \xrightarrow{S_*(f_{\mathbb{Q}})} & \langle \Lambda(V \otimes C_*)/K_{u_{f \circ g}} \rangle & & \\
\downarrow \omega & \downarrow \simeq & \downarrow \omega & & \downarrow \simeq \\
\langle \Lambda W \rangle & \xrightarrow{\langle \bar{\xi} \rangle} & \langle \Lambda V \rangle & & \\
& & \downarrow \omega & & \\
& & \langle \zeta \rangle & &
\end{array}$$

But this is immediate: Lateral faces are obtained by Theorem 21. The commutativity of the back face is an exercise and the rest are trivially commutative. \square

Finally, we describe the behavior at the homotopy group level of the distinguished maps studied in past sections. Let $(\Lambda S, \bar{d}) \cong (\Lambda(V \otimes B_*)/K_u, d)$ be the Sullivan model of the component $\mathcal{F}(X, Y; f)$ given in Corollary 22. Recall that $u: \Lambda(V \otimes B_*) \rightarrow \mathbb{Q}$ is the CDGA morphism corresponding to f and that $(\Lambda S, \bar{d}) = (\Lambda(\overline{V \otimes B_*})^1 \oplus (V \otimes B_*)^{\geq 2}, \bar{d})$. Now, by classical facts on rational homotopy theory of nilpotent spaces, we have:

Theorem 25. (1) $\pi_*\mathcal{F}(X, Y; f)_{\mathbb{Q}}$ is naturally isomorphic to the dual of $H^*(S, Q(\bar{d})) \cong H^*(\overline{V \otimes B_*})^1 \oplus (V \otimes B_*)^{\geq 2}, Q(\bar{d})$, being $S \cong Q(\Lambda S) = \Lambda S / (\Lambda^+ S \cdot \Lambda^+ S)$ the space of indecomposables. Moreover, the morphism $\pi_*(\omega_0): \pi_*\mathcal{F}(X, Y; f)_{\mathbb{Q}} \rightarrow Y_{\mathbb{Q}}$ is dual of

$$H^*(Q(\omega_0)): H^*(V, Q(d)) \rightarrow H^*(\overline{V \otimes B_*})^1 \oplus (V \otimes B_*)^{\geq 2}, Q(\bar{d}),$$

$$H^*(Q(\omega_0))(v) = [v \otimes 1^*].$$

(3) under the conditions and with the notation of Theorems 8 or 24, the

morphism induced in homotopy groups by

$$\begin{array}{ccc}
 \mathcal{F}(Z, X; g)_{\mathbb{Q}} & \xrightarrow{(f_*)_{\mathbb{Q}}} & \mathcal{F}(Z, Y; f \circ g)_{\mathbb{Q}} \\
 \omega_0 \downarrow & & \downarrow \omega_0 \\
 X_{\mathbb{Q}} & \xrightarrow{f_{\mathbb{Q}}} & Y_{\mathbb{Q}},
 \end{array}$$

is dual of

$$\begin{array}{ccc}
 H^*(S_{u_g}, Q(\bar{d})) & \xleftarrow{H^*(Q(\xi))} & H^*(S_{u_{fg}}, Q(\bar{d})) \\
 H^*(Q(\omega_0)) \uparrow & & \uparrow H^*(Q(\omega_0)) \\
 H^*(W, Q(d)) & \xleftarrow{H^*(Q(\zeta))} & H^*(V, Q(d)).
 \end{array}$$

□

Also, taking homology of indecomposables in Corollary 23 we obtain:

Corollary 26. $\pi_* \mathcal{F}^*(X, Y; f)_{\mathbb{Q}}$ is naturally isomorphic to the dual of $H^*(S/V, Q(\bar{d}))$. □

References

- [1] A. K. Bousfield and V. K. A. M. Gugenheim, On PL De Rham theory and rational homotopy type, *Memoirs of the Amer. Math. Soc.*, **179**(8) (1976).
- [2] E. H. Brown and R. H. Szczarba, Continuous cohomology and real homotopy type, *Trans. Amer. Math. Soc.*, **31** (1989), 57–106.
- [3] E. H. Brown and R. H. Szczarba, On the rational homotopy type of function spaces, *Trans. Amer. Math. Soc.*, **349** (1997), 4931–4951.
- [4] Y. Félix, Rational category of the space of sections of a nilpotent bundle, *Comment. Math. Helvetici*, **65** (1990), 615–622.
- [5] Y. Félix, S. Halperin and J.C. Thomas, Differential graded algebras in topology, in “Handbook of Algebraic Topology”, Elsevier, 1995.

- [6] Y. Félix, S. Halperin and J.C. Thomas, *Rational Homotopy Theory*, *G.T.M.* **205**, Springer, 2000.
- [7] P. G. Goerss and J. F. Jardine, *Simplicial Homotopy Theory*, Progress in Mathematics **174**, Birkhäuser Basel-Boston-Berlin, 1999.
- [8] A. Haefliger, Rational homotopy of the space of sections of a nilpotent bundle, *Trans. Amer. Math. Soc.*, **273** (1982), 609–620.
- [9] S. Halperin, Lecture on Minimal Models, *Mémoires de la Société Mathématique de France*, **230** (1983).
- [10] P. Hilton, G. Mislin and J. Roitberg, *Localization of nilpotent groups and spaces*, North Holland Mathematical Studies, **15**, North Holland, 1975.
- [11] K. Kuribayashi, A rational model for the evaluation map, *Preprint*, (2005).
- [12] J. P. May, *Simplicial Objects in Algebraic Topology*, University of Chicago Press, Chicago, 1992.
- [13] J. Milnor, On spaces having the homotopy type of a CW-complex, *Trans. Amer. Math. Soc.*, **342**(2) (1994), 895–915.
- [14] S. Smith, Rational homotopy of the space of self-maps of complexes with finitely many homotopy groups, *Trans. Amer. Math. Soc.*, **90** (1994), 272–280.
- [15] S. Smith, Rational evaluation subgroups, *Math. Zeit.*, **221** (1996), 387–400.
- [16] D. Sullivan, Infinitesimal computations in Topology, *Publ. Math. de l’I.H.E.S.*, **47** (1978), 269–331.
- [17] D. Tanré, Homotopie modérée et tempérée avec les coalgèbres. Applications aux espaces fonctionnels, *Arch. Math.*, **59** (1992), 130–145.
- [18] R. Thom, L’homologie des espaces fonctionnels, Colloque de topologie algébrique, Louvain, (1957), 299–39.

- [19] M. Vigué-Poirrier, Sur l'homotopie rationnelle des espaces fonctionnels,
Manuscripta Math., **56** (1986), 177–191.

Departamento de Álgebra, Geometría y Topología

Universidad de Málaga

Ap. 59, 29080 Málaga

Spain

e-mail addresses: aniceto@agt.cie.uma.es, urtzi@agt.cie.uma.es