

## INSEPARABLE EXTENSIONS OF ALGEBRAS OVER THE STEENROD ALGEBRA WITH APPLICATIONS TO MODULAR INVARIANT THEORY OF FINITE GROUPS II

MARA D. NEUSEL

ABSTRACT. We continue our study of the homological properties of the purely inseparable extensions  $H \hookrightarrow \mathcal{P}^*\sqrt{H}$  of integrally closed unstable Noetherian integral domains over the Steenrod algebra. It turns out that the projective dimension of  $H$  is a lower bound for the projective dimension of  $\mathcal{P}^*\sqrt{H}$ . Furthermore,  $\text{depth}(H) \geq \text{depth}(\mathcal{P}^*\sqrt{H})$ , where  $\text{depth}$  denotes the depth. Moreover, both algebras have the same global dimension. We apply these results to extension  $\mathbb{F}[V_\bullet]^G \hookrightarrow \mathbb{F}[V]^G$  of rings of invariants.

### 1. INTRODUCTION

Let  $H$  be a unstable reduced algebra over the Steenrod algebra of reduced powers  $\mathcal{P}^*$ . We denote the characteristic by  $p$ , and the order of the ground field  $\mathbb{F}$  by  $q$ . Recall that the Steenrod algebra contains an infinite sequence of derivations iteratively defined as

$$\begin{aligned} P^{\Delta_1} &= P^1, \\ P^{\Delta_i} &= P^{\Delta_{i-1}} P^{q^{i-1}} - P^{q^{i-1}} P^{\Delta_{i-1}} \quad \text{for } i \geq 2. \end{aligned}$$

We set

$$P^{\Delta_0}(h) = \deg(h)h \quad \forall h \in H.$$

Note that  $P^{\Delta_0}$  is not an element of the Steenrod algebra.

The algebra  $H$  is called  **$\mathcal{P}^*$ -inseparably closed**, if whenever  $h \in H$  and

$$P^{\Delta_i}(h) = 0 \quad \forall i \geq 0,$$

then there exists an element  $h' \in H$  such that

$$(h')^p = h.$$

The  **$\mathcal{P}^*$ -inseparable closure** of  $H$  is a  $\mathcal{P}^*$ -inseparably closed algebra  $\mathcal{P}^*\sqrt{H}$  containing  $H$  such that the following universal property holds: Whenever we have a  $\mathcal{P}^*$ -inseparably closed algebra  $H'$  containing  $H$  there exists an embedding  $\mathcal{P}^*\sqrt{H} \hookrightarrow H'$ .

---

Received by the editors May 24, 2007.

2000 *Mathematics Subject Classification*. Primary 55S10 Steenrod Algebra, Secondary 13A50 Invariant Theory.

*Key words and phrases*. Inseparable Extensions, Inseparable Closure, Cohen-Macaulay, Projective Dimension, Depth, Steenrod Algebra, Invariant Theory of Finite Groups.

I would like to thank Lucho Avramov, Lars Christensen and Arne Ledet for many good discussions.

In Section 4.1 of [5] an explicit algorithm to construct the inseparable closure is given. We collect known results in the following proposition.

**Proposition 1.1.** *Consider the natural inclusion*

$$\phi : H \hookrightarrow {}^{\mathcal{P}^*}\sqrt{H}$$

*of unstable reduced algebras over the Steenrod algebra. Then the following statements are valid:*

- (1) *H is an integral domain if and only if  ${}^{\mathcal{P}^*}\sqrt{H}$  is an integral domain.*
  - (2)  *$H \hookrightarrow {}^{\mathcal{P}^*}\sqrt{H}$  is an integral extension, and both algebras have the same Krull dimension.*
  - (3) *If H is integrally closed, then so is  ${}^{\mathcal{P}^*}\sqrt{H}$ .*
  - (4) *H is Noetherian if and only if  ${}^{\mathcal{P}^*}\sqrt{H}$  is Noetherian.*
- If in addition H is Noetherian, then*
- (5) *The extension  $\phi$  is finite.*
  - (6)  *$\overline{H}$  is Cohen-Macaulay if and only if  $\overline{{}^{\mathcal{P}^*}\sqrt{H}}$  is Cohen-Macaulay, where  $\overline{-}$  denotes the integral closure of  $-$ .*
  - (7)  *$\overline{H}$  is polynomial if and only if  $\overline{{}^{\mathcal{P}^*}\sqrt{H}}$  is polynomial.*

*Proof.* For (1) - (3) see Proposition 4.2.1 in [5], for (4) see part (2) of Lemma 4.1.3, Lemma 4.2.2, Proposition 4.2.4, and Theorem 6.3.1 loc.cit., for (5) see Proposition 4.2.4 [5] and [12]. Statement (6) was proven in<sup>1</sup> [8]. Statement (7) was conjectured by C. W. Wilkerson around 1980, [12], and proven in [8].  $\square$

In this paper we proceed with the investigation of the similarities of an unstable integrally closed algebra over the Steenrod algebra and its inseparable closure. The proofs of statements (6) and (7) in the above proposition led to the conjecture that H and its inseparable closure  ${}^{\mathcal{P}^*}\sqrt{H}$  share all properties that have homological characterizations, like, e.g., the depth, the projective dimension, the global dimension, and the Gorenstein property. In this generality this is not true. We illustrate this with the following two examples.

**Example 1.2.** Let  $\mathbb{F}$  be a finite field of characteristic two. Consider the  $\mathcal{P}^*$ -purely inseparable extensions

$$\mathbb{F}[x^2, y^2] \hookrightarrow \mathbb{F}[x^2, y^2, y^3, x^2y] \hookrightarrow \mathbb{F}[x^2, y] \hookrightarrow \mathbb{F}[x, y].$$

All four algebras have Krull dimension two. Moreover  $\mathbb{F}[x, y]$  is the  $\mathcal{P}^*$ -inseparable closure of the other three. Their respective fields of fractions are

$$\mathbb{F}(x^2, y^2) \hookrightarrow \mathbb{F}(x^2, y) = \mathbb{F}(x^2, y) \hookrightarrow \mathbb{F}(x, y).$$

Thus all of them are integrally closed except for  $\mathbb{F}[x^2, y^2, y^3, x^2y]$ . Observe that all of them have depth two, except  $\mathbb{F}[x^2, y^2, y^3, x^2y]$  which has depth one. Thus we see that the three integrally closed algebras are isomorphic as ungraded  $\mathbb{F}$ -algebras (even though not as algebras over the Steenrod algebra). However, we note that  $\mathbb{F}[x^2, y^2, y^3, x^2y]$  is not only not isomorphic to  $\mathbb{F}[x^2, y^2]$ , nor is it isomorphic to  $\mathbb{F}[x^2, y]$ , but also they do not have the same depth either.

Here is another example illustrating that we cannot expect good results for algebras that are not integrally closed.

<sup>1</sup>Note that the necessary assumption on H being integrally closed is missing in that reference.

**Example 1.3.** Consider the  $\mathcal{P}^*$ -purely inseparable extension

$$K = \mathbb{F}[x^2, y, xy] \hookrightarrow H = \mathbb{F}[x, y],$$

where  $|\mathbb{F}| = 2$ . Then  $\overline{K} = H$ , and its global dimension is

$$\text{gl} - \dim(H) = 2.$$

However

$$\text{gl} - \dim(K) = \text{proj} - \dim_K(\mathbb{F}) = \infty,$$

where  $\text{proj} - \dim$  denotes the projective dimension.

## 2. AN UNSTABLE ALGEBRA AND ITS INSEPARABLE CLOSURE

We assume from now on that  $H$  is an integral domain.

**Proposition 2.1.** *Let  $H$  be an integrally closed unstable algebra over the Steenrod algebra. Then*

$$\text{gl} - \dim(H) = \text{gl} - \dim({}^{\mathcal{P}^*}\sqrt{H}).$$

*Proof.* The global dimension of  $H$  is finite if and only if  $H$  is a Noetherian polynomial algebra. By Theorem 7.4 in [8] this is equivalent to  ${}^{\mathcal{P}^*}\sqrt{H}$  being Noetherian and polynomial. Thus the global dimensions of  $H$  and its inseparable closure are simultaneously finite and equal to their common Krull dimension by Theorem 6.3.1 in [5].  $\square$

We denote by  $H^{[p]} \subseteq H$  the subalgebra generated by the  $p$ th powers of elements in  $H$ . The classical Frobenius map

$$H \longrightarrow H^{[p]}, \quad h \mapsto h^p$$

provides us with an (ungraded) isomorphism between the two  $\mathbb{F}$ -algebras.

**Proposition 2.2.** *Let  $H$  be an integrally closed Noetherian integral domain. Then the extension  $H^{[p]} \hookrightarrow H$  splits as a modules<sup>2</sup> over  $H^{[p]} \odot \mathcal{P}^*$ .*

*Proof.* Since  $H$  is Noetherian the extension  $H^{[p]} \hookrightarrow H$  is finite. Thus we can pick a set of generators of  $H$  as a module over  $H^{[p]}$ , say  $\mathbf{t}_1, \dots, \mathbf{t}_k$ , and obtain

$$(\star) \quad H = \sum_{i=1}^k H^{[p]} \mathbf{t}_i.$$

By Proposition 5.1 in [7] we can choose the  $\mathbf{t}_i$ 's to be Thom classes, i.e., for all  $j = 1, \dots, k$

$$\sum_{i=1}^j H^{[p]} \mathbf{t}_i / \sum_{i=1}^{j-1} H^{[p]} \mathbf{t}_i = H^{[p]} \mathbf{t}_j / \left( \sum_{i=1}^{j-1} H^{[p]} \mathbf{t}_i \right) \cap H^{[p]} \mathbf{t}_j$$

is isomorphic to a suspension of an unstable cyclic module over  $H^{[p]}$ . Without loss of generality we can assume that  $\mathbf{t}_1 = 1$ . Consider the extension

$$\mathbb{F}\mathbb{F}(H^{[p]}) \hookrightarrow \sum_{i=1}^k \mathbb{F}\mathbb{F}(H^{[p]}) \mathbf{t}_i \hookrightarrow \mathbb{F}\mathbb{F}(H).$$

---

<sup>2</sup>The notation  $H^{[p]} \odot \mathcal{P}^*$ -module means that we are looking at modules over  $H^{[p]}$  that carry a Steenrod algebra action, that is compatible with the Steenrod algebra action of  $H^{[p]}$ .

We claim that  $\mathbf{FF}(\mathbf{H}) = \sum_{i=1}^k \mathbf{FF}(\mathbf{H}^{[p]})\mathbf{t}_i$ . To that end take an element  $\frac{h}{k} \in \mathbf{FF}(\mathbf{H})$  with  $h, k \in \mathbf{H}$ . Then

$$\frac{h}{k} = \frac{1}{k^p} h k^{p-1} = \frac{1}{k^p} \sum_{i=1}^k h_i \mathbf{t}_i = \sum_{i=1}^k \frac{h_i}{k^p} \mathbf{t}_i \in \sum_{i=1}^k \mathbf{FF}(\mathbf{H}^{[p]})\mathbf{t}_i,$$

for suitable  $h_i \in \mathbf{H}^{[p]}$ . Since  $\mathbf{FF}(\mathbf{H})$  is a finite dimensional vector space over  $\mathbf{FF}(\mathbf{H}^{[p]})$  and  $\{\mathbf{t}_1, \dots, \mathbf{t}_k\}$  forms a spanning set we find a basis among it and obtain

$$\mathbf{FF}(\mathbf{H}^{[p]}) \hookrightarrow \bigoplus_{i=1}^l \mathbf{FF}(\mathbf{H}^{[p]})\mathbf{t}_i = \mathbf{FF}(\mathbf{H})$$

for some  $l \leq k$ . By choice of the  $\mathbf{t}_i$ 's we can rewrite this and obtain a direct sum decomposition as  $\mathbf{FF}(\mathbf{H}^{[p]}) \odot \mathcal{P}^*$ -modules as follows

$$\mathbf{FF}(\mathbf{H}) = \mathbf{FF}(\mathbf{H}^{[p]})\mathbf{t}_1 \oplus \left( \bigoplus_{i=2}^l \mathbf{FF}(\mathbf{H}^{[p]})\mathbf{t}_i \right) / \mathbf{FF}(\mathbf{H}^{[p]})\mathbf{t}_1 \cap \bigoplus_{i=2}^l \mathbf{FF}(\mathbf{H}^{[p]})\mathbf{t}_i.$$

We take the unstable part of  $\mathbf{FF}(\mathbf{H})$ . By [4] we have that

$$\mathbf{H} = \mathbf{Un}(\mathbf{FF}(\mathbf{H}))$$

because  $\mathbf{H}$  is assume to be integrally closed. Since  $\mathbf{Un}$  commutes with direct sums (see [9]) we obtain

$$\mathbf{H} = \mathbf{Un}(\mathbf{FF}(\mathbf{H})) = \mathbf{Un}(\mathbf{FF}(\mathbf{H}^{[p]})\mathbf{t}_1) \oplus \mathbf{Un} \left( \bigoplus_{i=2}^l \mathbf{FF}(\mathbf{H}^{[p]})\mathbf{t}_i \right) / \mathbf{FF}(\mathbf{H}^{[p]})\mathbf{t}_1.$$

Since  $\mathbf{t}_1 = 1$  and  $\mathbf{H}^{[p]}$  is integrally closed, we find  $\mathbf{Un}(\mathbf{FF}(\mathbf{H}^{[p]})\mathbf{t}_1) = \mathbf{H}^{[p]}$ . Thus

$$\mathbf{H} = \mathbf{H}^{[p]}\mathbf{t}_1 \oplus \mathbf{Un} \left( \bigoplus_{i=2}^l \mathbf{FF}(\mathbf{H}^{[p]})\mathbf{t}_i \right) / \mathbf{FF}(\mathbf{H}^{[p]})\mathbf{t}_1$$

as desired.  $\square$

In Chapter 4 of [5] an explicite algorithm to construct the inseparable closure is given. We recollect the few steps we need in what follows:

Denote by  $\mathcal{C}(\mathbf{H}) \subseteq \mathbf{H}$  the subalgebra consisting of the so-called  $\mathcal{P}^*$ -constants:

$$\mathbf{H}^{[p]} \subseteq \mathcal{C}(\mathbf{H}) = \{h \in \mathbf{H} \mid P^{\Delta_i}(h) = 0 \ \forall i \geq 0\} \subseteq \mathbf{H}.$$

Let  $\{s_i, i \in I\}$  be a set of generators of  $\mathcal{C}(\mathbf{H})$  as a module over  $\mathbf{H}^{[p]}$ . We adjoin the  $p$ -th roots of the  $s_i$ 's and obtain

$$\mathbf{H}' = \mathbf{H}[\gamma_1, \gamma_2, \dots] / \sqrt{(\gamma_i^p - s_i, i = 1, 2, \dots)}.$$

Set  $\mathbf{H} = \mathbf{H}_0$  and  $\mathbf{H}' = \mathbf{H}_1$ . Then we define  $\mathbf{H}_i = (\mathbf{H}_{i-1})'$  and we obtain an ascending chain of unstable algebras

$$\mathbf{H} = \mathbf{H}_0 \hookrightarrow \mathbf{H}_1 \hookrightarrow \mathbf{H}_2 \hookrightarrow \dots$$

The  $\mathcal{P}^*$ -inseparable closure is then the colimit

$$\mathcal{P}^*\sqrt{\mathbf{H}} = \operatorname{colim}_i \{\mathbf{H}_i\};$$

see Proposition 4.1.5 in [5]. Furthermore, for the corresponding fields of fractions we have the following:

$$\mathbf{FF}(\mathbf{H}_{i+1}) = \mathbf{FF}(\mathbf{H}_i)[\gamma_1, \gamma_2, \dots] / \sqrt{(\gamma_i^p - s_i, i = 1, 2, \dots)};$$

see Proposition 2.4 in [8].

We note that  $\mathcal{C}(\mathbf{H})$  is itself an unstable Noetherian integral domain over the Steenrod algebra, see Lemmata 4.1.1 and 4.1.2 in [5] if  $\mathbf{H}$  is. We need another property of  $\mathcal{C}(\mathbf{H})$ :

**Lemma 2.3.** *If  $\mathbf{H}$  is an integrally closed integral domain then so is  $\mathcal{C}(\mathbf{H})$ .*

*Proof.* Consider the commutative diagram

$$\begin{array}{ccc} \mathbf{FF}(\mathcal{C}(\mathbf{H})) & \hookrightarrow & \mathbf{FF}(\mathbf{H}) \\ \cup & & \cup \\ \mathcal{C}(\mathbf{H}) & \hookrightarrow & \mathbf{H}. \end{array}$$

Let  $\frac{c}{d} \in \mathbf{FF}(\mathcal{C}(\mathbf{H}))$ , with  $c, d \in \mathcal{C}(\mathbf{H})$ , be integral over  $\mathcal{C}(\mathbf{H})$ . Thus  $\frac{c}{d} \in \mathbf{FF}(\mathbf{H})$  is integral over  $\mathbf{H}$ . Since  $\mathbf{H}$  is integrally closed we find that  $\frac{c}{d} = h \in \mathbf{H}$ . Thus  $c = dh$  and we have for all  $i$  that

$$0 = P^{\Delta_i}(c) = P^{\Delta_i}(dh) = P^{\Delta_i}(d)h + dP^{\Delta_i}(h) = dP^{\Delta_i}(h).$$

Since  $\mathbf{H}$  is an integral domain we have  $P^{\Delta_i}(h) = 0$  and thus  $\frac{c}{d} = h \in \mathcal{C}(\mathbf{H})$  as desired.  $\square$

**Lemma 2.4.** *Let  $\mathbf{H}$  be an unstable algebra over the Steenrod algebra. Then*

$$\mathcal{C}(\mathbf{H}) = (\mathbf{H}_1)^{[p]}.$$

*Proof.* By construction the extension  $\mathbf{H} \hookrightarrow \mathbf{H}_1$  is purely inseparable of exponent one. Thus  $(\mathbf{H}_1)^{[p]} \subseteq \mathbf{H}$ , and since this algebra consists of  $\mathcal{P}^*$ -constants we have

$$(\mathbf{H}_1)^{[p]} \subseteq \mathcal{C}(\mathbf{H}).$$

To prove the reverse inclusion note that every element in  $\mathcal{C}(\mathbf{H})$  has a  $p$ th root in  $\mathbf{H}_1$ , thus is contained in  $(\mathbf{H}_1)^{[p]}$ .  $\square$

**Theorem 2.5.** *Let  $\mathbf{H}$  be an integrally closed Noetherian integral domain. Then*

$$\text{proj} - \dim(\mathbf{H}_{i-1}) \leq \text{proj} - \dim(\mathbf{H}_i) \quad \forall i,$$

*where  $\text{proj} - \dim$  denotes the projective dimension.*

*Proof.* Since  $\mathbf{H}_i$  is an integrally closed integral domain whenever  $\mathbf{H}$  is (see Lemma 2.2 in [8]) it is enough to show the statement for  $i = 1$ . We note that the projective dimension of  $\mathbf{H}$  can be calculated by finding the projective dimension as a module over a system of parameters, say  $S$ . Since  $\mathbf{H} \hookrightarrow \mathbf{H}_1$  is finite  $S \subseteq \mathbf{H}_1$  is a Noether normalization as well. Consider the following commutative diagram of  $S$ -module

homomorphisms and exact rows and columns

$$\begin{array}{ccccccc}
& & & 0 & & 0 & \\
& & & \uparrow & & \uparrow & \\
& & 0 & \hookrightarrow & \mathbf{H}/\mathcal{C}(\mathbf{H}) & \longrightarrow & \mathbf{H}/\mathcal{C}(\mathbf{H}) & \longrightarrow & 0 \\
& & \uparrow & & \uparrow & & \uparrow & & \\
0 & \longrightarrow & \mathbf{H}^{[p]} & \hookrightarrow & \mathbf{H} & \longrightarrow & \mathbf{H}/\mathbf{H}^{[p]} & \longrightarrow & 0 \\
& & \parallel & & \uparrow & & \uparrow & & \\
0 & \longrightarrow & \mathbf{H}^{[p]} & \hookrightarrow & \mathcal{C}(\mathbf{H}) & \longrightarrow & \mathcal{C}(\mathbf{H})/\mathbf{H}^{[p]} & \longrightarrow & 0 \\
& & \uparrow & & \uparrow & & \uparrow & & \\
& & 0 & & 0 & & 0 & & 
\end{array}$$

By Lemma 6.3 in [8] the algebras  $\mathbf{H}$  and  $\mathbf{H}^{[p]}$  have the same projective dimension. Set  $\text{proj-dim}(\mathbf{H}) = \text{proj-dim}(\mathbf{H}^{[p]}) = d$  and  $\text{proj-dim}(\mathcal{C}(\mathbf{H})) = t$ . Since  $\mathbf{H}^{[p]} \hookrightarrow \mathbf{H}$  splits by Proposition 2.2, we read off the second exact row that  $\text{proj-dim}(\mathbf{H}/\mathbf{H}^{[p]}) \leq d$ .

We want to show that  $d \leq t$ . Assume to the contrary that  $d > t$ . We proceed by depth chasing: The last row tells us that  $\text{proj-dim}(\mathcal{C}(\mathbf{H})/\mathbf{H}^{[p]}) = d + 1$ . Thus the last column gives that  $\text{proj-dim}(\mathbf{H}/\mathcal{C}(\mathbf{H})) = d + 2$ . However the middle column says  $\text{proj-dim}(\mathbf{H}/\mathcal{C}(\mathbf{H})) = d$ . This is the desired contradiction. Thus we have

$$\text{proj-dim}(\mathbf{H}^{[p]}) = \text{proj-dim}(\mathbf{H}) \leq \text{proj-dim}(\mathcal{C}(\mathbf{H})).$$

To conclude the proof note that  $\mathcal{C}(\mathbf{H}) = (\mathbf{H}_1)^{[p]}$  by Lemma 2.4 and thus

$$\text{proj-dim}(\mathbf{H}) \leq \text{proj-dim}((\mathbf{H}_1)^{[p]}) = \text{proj-dim}(\mathbf{H}_1)$$

as claimed.  $\square$

**Corollary 2.6.** *Let  $\mathbf{H}$  be a Noetherian integrally closed integral domain. Then*

$$\text{proj-dim}(\mathbf{H}) \leq \text{proj-dim}({}^{\mathcal{P}^*}\sqrt{\mathbf{H}}).$$

*Proof.* Since  $\mathbf{H}$  is Noetherian the chain of algebras

$$\mathbf{H} = \mathbf{H}_0 \hookrightarrow \mathbf{H}_1 \hookrightarrow \mathbf{H}_2 \hookrightarrow \cdots \hookrightarrow \mathbf{H}_r = {}^{\mathcal{P}^*}\sqrt{\mathbf{H}}$$

stabilizes at some  $r \in \mathbb{N}$ ; see Theorem 6.3.1 in [5]. Furthermore, if  $\mathbf{H}$  is integrally closed, then so is  $\mathbf{H}_i$  for all  $i$ ; see Proposition 4.2.1 (5) in [5]. Thus the result follows from the preceding by induction on  $r$ .  $\square$

We have the following immediate corollary:

**Corollary 2.7.** *Let  $\mathbf{H}$  be a Noetherian integrally closed unstable integral domain over the Steenrod algebra and let  ${}^{\mathcal{P}^*}\sqrt{\mathbf{H}}$  be its  $\mathcal{P}^*$ -inseparable closure. Then*

$$\text{depth}(\mathbf{H}) \geq \text{depth}(\mathbf{H}_i) \geq \text{depth}({}^{\mathcal{P}^*}\sqrt{\mathbf{H}})$$

for all  $i$ .

*Proof.* Since  $\mathbf{H}$  is Noetherian, the extensions  $\mathbf{H} \hookrightarrow \mathbf{H}_i \hookrightarrow {}^{\mathcal{P}^*}\sqrt{\mathbf{H}}$  is finite. Thus a Noether normalization  $\mathbf{S} \subseteq \mathbf{H}$  of  $\mathbf{H}$  is a Noether normalization for  $\mathbf{H}_i$  and  ${}^{\mathcal{P}^*}\sqrt{\mathbf{H}}$  as well. Thus the statement follows from the Auslander-Buchsbaum formula.  $\square$

*Remark 2.8.* Note that the above result contains as a special case Corollary 2.2 in [6], where the above inequality was proven for Cohen-Macaulay  ${}^{\mathcal{P}^*}\sqrt{\mathbf{H}}$ .

## 3. APPLICATIONS TO MODULAR INVARIANT THEORY

Let  $\rho : G \hookrightarrow \mathrm{GL}(n, \mathbb{F})$  be a faithful representation of a finite group over a finite field  $\mathbb{F}$ . Denote by  $V = \mathbb{F}^n$  the  $n$ -dimensional vector space over  $\mathbb{F}$ , and by  $\mathbb{F}[V]$  the symmetric algebra on the dual  $V^*$ . The representation  $\rho$  induces a linear action of  $G$  on  $\mathbb{F}[V]$ . Denote by  $\mathbb{F}[V]^G \subseteq \mathbb{F}[V]$  the subring of  $G$ -invariant polynomials. By the Galois Embedding Theorem an integrally closed  $\mathcal{P}^*$ -inseparably closed Noetherian unstable integral domain over the Steenrod algebra is such a ring of invariants  $\mathbb{F}[V]^G$  for a suitable representation  $\rho$  of some group  $G$ , see [1] and Theorem 7.1.1 in [5].

Let  $V = W_0 \oplus \cdots \oplus W_e$  be a vector space decomposition. Set

$$\mathbb{F}[V_\bullet] = \mathbb{F}[W_0] \otimes_{\mathbb{F}} \mathbb{F}[W_1]^{[p]} \otimes_{\mathbb{F}} \cdots \otimes_{\mathbb{F}} \mathbb{F}[W_e]^{[p^e]}.$$

The generalized Galois Embedding Theorem states that  $H$  is isomorphic to  $\mathbb{F}[V_\bullet]^G$  as an algebra over the Steenrod algebra for some suitable flag  $V_\bullet$ , group  $G$ , and representation  $\rho$  if and only if  $H$  is an integrally closed Noetherian unstable integral domain over the Steenrod algebra, see [12] and Theorem 5.2 in [8]. Furthermore we have a commutative diagram

$$\begin{array}{ccc} H = \mathbb{F}[V_\bullet]^G & \hookrightarrow & \mathbb{F}[V_\bullet] \\ \downarrow & & \downarrow \\ \mathcal{P}^*\sqrt{H} = \mathbb{F}[V]^G & \hookrightarrow & \mathbb{F}[V] \end{array},$$

where the horizontal inclusions are Galois and the vertical inclusions are purely  $\mathcal{P}^*$ -inseparable.

In Proposition 6.4 and Theorem 7.4 in [8] we saw that  $\mathbb{F}[V_\bullet]^G$  and  $\mathbb{F}[V]^G$  are simultaneously Cohen-Macaulay, or polynomial. Based on the results of the preceding section we can add to that list the following properties.

**Theorem 3.1.** *Let  $\mathbb{F}[V_\bullet]^G \hookrightarrow \mathbb{F}[V]^G$  be an extension of rings of invariants, where  $V_\bullet$  is a  $G$ -flag in  $V$ , and  $\mathbb{F}$  is the prime field of characteristic  $p$ . Then*

- (1)  $\mathbb{F}[V_\bullet]^G$  and  $\mathbb{F}[V]^G$  have the same global dimension,
- (2)  $\mathrm{proj} - \dim(\mathbb{F}[V_\bullet]^G) \leq \mathrm{proj} - \dim(\mathbb{F}[V]^G)$ , and
- (3)  $\mathrm{depth}(\mathbb{F}[V_\bullet]^G) \geq \mathrm{depth}(\mathbb{F}[V]^G)$ .

*Proof.* The first statement follows from Proposition 2.1. The second statement follows from Corollary 2.6 and the last from Corollary 2.7.  $\square$

*Remark 3.2.* We note that the preceding result can be refined for the extension

$$\mathbb{F}[V_\star]^G \hookrightarrow \mathbb{F}[V_\bullet]^G,$$

where  $V_\star \subseteq V_\bullet$  denotes a subflag. We find

- (1)  $\mathbb{F}[V_\bullet]^G$  and  $\mathbb{F}[V_\star]^G$  have the same global dimension by Proposition 2.1,
- (2)  $\mathrm{proj} - \dim(\mathbb{F}[V_\star]^G) \leq \mathrm{proj} - \dim(\mathbb{F}[V_\bullet]^G)$  by Theorem 2.5, since  $\mathbb{F}[V_\bullet]^G = (\mathbb{F}[V_\star]^G)_i$  for some  $i \geq 0$ , and similarly
- (3)  $\mathrm{depth}(\mathbb{F}[V_\star]^G) \geq \mathrm{depth}(\mathbb{F}[V_\bullet]^G)$  by Corollary 2.7.

We note that in the case of the preceding result the image of  $G$  under  $\rho$  necessarily consists of matrices of the form

$$\begin{bmatrix} A_0 & 0 & \cdots & 0 \\ * & A_1 & 0 & \cdots & 0 \\ & * & \ddots & & \vdots \\ \cdots & & \ddots & & 0 \\ * & & \cdots & * & A_e \end{bmatrix}$$

where  $A_i$  is an invertible  $n_i \times n_i$ -matrix with  $n_i = \dim(W_i)$ .

**Proposition 3.3.** *If  $\rho(G)$  consists of matrices of block diagonal form*

$$\begin{bmatrix} A_0 & 0 & \cdots & 0 \\ 0 & A_1 & 0 & \cdots & 0 \\ & 0 & \ddots & & \vdots \\ \cdots & & \ddots & & 0 \\ 0 & & \cdots & 0 & A_e \end{bmatrix},$$

then  $\mathbb{H}$  and  ${}^{\mathcal{P}}\sqrt{\mathbb{H}}$  are ungraded isomorphic.

*Proof.* Consider the (ungraded) isomorphism

$$\phi : \mathbb{F}[V] \longrightarrow \mathbb{F}[V_{\bullet}], x_i \mapsto x_i^{p^j}$$

for  $x_i \in \mathbb{F}[W_j]$  as basis element. Since  $G$  acts on  $\mathbb{F}[W_j]$  for all  $j$ , the map  $\phi$  commutes with the group action. Thus the result follows.  $\square$

*Remark 3.4.* Obviously the preceding result remains true over any field of finite characteristic.

We want to illustrate these results with an example taken from [8]; see Example 7.6 loc.cit.

**Example 3.5.** Let  $p$  be odd, and  $\mathbb{F}$  a field of characteristic  $p$ . Consider the four dimensional modular representation  $\mathbb{Z}/p \hookrightarrow \mathrm{GL}(4, \mathbb{F})$  afforded by the matrix

$$\begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Its ring of invariants turns out to be a hypersurface

$$\mathbb{F}[x_1, y_1, x_2, y_2]^{\mathbb{Z}/p} = \mathbb{F}[c_1, y_1, c_2, y_2, q]/(r),$$

where  $c_i = x_i^p - x_i y_i^{p-1}$  are the top orbit Chern classes of  $x_i$ ,  $i = 1, 2$ , and  $q = x_1 y_2 - x_2 y_1$  is an invariant quadratic form. The relation is given by

$$r = q^p - c_1 y_2^p + c_2 y_1^p + q y_1^{p-1} y_2^{p-1},$$

see Theorem 2.1 in [3]. Certainly,  $\mathbb{Z}/p$  acts also on  $\mathbb{F}[x_1, y_1] \otimes \mathbb{F}[x_2^p, y_2^p]$  and we find that

$$(\mathbb{F}[x_1, y_1] \otimes \mathbb{F}[x_2^p, y_2^p])^{\mathbb{Z}/p} = \mathbb{F}[c_1, y_1, c_2^p, y_2^p, q']/(r'),$$



where  $q' = x_1 y_2^p - x_2^p y_1$  and  $r' = (q')^p - c_1 y_2^{p^2} + c_2^p y_1^p - q' y_1^{p-1} y_2^{p(p-1)}$ . We note that the two rings are isomorphic, but not graded isomorphic, nor (in the case of a finite ground field  $\mathbb{F}$ ) isomorphic as algebras over the Steenrod algebra.

## REFERENCES

1. J. F. Adams and C. W. Wilkerson: Finite  $H$ -Spaces and Algebras over the Steenrod Algebra, *Annals of Mathematics* 111 (1980), 95-143.
2. H. Matsumura: *Commutative Ring Theory*, Cambridge Studies in Advanced Mathematics 8, Cambridge University Press, Cambridge 1986.
3. M. D. Neusel: Invariants of some Abelian  $p$ -Groups in Characteristic  $p$ , *Proc. of the AMS* 125 (1997), 1921-1931.
4. M. D. Neusel: Localizations over the Steenrod Algebra. The lost Chapter, *Mathematische Zeitschrift* 235 (2000), 353-378.
5. M. D. Neusel: *Inverse Invariant Theory and Steenrod Operations*, Memoirs of the AMS 146, AMS, Providence RI 2000.
6. M. D. Neusel: Unstable Cohen-Macaulay Algebras, *Math. Research Letters* 8 (2001), 347-360.
7. M. D. Neusel: The Existence of Thom Classes, *Journal of Pure and Applied Algebra* 191 (2004), 265-283.
8. M. D. Neusel: Inseparable Extensions of Algebras over the Steenrod Algebra with Applications to Modular Invariant Theory of Finite Groups, *Transactions of the AMS* 358 (2006), 4689-4720.
9. M. D. Neusel: On the Unstable Parts Functor, in preparation.
10. C. A. Weibel: *Introduction to Homological Algebra*, Cambridge Studies in Advanced Mathematics 38, Cambridge University Press, Cambridge 1994.
11. C. W. Wilkerson: Integral Closure of Unstable Steenrod Algebra Actions, *Journal of Pure and Applied Algebra* 13 (1978), 49-55.
12. C. W. Wilkerson: Rings of Invariants and Inseparable Forms of Algebras over the Steenrod Algebra, pp. 381-396 in: *Recent progress in homotopy theory (Baltimore, MD, 2001)*, Contemp. Math. 293, AMS, Providence RI 2002

DEPARTMENT OF MATHEMATICS AND STATISTICS, MS 1042, TEXAS TECH UNIVERSITY, LUBBOCK, TEXAS 79409

*E-mail address:* Mara.D.Neusel@ttu.edu